


# Polynomials

## Exercise 2.1


Question 1:

The graphs of  $y = p(x)$  are given in following figure, for some polynomials  $p(x)$ . Find the number of zeroes of  $p(x)$ , in each case.


(i)




(ii)




(iii)




(iv)



(v)



(v)



**Answer:**

(i) The number of zeroes is 0 as the graph does not cut the x-axis at any point.

(ii) The number of zeroes is 1 as the graph intersects the x-axis at only 1 point.

(iii) The number of zeroes is 3 as the graph intersects the x-axis at 3 points.

(iv) The number of zeroes is 2 as the graph intersects the x-axis at 2 points.

(v) The number of zeroes is 4 as the graph intersects the x-axis at 4 points.

(vi) The number of zeroes is 3 as the graph intersects the x-axis at 3 points.

## Exercise 2.2

### Question 1:

Find the zeroes of the following quadratic polynomials and verify the relationship between the zeroes and the coefficients.

$$(i) x^2 - 2x - 8 \quad (ii) 4s^2 - 4s + 1 \quad (iii) 6x^2 - 3 - 7x$$

$$(iv) 4u^2 + 8u \quad (v) t^2 - 15 \quad (vi) 3x^2 - x - 4$$

### Answer:

$$(i) \quad x^2 - 2x - 8 = (x - 4)(x + 2)$$

The value of  $x^2 - 2x - 8$  is zero when  $x - 4 = 0$  or  $x + 2 = 0$ , i.e., when  $x = 4$  or  $x = -2$

Therefore, the zeroes of  $x^2 - 2x - 8$  are 4 and -2.

$$\text{Sum of zeroes} = 4 - 2 = 2 = \frac{-(-2)}{1} = \frac{-(\text{Coefficient of } x)}{\text{Coefficient of } x^2}$$

$$\text{Product of zeroes} = 4 \times (-2) = -8 = \frac{(-8)}{1} = \frac{\text{Constant term}}{\text{Coefficient of } x^2}$$

$$(ii) \quad 4s^2 - 4s + 1 = (2s - 1)^2$$

The value of  $4s^2 - 4s + 1$  is zero when  $2s - 1 = 0$ , i.e.,  $s = \frac{1}{2}$

Therefore, the zeroes of  $4s^2 - 4s + 1$  are  $\frac{1}{2}$  and  $\frac{1}{2}$ .

$$\text{Sum of zeroes} = \frac{1}{2} + \frac{1}{2} = 1 = \frac{-(-4)}{4} = \frac{-(\text{Coefficient of } s)}{(\text{Coefficient of } s^2)}$$

$$\text{Product of zeroes} = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4} = \frac{\text{Constant term}}{\text{Coefficient of } s^2}$$

$$(iii) \quad 6x^2 - 3 - 7x = 6x^2 - 7x - 3 = (3x+1)(2x-3)$$

The value of  $6x^2 - 3 - 7x$  is zero when  $3x + 1 = 0$  or  $2x - 3 = 0$ , i.e.,  $x = \frac{-1}{3}$  or  $x = \frac{3}{2}$

Therefore, the zeroes of  $6x^2 - 3 - 7x$  are  $\frac{-1}{3}$  and  $\frac{3}{2}$ .

$$\text{Sum of zeroes} = \frac{-1}{3} + \frac{3}{2} = \frac{7}{6} = \frac{-(-7)}{6} = \frac{-(\text{Coefficient of } x)}{\text{Coefficient of } x^2}$$

$$\text{Product of zeroes} = \frac{-1}{3} \times \frac{3}{2} = \frac{-1}{2} = \frac{-3}{6} = \frac{\text{Constant term}}{\text{Coefficient of } x^2}$$

$$(iv) \quad 4u^2 + 8u = 4u^2 + 8u + 0 \\ = 4u(u + 2)$$

The value of  $4u^2 + 8u$  is zero when  $4u = 0$  or  $u + 2 = 0$ , i.e.,  $u = 0$  or  $u = -2$

Therefore, the zeroes of  $4u^2 + 8u$  are 0 and -2.

$$\text{Sum of zeroes} = 0 + (-2) = -2 = \frac{-(8)}{4} = \frac{-(\text{Coefficient of } u)}{\text{Coefficient of } u^2}$$

$$\text{Product of zeroes} = 0 \times (-2) = 0 = \frac{0}{4} = \frac{\text{Constant term}}{\text{Coefficient of } u^2}$$

$$(v) \quad t^2 - 15 \\ = t^2 - 0t - 15 \\ = (t - \sqrt{15})(t + \sqrt{15})$$

The value of  $t^2 - 15$  is zero when  $t - \sqrt{15} = 0$  or  $t + \sqrt{15} = 0$ , i.e., when  $t = \sqrt{15}$  or  $t = -\sqrt{15}$

Therefore, the zeroes of  $t^2 - 15$  are  $\sqrt{15}$  and  $-\sqrt{15}$ .

$$\text{Sum of zeroes} = \sqrt{15} + (-\sqrt{15}) = 0 = \frac{-0}{1} = \frac{-(\text{Coefficient of } t)}{(\text{Coefficient of } t^2)}$$

$$\text{Product of zeroes} = (\sqrt{15})(-\sqrt{15}) = -15 = \frac{-15}{1} = \frac{\text{Constant term}}{\text{Coefficient of } x^2}$$

$$\begin{aligned} \text{(vi)} \quad 3x^2 - x - 4 \\ = (3x - 4)(x + 1) \end{aligned}$$

The value of  $3x^2 - x - 4$  is zero when  $3x - 4 = 0$  or  $x + 1 = 0$ , i.e., when  $x = \frac{4}{3}$  or  $x = -1$

Therefore, the zeroes of  $3x^2 - x - 4$  are  $\frac{4}{3}$  and  $-1$ .

$$\text{Sum of zeroes} = \frac{4}{3} + (-1) = \frac{1}{3} = \frac{-(-1)}{3} = \frac{-(\text{Coefficient of } x)}{\text{Coefficient of } x^2}$$

$$\text{Product of zeroes} = \frac{4}{3}(-1) = \frac{-4}{3} = \frac{\text{Constant term}}{\text{Coefficient of } x^2}$$

### Question 2:

**Find a quadratic polynomial each with the given numbers as the sum and product of its zeroes respectively.**

$$\text{(i)} \quad \frac{1}{4}, -1 \quad \text{(ii)} \quad \sqrt{2}, \frac{1}{3} \quad \text{(iii)} \quad 0, \sqrt{5}$$

$$\text{(iv)} \quad 1, 1 \quad \text{(v)} \quad -\frac{1}{4}, \frac{1}{4} \quad \text{(vi)} \quad 4, 1$$

**Answer:**

(i)  $\frac{1}{4}, -1$

Let the polynomial be  $ax^2 + bx + c$ , and its zeroes be  $\alpha$  and  $\beta$ .

$$\alpha + \beta = \frac{1}{4} = \frac{-b}{a}$$

$$\alpha\beta = -1 = \frac{-4}{4} = \frac{c}{a}$$

If  $a = 4$ , then  $b = -1$ ,  $c = -4$

Therefore, the quadratic polynomial is  $4x^2 - x - 4$ .

(ii)  $\sqrt{2}, \frac{1}{3}$

Let the polynomial be  $ax^2 + bx + c$ , and its zeroes be  $\alpha$  and  $\beta$ .

$$\alpha + \beta = \sqrt{2} = \frac{3\sqrt{2}}{3} = \frac{-b}{a}$$

$$\alpha\beta = \frac{1}{3} = \frac{c}{a}$$

If  $a = 3$ , then  $b = -3\sqrt{2}$ ,  $c = 1$

Therefore, the quadratic polynomial is  $3x^2 - 3\sqrt{2}x + 1$ .

(iii)  $0, \sqrt{5}$

Let the polynomial be  $ax^2 + bx + c$ , and its zeroes be  $\alpha$  and  $\beta$ .

$$\alpha + \beta = 0 = \frac{0}{1} = \frac{-b}{a}$$

$$\alpha \times \beta = \sqrt{5} = \frac{\sqrt{5}}{1} = \frac{c}{a}$$

If  $a = 1$ , then  $b = 0$ ,  $c = \sqrt{5}$

Therefore, the quadratic polynomial is  $x^2 + \sqrt{5}$ .

(iv) 1, 1

Let the polynomial be  $ax^2 + bx + c$ , and its zeroes be  $\alpha$  and  $\beta$ .

$$\alpha + \beta = 1 = \frac{1}{1} = \frac{-b}{a}$$

$$\alpha \times \beta = 1 = \frac{1}{1} = \frac{c}{a}$$

If  $a = 1$ , then  $b = -1$ ,  $c = 1$

Therefore, the quadratic polynomial is  $x^2 - x + 1$ .

(v)  $-\frac{1}{4}, \frac{1}{4}$

Let the polynomial be  $ax^2 + bx + c$ , and its zeroes be  $\alpha$  and  $\beta$ .

$$\alpha + \beta = \frac{-1}{4} = \frac{-b}{a}$$

$$\alpha \times \beta = \frac{1}{4} = \frac{c}{a}$$

If  $a = 4$ , then  $b = 1$ ,  $c = 1$

Therefore, the quadratic polynomial is  $4x^2 + x + 1$ .

(vi) 4, 1

Let the polynomial be  $ax^2 + bx + c$ .

$$\alpha + \beta = 4 = \frac{4}{1} = \frac{-b}{a}$$

$$\alpha \times \beta = 1 = \frac{1}{1} = \frac{c}{a}$$

If  $a = 1$ , then  $b = -4$ ,  $c = 1$

Therefore, the quadratic polynomial is  $x^2 - 4x + 1$ .

## Exercise 2.3

### Question 1:

Divide the polynomial  $p(x)$  by the polynomial  $g(x)$  and find the quotient and remainder in each of the following:

(i)  $p(x) = x^3 - 3x^2 + 5x - 3$ ,  $g(x) = x^2 - 2$

(ii)  $p(x) = x^4 - 3x^2 + 4x + 5$ ,  $g(x) = x^2 + 1 - x$

(iii)  $p(x) = x^4 - 5x + 6$ ,  $g(x) = 2 - x^2$

### Answer:

(i)  $p(x) = x^3 - 3x^2 + 5x - 3$   
 $q(x) = x^2 - 2$

$$\begin{array}{r} x-3 \\ x^2-2 \overline{) x^3-3x^2+5x-3} \\ \underline{x^3 \quad -2x} \phantom{-3} \\ -3x^2+7x-3 \\ \underline{-3x^2 \quad +6} \phantom{-3} \\ + \phantom{-3x^2} -9 \\ \hline 7x-9 \end{array}$$

Quotient =  $x - 3$

Remainder =  $7x - 9$



$$(ii) \quad p(x) = x^4 - 3x^2 + 4x + 5 = x^4 + 0x^3 - 3x^2 + 4x + 5$$

$$q(x) = x^2 + 1 - x = x^2 - x + 1$$

$$\begin{array}{r}
 x^2 - x + 1 \overline{) x^4 + 0x^3 - 3x^2 + 4x + 5} \\
 \underline{x^4 - x^3 + x^2} \phantom{+ 5} \\
 x^3 - 4x^2 + 4x + 5 \\
 \underline{x^3 - x^2 + x} \phantom{+ 5} \\
 -3x^2 + 3x + 5 \\
 \underline{-3x^2 + 3x - 3} \phantom{+ 5} \\
 8
 \end{array}$$

$$\text{Quotient} = x^2 + x - 3$$

$$\text{Remainder} = 8$$

$$(iii) \quad p(x) = x^4 - 5x + 6 = x^4 + 0x^2 - 5x + 6$$

$$q(x) = 2 - x^2 = -x^2 + 2$$

$$\begin{array}{r}
 -x^2 + 2 \overline{) x^4 + 0x^2 - 5x + 6} \\
 \underline{x^4 - 2x^2} \phantom{+ 6} \\
 2x^2 - 5x + 6 \\
 \underline{2x^2 - 4} \phantom{+ 6} \\
 -5x + 10
 \end{array}$$

$$\text{Quotient} = -x^2 - 2$$

$$\text{Remainder} = -5x + 10$$

### Question 2:

Check whether the first polynomial is a factor of the second polynomial by dividing the second polynomial by the first polynomial:

- (i)  $t^2 - 3, 2t^4 + 3t^3 - 2t^2 - 9t - 12$   
(ii)  $x^2 + 3x + 1, 3x^4 + 5x^3 - 7x^2 + 2x + 2$   
(iii)  $x^3 - 3x + 1, x^5 - 4x^3 + x^2 + 3x + 1$

**Answer:**

(i)  $t^2 - 3, 2t^4 + 3t^3 - 2t^2 - 9t - 12$

$$t^2 - 3 = t^2 + 0t - 3$$

$$\begin{array}{r}
 t^2 + 0t - 3 \overline{) 2t^4 + 3t^3 - 2t^2 - 9t - 12} \\
 \underline{2t^4 + 0t^3 - 6t^2} \phantom{- 9t - 12} \\
 \phantom{2t^4 + } 3t^3 + 4t^2 - 9t - 12 \\
 \phantom{2t^4 + } \underline{3t^3 + 0t^2 - 9t} \phantom{- 12} \\
 \phantom{2t^4 + } \phantom{3t^3 + } 4t^2 + 0t - 12 \\
 \phantom{2t^4 + } \phantom{3t^3 + } \underline{4t^2 + 0t - 12} \\
 \phantom{2t^4 + } \phantom{3t^3 + } \phantom{4t^2 + } 0
 \end{array}$$

Since the remainder is 0,

Hence,  $t^2 - 3$  is a factor of  $2t^4 + 3t^3 - 2t^2 - 9t - 12$ .

(ii)  $x^2 + 3x + 1, 3x^4 + 5x^3 - 7x^2 + 2x + 2$

$$\begin{array}{r}
 3x^2 - 4x + 2 \\
 x^2 + 3x + 1 \overline{) 3x^4 + 5x^3 - 7x^2 + 2x + 2} \\
 \underline{3x^4 + 9x^3 + 3x^2} \phantom{+ 2x + 2} \\
 -4x^3 - 10x^2 + 2x + 2 \\
 \underline{-4x^3 - 12x^2 - 4x} \phantom{+ 2} \\
 2x^2 + 6x + 2 \\
 \underline{2x^2 + 6x + 2} \\
 0
 \end{array}$$

Since the remainder is 0,

Hence,  $x^2 + 3x + 1$  is a factor of  $3x^4 + 5x^3 - 7x^2 + 2x + 2$ .

(iii)  $x^3 - 3x + 1$ ,  $x^5 - 4x^3 + x^2 + 3x + 1$

$$\begin{array}{r}
 x^2 - 1 \\
 x^3 - 3x + 1 \overline{) x^5 - 4x^3 + x^2 + 3x + 1} \\
 \underline{x^5 - 3x^3 + x^2} \phantom{+ 3x + 1} \\
 -x^3 \phantom{+ 3x} + 1 \\
 \underline{-x^3 \phantom{+ 3x} - 1} \\
 2x \phantom{+ 1} \\
 \underline{2x} \\
 0
 \end{array}$$

Since the remainder  $\neq 0$ ,

Hence,  $x^3 - 3x + 1$  is not a factor of  $x^5 - 4x^3 + x^2 + 3x + 1$ .

### Question 3:

Obtain all other zeroes of  $3x^4 + 6x^3 - 2x^2 - 10x - 5$ , if two of its zeroes are

$$\sqrt{\frac{5}{3}} \text{ and } -\sqrt{\frac{5}{3}}.$$

**Answer:**

$$p(x) = 3x^4 + 6x^3 - 2x^2 - 10x - 5$$

Since the two zeroes are  $\sqrt{\frac{5}{3}}$  and  $-\sqrt{\frac{5}{3}}$ ,

$\therefore \left(x - \sqrt{\frac{5}{3}}\right)\left(x + \sqrt{\frac{5}{3}}\right) = \left(x^2 - \frac{5}{3}\right)$  is a factor of  $3x^4 + 6x^3 - 2x^2 - 10x - 5$ .

Therefore, we divide the given polynomial by  $x^2 - \frac{5}{3}$ .

$$\begin{array}{r}
 x^2 + 0x - \frac{5}{3} \overline{) 3x^4 + 6x^3 - 2x^2 - 10x - 5} \\
 \underline{3x^4 + 0x^3 - 5x^2} \phantom{- 10x - 5} \\
 6x^3 + 3x^2 - 10x - 5 \\
 \underline{6x^3 + 0x^2 - 10x} \phantom{- 5} \\
 3x^2 + 0x - 5 \\
 \underline{3x^2 + 0x - 5} \\
 0
 \end{array}$$

$$\begin{aligned}
 3x^4 + 6x^3 - 2x^2 - 10x - 5 &= \left(x^2 - \frac{5}{3}\right)(3x^2 + 6x + 3) \\
 &= 3\left(x^2 - \frac{5}{3}\right)(x^2 + 2x + 1)
 \end{aligned}$$

We factorize  $x^2 + 2x + 1$

$$= (x+1)^2$$

Therefore, its zero is given by  $x + 1 = 0$

$$x = -1$$

As it has the term  $(x+1)^2$ , therefore, there will be 2 zeroes at  $x = -1$ .

Hence, the zeroes of the given polynomial are  $\sqrt{\frac{5}{3}}$ ,  $-\sqrt{\frac{5}{3}}$ ,  $-1$  and  $-1$ .

#### Question 4:

On dividing  $x^3 - 3x^2 + x + 2$  by a polynomial  $g(x)$ , the quotient and remainder were  $x - 2$  and  $-2x + 4$ , respectively. Find  $g(x)$ .

**Answer:**

$$p(x) = x^3 - 3x^2 + x + 2 \quad (\text{Dividend})$$

$$g(x) = ? \quad (\text{Divisor})$$

$$\text{Quotient} = (x - 2)$$

$$\text{Remainder} = (-2x + 4)$$

$$\text{Dividend} = \text{Divisor} \times \text{Quotient} + \text{Remainder}$$

$$x^3 - 3x^2 + x + 2 = g(x) \times (x - 2) + (-2x + 4)$$

$$x^3 - 3x^2 + x + 2 + 2x - 4 = g(x)(x - 2)$$

$$x^3 - 3x^2 + 3x - 2 = g(x)(x - 2)$$

$g(x)$  is the quotient when we divide  $(x^3 - 3x^2 + 3x - 2)$  by  $(x - 2)$

$$\begin{array}{r}
 x^2 - x + 1 \\
 x - 2 \overline{) x^3 - 3x^2 + 3x - 2} \\
 \underline{x^3 - 2x^2} \phantom{+ 3x - 2} \\
 -x^2 + 3x - 2 \\
 \underline{-x^2 + 2x} \phantom{- 2} \\
 +x - 2 \\
 \underline{x - 2} \\
 0
 \end{array}$$

$$\therefore g(x) = (x^2 - x + 1)$$

### Question 5:

Give examples of polynomial  $p(x)$ ,  $g(x)$ ,  $q(x)$  and  $r(x)$ , which satisfy the division algorithm and

(i)  $\deg p(x) = \deg q(x)$

(ii)  $\deg q(x) = \deg r(x)$

(iii)  $\deg r(x) = 0$

### Answer:

According to the division algorithm, if  $p(x)$  and  $g(x)$  are two polynomials with

$g(x) \neq 0$ , then we can find polynomials  $q(x)$  and  $r(x)$  such that

$$p(x) = g(x) \times q(x) + r(x),$$

where  $r(x) = 0$  or degree of  $r(x) <$  degree of  $g(x)$

Degree of a polynomial is the highest power of the variable in the polynomial.

(i)  $\deg p(x) = \deg q(x)$

Degree of quotient will be equal to degree of dividend when divisor is constant ( i.e., when any polynomial is divided by a constant).

Let us assume the division of  $6x^2 + 2x + 2$  by 2.

$$\text{Here, } p(x) = 6x^2 + 2x + 2$$

$$g(x) = 2$$

$$q(x) = 3x^2 + x + 1 \text{ and } r(x) = 0$$

Degree of  $p(x)$  and  $q(x)$  is the same i.e., 2.

Checking for division algorithm,

$$p(x) = g(x) \times q(x) + r(x)$$

$$6x^2 + 2x + 2 = 2(3x^2 + x + 1)$$

$$= 6x^2 + 2x + 2$$

Thus, the division algorithm is satisfied.

$$\text{(ii) } \deg q(x) = \deg r(x)$$

Let us assume the division of  $x^3 + x$  by  $x^2$ ,

$$\text{Here, } p(x) = x^3 + x$$

$$g(x) = x^2$$

$$q(x) = x \text{ and } r(x) = x$$

Clearly, the degree of  $q(x)$  and  $r(x)$  is the same i.e., 1.

Checking for division algorithm,

$$p(x) = g(x) \times q(x) + r(x)$$

$$x^3 + x = (x^2) \times x + x$$

$$x^3 + x = x^3 + x$$

Thus, the division algorithm is satisfied.

$$(iii) \deg r(x) = 0$$

Degree of remainder will be 0 when remainder comes to a constant.

Let us assume the division of  $x^3 + 1$  by  $x^2$ .

$$\text{Here, } p(x) = x^3 + 1$$

$$g(x) = x^2$$

$$q(x) = x \text{ and } r(x) = 1$$

Clearly, the degree of  $r(x)$  is 0.

Checking for division algorithm,

$$p(x) = g(x) \times q(x) + r(x)$$

$$x^3 + 1 = (x^2) \times x + 1$$

$$x^3 + 1 = x^3 + 1$$

Thus, the division algorithm is satisfied.

## Exercise 2.4

### Question 1:

**Verify that the numbers given alongside of the cubic polynomials below are their zeroes. Also verify the relationship between the zeroes and the coefficients in each case:**

$$(i) \quad 2x^3 + x^2 - 5x + 2; \quad \frac{1}{2}, 1, -2$$

$$(ii) \quad x^3 - 4x^2 + 5x - 2; \quad 2, 1, 1$$



**Answer:**

(i)  $p(x) = 2x^3 + x^2 - 5x + 2$ .

Zeroes for this polynomial are  $\frac{1}{2}, 1, -2$

$$\begin{aligned} p\left(\frac{1}{2}\right) &= 2\left(\frac{1}{2}\right)^3 + \left(\frac{1}{2}\right)^2 - 5\left(\frac{1}{2}\right) + 2 \\ &= \frac{1}{4} + \frac{1}{4} - \frac{5}{2} + 2 \\ &= 0 \end{aligned}$$

$$\begin{aligned} p(1) &= 2 \times 1^3 + 1^2 - 5 \times 1 + 2 \\ &= 0 \end{aligned}$$

$$\begin{aligned} p(-2) &= 2(-2)^3 + (-2)^2 - 5(-2) + 2 \\ &= -16 + 4 + 10 + 2 = 0 \end{aligned}$$

Therefore,  $\frac{1}{2}$ , 1, and -2 are the zeroes of the given polynomial.

Comparing the given polynomial with  $ax^3 + bx^2 + cx + d$ , we obtain  $a = 2, b = 1, c = -5, d = 2$

We can take  $\alpha = \frac{1}{2}, \beta = 1, \gamma = -2$

$$\alpha + \beta + \gamma = \frac{1}{2} + 1 + (-2) = -\frac{1}{2} = \frac{-b}{a}$$

$$\alpha\beta + \beta\gamma + \alpha\gamma = \frac{1}{2} \times 1 + 1(-2) + \frac{1}{2}(-2) = \frac{-5}{2} = \frac{c}{a}$$

$$\alpha\beta\gamma = \frac{1}{2} \times 1 \times (-2) = \frac{-1}{1} = \frac{-(-2)}{2} = \frac{-d}{a}$$

Therefore, the relationship between the zeroes and the coefficients is verified.

(ii)  $p(x) = x^3 - 4x^2 + 5x - 2$

Zeroes for this polynomial are 2, 1, 1.

$$p(2) = 2^3 - 4(2^2) + 5(2) - 2$$

$$= 8 - 16 + 10 - 2 = 0$$

$$p(1) = 1^3 - 4(1)^2 + 5(1) - 2$$

$$= 1 - 4 + 5 - 2 = 0$$

Therefore, 2, 1, 1 are the zeroes of the given polynomial.

Comparing the given polynomial with  $ax^3 + bx^2 + cx + d$ , we obtain  $a = 1$ ,  $b = -4$ ,  $c = 5$ ,  $d = -2$ .

Verification of the relationship between zeroes and coefficient of the given polynomial

$$\text{Sum of zeroes} = 2 + 1 + 1 = 4 = \frac{-(-4)}{1} = \frac{-b}{a}$$

$$\text{Multiplication of zeroes taking two at a time} = (2)(1) + (1)(1) + (2)(1) = 2 + 1 + 2 = 5$$

$$= \frac{(5)}{1} = \frac{c}{a}$$

$$\text{Multiplication of zeroes} = 2 \times 1 \times 1 = 2 = \frac{-(-2)}{1} = \frac{-d}{a}$$

Hence, the relationship between the zeroes and the coefficients is verified.

### Question 2:

**Find a cubic polynomial with the sum, sum of the product of its zeroes taken two at a time, and the product of its zeroes as 2, -7, -14 respectively.**

**Answer:**

Let the polynomial be  $ax^3 + bx^2 + cx + d$  and the zeroes be  $\alpha, \beta$ , and  $\gamma$ .

It is given that

$$\alpha + \beta + \gamma = \frac{2}{1} = \frac{-b}{a}$$

$$\alpha\beta + \beta\gamma + \alpha\gamma = \frac{-7}{1} = \frac{c}{a}$$

$$\alpha\beta\gamma = \frac{-14}{1} = \frac{-d}{a}$$

If  $a = 1$ , then  $b = -2$ ,  $c = -7$ ,  $d = 14$

Hence, the polynomial is  $x^3 - 2x^2 - 7x + 14$ .

### Question 3:

If the zeroes of polynomial  $x^3 - 3x^2 + x + 1$  are  $a - b, a, a + b$ , find  $a$  and  $b$ .

**Answer:**

$$p(x) = x^3 - 3x^2 + x + 1$$

Zeroes are  $a - b, a, a + b$

Comparing the given polynomial with  $px^3 + qx^2 + rx + t$ , we obtain

$$p = 1, q = -3, r = 1, t = 1$$

$$\text{Sum of zeroes} = a - b + a + a + b$$

$$\frac{-q}{p} = 3a$$

$$\frac{-(-3)}{1} = 3a$$

$$3 = 3a$$

$$a = 1$$

The zeroes are  $1 - b, 1, 1 + b$ .

Multiplication of zeroes =  $1(1-b)(1+b)$

$$\frac{-t}{p} = 1 - b^2$$

$$\frac{-1}{1} = 1 - b^2$$

$$1 - b^2 = -1$$

$$1 + 1 = b^2$$

$$b = \pm\sqrt{2}$$

Hence,  $a = 1$  and  $b = \sqrt{2}$  or  $-\sqrt{2}$ .

#### Question 4:

**It two zeroes of the polynomial  $x^4 - 6x^3 - 26x^2 + 138x - 35$  are  $2 \pm \sqrt{3}$ , find other zeroes.**

#### Answer:

Given that  $2 + \sqrt{3}$  and  $2 - \sqrt{3}$  are zeroes of the given polynomial.

Therefore,  $(x - 2 - \sqrt{3})(x - 2 + \sqrt{3}) = x^2 + 4 - 4x - 3$

$= x^2 - 4x + 1$  is a factor of the given polynomial

For finding the remaining zeroes of the given polynomial, we will find the quotient by dividing  $x^4 - 6x^3 - 26x^2 + 138x - 35$  by  $x^2 - 4x + 1$ .

$$\begin{array}{r}
 x^2 - 2x - 35 \\
 x^2 - 4x + 1 \overline{) x^4 - 6x^3 - 26x^2 + 138x - 35} \\
 \underline{x^4 - 4x^3 + \phantom{x^2} \phantom{x} \phantom{-35}} \\
 -2x^3 - 27x^2 + 138x - 35 \\
 \underline{-2x^3 + 8x^2 - 2x \phantom{-35}} \\
 -35x^2 + 140x - 35 \\
 \underline{-35x^2 + 140x - 35} \\
 0
 \end{array}$$

Clearly,  $x^4 - 6x^3 - 26x^2 + 138x - 35 = (x^2 - 4x + 1)(x^2 - 2x - 35)$

It can be observed that  $(x^2 - 2x - 35)$  is also a factor of the given polynomial.

And  $(x^2 - 2x - 35) = (x - 7)(x + 5)$

Therefore, the value of the polynomial is also zero when  $x - 7 = 0$  or  $x + 5 = 0$

Or  $x = 7$  or  $-5$

Hence, 7 and -5 are also zeroes of this polynomial.

### Question 5:

**If the polynomial  $x^4 - 6x^3 + 16x^2 - 25x + 10$  is divided by another polynomial  $x^2 - 2x + k$ , the remainder comes out to be  $x + a$ , find  $k$  and  $a$ .**

**Answer:**

By division algorithm,

Dividend = Divisor  $\times$  Quotient + Remainder



$$-5 - a = 0$$

Therefore,  $a = -5$

Hence,  $k = 5$  and  $a = -5$