

## Chapter 13 – Limits and Derivatives

---

Page No 301:

Question 1:

Evaluate the Given limit:  $\lim_{x \rightarrow 3} x + 3$

Answer:

$$\lim_{x \rightarrow 3} x + 3 = 3 + 3 = 6$$

---

Question 2:

Evaluate the Given limit:  $\lim_{x \rightarrow \pi} \left( x - \frac{22}{7} \right)$

Answer:

$$\lim_{x \rightarrow \pi} \left( x - \frac{22}{7} \right) = \left( \pi - \frac{22}{7} \right)$$

---

Question 3:

Evaluate the Given limit:  $\lim_{r \rightarrow 1} \pi r^2$

Answer:

$$\lim_{r \rightarrow 1} \pi r^2 = \pi (1)^2 = \pi$$

---

Question 4:

Evaluate the Given limit:  $\lim_{x \rightarrow 4} \frac{4x+3}{x-2}$

Answer:

$$\lim_{x \rightarrow 4} \frac{4x+3}{x-2} = \frac{4(4)+3}{4-2} = \frac{16+3}{2} = \frac{19}{2}$$

---

Question 5:

Evaluate the Given limit:  $\lim_{x \rightarrow -1} \frac{x^{10} + x^5 + 1}{x - 1}$

Answer:

$$\lim_{x \rightarrow -1} \frac{x^{10} + x^5 + 1}{x - 1} = \frac{(-1)^{10} + (-1)^5 + 1}{-1 - 1} = \frac{1 - 1 + 1}{-2} = -\frac{1}{2}$$

---

Question 6:

$$\lim_{x \rightarrow 0} \frac{(x+1)^5 - 1}{x}$$

Evaluate the Given limit:

Answer:

$$\lim_{x \rightarrow 0} \frac{(x+1)^5 - 1}{x}$$

Put  $x + 1 = y$  so that  $y \rightarrow 1$  as  $x \rightarrow 0$ .

$$\begin{aligned} \text{Accordingly, } \lim_{x \rightarrow 0} \frac{(x+1)^5 - 1}{x} &= \lim_{y \rightarrow 1} \frac{y^5 - 1}{y - 1} \\ &= \lim_{y \rightarrow 1} \frac{y^5 - 1^5}{y - 1} \\ &= 5 \cdot 1^{5-1} && \left[ \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1} \right] \\ &= 5 \end{aligned}$$

$$\therefore \lim_{x \rightarrow 0} \frac{(x+5)^5 - 1}{x} = 5$$

---

Question 7:

$$\lim_{x \rightarrow 2} \frac{3x^2 - x - 10}{x^2 - 4}$$

Evaluate the Given limit:

Answer:

At  $x = 2$ , the value of the given rational function takes the form  $\frac{0}{0}$ .

$$\begin{aligned}\therefore \lim_{x \rightarrow 2} \frac{3x^2 - x - 10}{x^2 - 4} &= \lim_{x \rightarrow 2} \frac{(x-2)(3x+5)}{(x-2)(x+2)} \\&= \lim_{x \rightarrow 2} \frac{3x+5}{x+2} \\&= \frac{3(2)+5}{2+2} \\&= \frac{11}{4}\end{aligned}$$


---

Question 8:

$$\lim_{x \rightarrow 3} \frac{x^4 - 81}{2x^2 - 5x - 3}$$

Evaluate the Given limit:

Answer:

At  $x = 2$ , the value of the given rational function takes the form  $\frac{0}{0}$ .

$$\begin{aligned}\therefore \lim_{x \rightarrow 3} \frac{x^4 - 81}{2x^2 - 5x - 3} &= \lim_{x \rightarrow 3} \frac{(x-3)(x+3)(x^2 + 9)}{(x-3)(2x+1)} \\&= \lim_{x \rightarrow 3} \frac{(x+3)(x^2 + 9)}{2x+1} \\&= \frac{(3+3)(3^2 + 9)}{2(3)+1} \\&= \frac{6 \times 18}{7} \\&= \frac{108}{7}\end{aligned}$$


---

Question 9:

$$\lim_{x \rightarrow 0} \frac{ax+b}{cx+1}$$

Evaluate the Given limit:

Answer:

$$\lim_{x \rightarrow 0} \frac{ax+b}{cx+1} = \frac{a(0)+b}{c(0)+1} = b$$


---

Question 10:

$$\lim_{z \rightarrow 1} \frac{z^{\frac{1}{3}} - 1}{z^{\frac{1}{6}} - 1}$$

Evaluate the Given limit:

Answer:

$$\lim_{z \rightarrow 1} \frac{z^{\frac{1}{3}} - 1}{z^{\frac{1}{6}} - 1}$$

$$\frac{0}{0}$$

At  $z = 1$ , the value of the given function takes the form  $\frac{0}{0}$ .

Put  $z^{\frac{1}{6}} = x$  so that  $z \rightarrow 1$  as  $x \rightarrow 1$ .

$$\begin{aligned} \text{Accordingly, } \lim_{z \rightarrow 1} \frac{z^{\frac{1}{3}} - 1}{z^{\frac{1}{6}} - 1} &= \lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1} \\ &= \lim_{x \rightarrow 1} \frac{x^2 - 1^2}{x - 1} \\ &= 2 \cdot 1^{2-1} && \left[ \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1} \right] \\ &= 2 \end{aligned}$$

$$\therefore \lim_{z \rightarrow 1} \frac{z^{\frac{1}{3}} - 1}{z^{\frac{1}{6}} - 1} = 2$$

Question 11:

$$\lim_{x \rightarrow 1} \frac{ax^2 + bx + c}{cx^2 + bx + a}, a + b + c \neq 0$$

Evaluate the Given limit:

Answer:

$$\begin{aligned} \lim_{x \rightarrow 1} \frac{ax^2 + bx + c}{cx^2 + bx + a} &= \frac{a(1)^2 + b(1) + c}{c(1)^2 + b(1) + a} \\ &= \frac{a+b+c}{a+b+c} \\ &= 1 && [a+b+c \neq 0] \end{aligned}$$

Question 12:

$$\lim_{x \rightarrow -2} \frac{\frac{1}{x} + \frac{1}{2}}{x+2}$$

Evaluate the Given limit:

Answer:

$$\lim_{x \rightarrow -2} \frac{\frac{1}{x} + \frac{1}{2}}{x+2}$$

At  $x = -2$ , the value of the given function takes the form  $\frac{0}{0}$ .

$$\begin{aligned} \text{Now, } \lim_{x \rightarrow -2} \frac{\frac{1}{x} + \frac{1}{2}}{x+2} &= \lim_{x \rightarrow -2} \frac{\left( \frac{2+x}{2x} \right)}{x+2} \\ &= \lim_{x \rightarrow -2} \frac{1}{2x} \\ &= \frac{1}{2(-2)} = \frac{-1}{4} \end{aligned}$$


---

Question 13:

$$\lim_{x \rightarrow 0} \frac{\sin ax}{bx}$$

Evaluate the Given limit:

$$\lim_{x \rightarrow 0} \frac{\sin ax}{bx}$$

At  $x = 0$ , the value of the given function takes the form  $\frac{0}{0}$ .

$$\begin{aligned} \text{Now, } \lim_{x \rightarrow 0} \frac{\sin ax}{bx} &= \lim_{x \rightarrow 0} \frac{\sin ax}{ax} \times \frac{ax}{bx} \\ &= \lim_{x \rightarrow 0} \left( \frac{\sin ax}{ax} \right) \times \left( \frac{a}{b} \right) \\ &= \frac{a}{b} \lim_{ax \rightarrow 0} \left( \frac{\sin ax}{ax} \right) \quad [x \rightarrow 0 \Rightarrow ax \rightarrow 0] \\ &= \frac{a}{b} \times 1 \quad \left[ \lim_{y \rightarrow 0} \frac{\sin y}{y} = 1 \right] \\ &= \frac{a}{b} \end{aligned}$$


---

Question 14:

Evaluate the Given limit:  $\lim_{x \rightarrow 0} \frac{\sin ax}{\sin bx}$ ,  $a, b \neq 0$

Answer:

$$\lim_{x \rightarrow 0} \frac{\sin ax}{\sin bx}, a, b \neq 0$$

$\frac{0}{0}$

At  $x = 0$ , the value of the given function takes the form  $\frac{0}{0}$ .

$$\begin{aligned} \text{Now, } \lim_{x \rightarrow 0} \frac{\sin ax}{\sin bx} &= \lim_{x \rightarrow 0} \frac{\left( \frac{\sin ax}{ax} \right) \times ax}{\left( \frac{\sin bx}{bx} \right) \times bx} \\ &= \left( \frac{a}{b} \right) \times \frac{\lim_{ax \rightarrow 0} \left( \frac{\sin ax}{ax} \right)}{\lim_{bx \rightarrow 0} \left( \frac{\sin bx}{bx} \right)} && \left[ \begin{array}{l} x \rightarrow 0 \Rightarrow ax \rightarrow 0 \\ \text{and } x \rightarrow 0 \Rightarrow bx \rightarrow 0 \end{array} \right] \\ &= \left( \frac{a}{b} \right) \times \frac{1}{1} && \left[ \lim_{y \rightarrow 0} \frac{\sin y}{y} = 1 \right] \\ &= \frac{a}{b} \end{aligned}$$


---

Page No 302:

Question 15:

$$\lim_{x \rightarrow \pi} \frac{\sin(\pi - x)}{\pi(\pi - x)}$$

Evaluate the Given limit:

Answer:

$$\lim_{x \rightarrow \pi} \frac{\sin(\pi - x)}{\pi(\pi - x)}$$

It is seen that  $x \rightarrow \pi \Rightarrow (\pi - x) \rightarrow 0$

$$\begin{aligned} \therefore \lim_{x \rightarrow \pi} \frac{\sin(\pi - x)}{\pi(\pi - x)} &= \frac{1}{\pi} \lim_{(\pi-x) \rightarrow 0} \frac{\sin(\pi - x)}{(\pi - x)} \\ &= \frac{1}{\pi} \times 1 && \left[ \lim_{y \rightarrow 0} \frac{\sin y}{y} = 1 \right] \\ &= \frac{1}{\pi} \end{aligned}$$


---

Question 16:

$$\lim_{x \rightarrow 0} \frac{\cos x}{\pi - x}$$

Answer:

$$\lim_{x \rightarrow 0} \frac{\cos x}{\pi - x} = \frac{\cos 0}{\pi - 0} = \frac{1}{\pi}$$

---

Question 17:

$$\lim_{x \rightarrow 0} \frac{\cos 2x - 1}{\cos x - 1}$$

Answer:

$$\lim_{x \rightarrow 0} \frac{\cos 2x - 1}{\cos x - 1}$$

At  $x = 0$ , the value of the given function takes the form  $\frac{0}{0}$ .

Now,

$$\begin{aligned}
\lim_{x \rightarrow 0} \frac{\cos 2x - 1}{\cos x - 1} &= \lim_{x \rightarrow 0} \frac{1 - 2\sin^2 x - 1}{1 - 2\sin^2 \frac{x}{2} - 1} \quad \left[ \cos x = 1 - 2\sin^2 \frac{x}{2} \right] \\
&= \lim_{x \rightarrow 0} \frac{\sin^2 x}{\sin^2 \frac{x}{2}} = \lim_{x \rightarrow 0} \frac{\left( \frac{\sin^2 x}{x^2} \right) \times x^2}{\left( \frac{\sin^2 \frac{x}{2}}{\left( \frac{x}{2} \right)^2} \right) \times \frac{x^2}{4}} \\
&= 4 \lim_{x \rightarrow 0} \left( \frac{\sin^2 x}{x^2} \right) \\
&\quad \lim_{x \rightarrow 0} \left( \frac{\sin^2 \frac{x}{2}}{\left( \frac{x}{2} \right)^2} \right) \\
&= 4 \frac{\left( \lim_{x \rightarrow 0} \frac{\sin x}{x} \right)^2}{\left( \lim_{\frac{x}{2} \rightarrow 0} \frac{\sin \frac{x}{2}}{\frac{x}{2}} \right)^2} \quad \left[ x \rightarrow 0 \Rightarrow \frac{x}{2} \rightarrow 0 \right] \\
&= 4 \frac{1^2}{1^2} \quad \left[ \lim_{y \rightarrow 0} \frac{\sin y}{y} = 1 \right] \\
&= 4
\end{aligned}$$


---

Question 18:

$$\lim_{x \rightarrow 0} \frac{ax + x \cos x}{b \sin x}$$

Evaluate the Given limit:

Answer:

$$\lim_{x \rightarrow 0} \frac{ax + x \cos x}{b \sin x}$$

At  $x = 0$ , the value of the given function takes the form  $\frac{0}{0}$ .

Now,

$$\begin{aligned}
\lim_{x \rightarrow 0} \frac{ax + x \cos x}{b \sin x} &= \frac{1}{b} \lim_{x \rightarrow 0} \frac{x(a + \cos x)}{\sin x} \\
&= \frac{1}{b} \lim_{x \rightarrow 0} \left( \frac{x}{\sin x} \right) \times \lim_{x \rightarrow 0} (a + \cos x) \\
&= \frac{1}{b} \times \left( \lim_{x \rightarrow 0} \frac{\sin x}{x} \right)^{-1} \times \lim_{x \rightarrow 0} (a + \cos x) \\
&= \frac{1}{b} \times (a + \cos 0) \quad \left[ \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right] \\
&= \frac{a+1}{b}
\end{aligned}$$


---

Question 19:

Evaluate the Given limit:  $\lim_{x \rightarrow 0} x \sec x$

Answer:

$$\lim_{x \rightarrow 0} x \sec x = \lim_{x \rightarrow 0} \frac{x}{\cos x} = \frac{0}{\cos 0} = \frac{0}{1} = 0$$


---

Question 20:

Evaluate the Given limit:  $\lim_{x \rightarrow 0} \frac{\sin ax + bx}{ax + \sin bx}$   $a, b, a+b \neq 0$

Answer:

At  $x = 0$ , the value of the given function takes the form  $\frac{0}{0}$ .

Now,

$$\begin{aligned}
& \lim_{x \rightarrow 0} \frac{\sin ax + bx}{ax + \sin bx} \\
&= \lim_{x \rightarrow 0} \frac{\left(\frac{\sin ax}{ax}\right)ax + bx}{ax + bx\left(\frac{\sin bx}{bx}\right)} \\
&= \frac{\left(\lim_{ax \rightarrow 0} \frac{\sin ax}{ax}\right) \times \lim_{x \rightarrow 0}(ax) + \lim_{x \rightarrow 0} bx}{\lim_{x \rightarrow 0} ax + \lim_{x \rightarrow 0} bx \left(\lim_{bx \rightarrow 0} \frac{\sin bx}{bx}\right)} \quad [\text{As } x \rightarrow 0 \Rightarrow ax \rightarrow 0 \text{ and } bx \rightarrow 0] \\
&= \frac{\lim_{x \rightarrow 0}(ax) + \lim_{x \rightarrow 0} bx}{\lim_{x \rightarrow 0} ax + \lim_{x \rightarrow 0} bx} \quad \left[ \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right] \\
&= \frac{\lim_{x \rightarrow 0}(ax + bx)}{\lim_{x \rightarrow 0}(ax + bx)} \\
&= \lim_{x \rightarrow 0}(1) \\
&= 1
\end{aligned}$$


---

Question 21:

Evaluate the Given limit:  $\lim_{x \rightarrow 0} (\cosec x - \cot x)$

Answer:

At  $x = 0$ , the value of the given function takes the form  $\infty - \infty$ .

Now,

$$\begin{aligned}
& \lim_{x \rightarrow 0} (\cosec x - \cot x) \\
&= \lim_{x \rightarrow 0} \left( \frac{1}{\sin x} - \frac{\cos x}{\sin x} \right) \\
&= \lim_{x \rightarrow 0} \left( \frac{1 - \cos x}{\sin x} \right) \\
&= \lim_{x \rightarrow 0} \frac{\left( \frac{1 - \cos x}{x} \right)}{\left( \frac{\sin x}{x} \right)} \\
&= \frac{\lim_{x \rightarrow 0} \frac{1 - \cos x}{x}}{\lim_{x \rightarrow 0} \frac{\sin x}{x}} \\
&= \frac{0}{1} \quad \left[ \lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0 \text{ and } \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right] \\
&= 0
\end{aligned}$$


---

Question 22:

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{\tan 2x}{x - \frac{\pi}{2}}$$

Answer:

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{\tan 2x}{x - \frac{\pi}{2}}$$

At  $x = \frac{\pi}{2}$ , the value of the given function takes the form  $\frac{0}{0}$ .

Now, put  $x - \frac{\pi}{2} = y$  so that  $x \rightarrow \frac{\pi}{2}, y \rightarrow 0$ .

$$\begin{aligned}
& \therefore \lim_{x \rightarrow \frac{\pi}{2}^-} \frac{\tan 2x}{x - \frac{\pi}{2}} = \lim_{y \rightarrow 0} \frac{\tan 2\left(y + \frac{\pi}{2}\right)}{y} \\
&= \lim_{y \rightarrow 0} \frac{\tan(\pi + 2y)}{y} \\
&= \lim_{y \rightarrow 0} \frac{\tan 2y}{y} \quad [\tan(\pi + 2y) = \tan 2y] \\
&= \lim_{y \rightarrow 0} \frac{\sin 2y}{y \cos 2y} \\
&= \lim_{y \rightarrow 0} \left( \frac{\sin 2y}{2y} \times \frac{2}{\cos 2y} \right) \\
&= \left( \lim_{2y \rightarrow 0} \frac{\sin 2y}{2y} \right) \times \lim_{y \rightarrow 0} \left( \frac{2}{\cos 2y} \right) \quad [y \rightarrow 0 \Rightarrow 2y \rightarrow 0] \\
&= 1 \times \frac{2}{\cos 0} \quad \left[ \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right] \\
&= 1 \times \frac{2}{1} \\
&= 2
\end{aligned}$$


---

Question 23:

$$\text{Find } \lim_{x \rightarrow 0^-} f(x) \text{ and } \lim_{x \rightarrow 1^+} f(x), \text{ where } f(x) = \begin{cases} 2x + 3, & x \leq 0 \\ 3(x+1), & x > 0 \end{cases}$$

Answer:

The given function is

$$f(x) = \begin{cases} 2x + 3, & x \leq 0 \\ 3(x+1), & x > 0 \end{cases}$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} [2x + 3] = 2(0) + 3 = 3$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} 3(x+1) = 3(0+1) = 3$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} f(x) = 3$$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1} 3(x+1) = 3(1+1) = 6$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} 3(x+1) = 3(1+1) = 6$$

$$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} f(x) = 6$$


---

Question 24:

Find  $\lim_{x \rightarrow 1} f(x)$ , where  $f(x) = \begin{cases} x^2 - 1, & x \leq 1 \\ -x^2 - 1, & x > 1 \end{cases}$

Answer:

The given function is

$$f(x) = \begin{cases} x^2 - 1, & x \leq 1 \\ -x^2 - 1, & x > 1 \end{cases}$$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1} [x^2 - 1] = 1^2 - 1 = 1 - 1 = 0$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} [-x^2 - 1] = -1^2 - 1 = -1 - 1 = -2$$

It is observed that  $\lim_{x \rightarrow 1^-} f(x) \neq \lim_{x \rightarrow 1^+} f(x)$ .

Hence,  $\lim_{x \rightarrow 1} f(x)$  does not exist.

---

Question 25:

Evaluate  $\lim_{x \rightarrow 0} f(x)$ , where  $f(x) = \begin{cases} \frac{|x|}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$

Answer:

The given function is

$$f(x) = \begin{cases} \frac{|x|}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

$$\begin{aligned}
\lim_{x \rightarrow 0^-} f(x) &= \lim_{x \rightarrow 0^-} \left[ \frac{|x|}{x} \right] \\
&= \lim_{x \rightarrow 0^-} \left( \frac{-x}{x} \right) \quad [\text{When } x \text{ is negative, } |x| = -x] \\
&= \lim_{x \rightarrow 0^-} (-1) \\
&= -1
\end{aligned}$$

$$\begin{aligned}
\lim_{x \rightarrow 0^+} f(x) &= \lim_{x \rightarrow 0^+} \left[ \frac{|x|}{x} \right] \\
&= \lim_{x \rightarrow 0^+} \left[ \frac{x}{x} \right] \quad [\text{When } x \text{ is positive, } |x| = x] \\
&= \lim_{x \rightarrow 0^+} (1) \\
&= 1
\end{aligned}$$

It is observed that  $\lim_{x \rightarrow 0^-} f(x) \neq \lim_{x \rightarrow 0^+} f(x)$ .

Hence,  $\lim_{x \rightarrow 0} f(x)$  does not exist.

---

**Question 26:**

$$\text{Find } \lim_{x \rightarrow 0} f(x), \text{ where } f(x) = \begin{cases} \frac{x}{|x|}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

**Answer:**

The given function is

$$f(x) = \begin{cases} \frac{x}{|x|}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

$$\begin{aligned}\lim_{x \rightarrow 0^-} f(x) &= \lim_{x \rightarrow 0^-} \left[ \frac{x}{|x|} \right] \\ &= \lim_{x \rightarrow 0^-} \left[ \frac{x}{-x} \right] \quad [\text{When } x < 0, |x| = -x] \\ &= \lim_{x \rightarrow 0^-} (-1) \\ &= -1\end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow 0^+} f(x) &= \lim_{x \rightarrow 0^+} \left[ \frac{x}{|x|} \right] \\ &= \lim_{x \rightarrow 0^+} \left[ \frac{x}{x} \right] \quad [\text{When } x > 0, |x| = x] \\ &= \lim_{x \rightarrow 0^+} (1) \\ &= 1\end{aligned}$$

It is observed that  $\lim_{x \rightarrow 0^-} f(x) \neq \lim_{x \rightarrow 0^+} f(x)$ .

Hence,  $\lim_{x \rightarrow 0} f(x)$  does not exist.

---

**Question 27:**

Find  $\lim_{x \rightarrow 5} f(x)$ , where  $f(x) = |x| - 5$

**Answer:**

The given function is  $f(x) = |x| - 5$ .

$$\begin{aligned}\lim_{x \rightarrow 5^-} f(x) &= \lim_{x \rightarrow 5^-} [|x| - 5] \\ &= \lim_{x \rightarrow 5^-} (x - 5) \quad [\text{When } x < 0, |x| = x] \\ &= 5 - 5 \\ &= 0\end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow 5^+} f(x) &= \lim_{x \rightarrow 5^+} (|x| - 5) \\ &= \lim_{x \rightarrow 5^+} (x - 5) \quad [\text{When } x > 0, |x| = x] \\ &= 5 - 5 \\ &= 0\end{aligned}$$

$$\therefore \lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^+} f(x) = 0$$

Hence,  $\lim_{x \rightarrow 5} f(x) = 0$

---

Question 28:

$$\text{Suppose } f(x) = \begin{cases} a+bx, & x < 1 \\ 4, & x = 1 \\ b-ax, & x > 1 \end{cases} \text{ and if } \lim_{x \rightarrow 1} f(x) = f(1) \text{ what are possible values of } a \text{ and } b?$$

Answer:

The given function is

$$f(x) = \begin{cases} a+bx, & x < 1 \\ 4, & x = 1 \\ b-ax, & x > 1 \end{cases}$$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1} (a+bx) = a+b$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} (b-ax) = b-a$$

$$f(1) = 4$$

It is given that  $\lim_{x \rightarrow 1} f(x) = f(1)$ .

$$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} f(x) = f(1)$$

$$\Rightarrow a+b = 4 \text{ and } b-a = 4$$

On solving these two equations, we obtain  $a = 0$  and  $b = 4$ .

Thus, the respective possible values of  $a$  and  $b$  are 0 and 4.

---

Page No 303:

Question 29:

Let  $a_1, a_2, \dots, a_n$  be fixed real numbers and define a function

$$f(x) = (x - a_1)(x - a_2) \dots (x - a_n)$$

What is  $\lim_{x \rightarrow a_i} f(x)$ ? For some  $a \neq a_1, a_2, \dots, a_n$ , compute  $\lim_{x \rightarrow a} f(x)$ .

Answer:

The given function is  $f(x) = (x - a_1)(x - a_2) \dots (x - a_n)$

$$\begin{aligned}\lim_{x \rightarrow a_1} f(x) &= \lim_{x \rightarrow a_1} [(x - a_1)(x - a_2) \dots (x - a_n)] \\ &= \left[ \lim_{x \rightarrow a_1} (x - a_1) \right] \left[ \lim_{x \rightarrow a_1} (x - a_2) \right] \dots \left[ \lim_{x \rightarrow a_1} (x - a_n) \right] \\ &= (a_1 - a_1)(a_1 - a_2) \dots (a_1 - a_n) = 0\end{aligned}$$

$$\therefore \lim_{x \rightarrow a_1} f(x) = 0$$

$$\begin{aligned}\text{Now, } \lim_{x \rightarrow a} f(x) &= \lim_{x \rightarrow a} [(x - a_1)(x - a_2) \dots (x - a_n)] \\ &= \left[ \lim_{x \rightarrow a} (x - a_1) \right] \left[ \lim_{x \rightarrow a} (x - a_2) \right] \dots \left[ \lim_{x \rightarrow a} (x - a_n) \right] \\ &= (a - a_1)(a - a_2) \dots (a - a_n)\end{aligned}$$

$$\therefore \lim_{x \rightarrow a} f(x) = (a - a_1)(a - a_2) \dots (a - a_n)$$


---

Question 30:

$$\text{If } f(x) = \begin{cases} |x| + 1, & x < 0 \\ 0, & x = 0 \\ |x| - 1, & x > 0 \end{cases}$$

For what value (s) of  $a$  does  $\lim_{x \rightarrow a} f(x)$  exists?

Answer:

The given function is

$$f(x) = \begin{cases} |x| + 1, & x < 0 \\ 0, & x = 0 \\ |x| - 1, & x > 0 \end{cases}$$

When  $a = 0$ ,

$$\begin{aligned}\lim_{x \rightarrow 0^-} f(x) &= \lim_{x \rightarrow 0^-} (|x| + 1) \\ &= \lim_{x \rightarrow 0^-} (-x + 1) \quad [\text{If } x < 0, |x| = -x] \\ &= -0 + 1 \\ &= 1\end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow 0^+} f(x) &= \lim_{x \rightarrow 0^+} (|x| - 1) \\ &= \lim_{x \rightarrow 0^+} (x - 1) \quad [\text{If } x > 0, |x| = x] \\ &= 0 - 1 \\ &= -1\end{aligned}$$

Here, it is observed that  $\lim_{x \rightarrow 0^-} f(x) \neq \lim_{x \rightarrow 0^+} f(x)$ .

$\therefore \lim_{x \rightarrow 0} f(x)$  does not exist.

When  $a < 0$ ,

$$\begin{aligned}\lim_{x \rightarrow a^-} f(x) &= \lim_{x \rightarrow a^-} (|x| + 1) \\ &= \lim_{x \rightarrow a} (-x + 1) \quad [x < a < 0 \Rightarrow |x| = -x] \\ &= -a + 1\end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow a^+} f(x) &= \lim_{x \rightarrow a^+} (|x| + 1) \\ &= \lim_{x \rightarrow a} (-x + 1) \quad [a < x < 0 \Rightarrow |x| = -x] \\ &= -a + 1\end{aligned}$$

$$\therefore \lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = -a + 1$$

Thus, limit of  $f(x)$  exists at  $x = a$ , where  $a < 0$ .

When  $a > 0$

$$\begin{aligned}\lim_{x \rightarrow a^-} f(x) &= \lim_{x \rightarrow a^-} (|x| - 1) \\ &= \lim_{x \rightarrow a} (x - 1) \quad [0 < x < a \Rightarrow |x| = x] \\ &= a - 1\end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow a^+} f(x) &= \lim_{x \rightarrow a^+} (|x| - 1) \\ &= \lim_{x \rightarrow a} (x - 1) \quad [0 < a < x \Rightarrow |x| = x] \\ &= a - 1\end{aligned}$$

$$\therefore \lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = a - 1$$

Thus, limit of  $f(x)$  exists at  $x = a$ , where  $a > 0$ .

Thus,  $\lim_{x \rightarrow a} f(x)$  exists for all  $a \neq 0$ .

---

Question 31:

If the function  $f(x)$  satisfies  $\lim_{x \rightarrow 1} \frac{f(x) - 2}{x^2 - 1} = \pi$ , evaluate  $\lim_{x \rightarrow 1} f(x)$ .

Answer:

$$\begin{aligned}
& \lim_{x \rightarrow 1} \frac{f(x) - 2}{x^2 - 1} = \pi \\
& \Rightarrow \frac{\lim_{x \rightarrow 1}(f(x) - 2)}{\lim_{x \rightarrow 1}(x^2 - 1)} = \pi \\
& \Rightarrow \lim_{x \rightarrow 1}(f(x) - 2) = \pi \lim_{x \rightarrow 1}(x^2 - 1) \\
& \Rightarrow \lim_{x \rightarrow 1}(f(x) - 2) = \pi(1^2 - 1) \\
& \Rightarrow \lim_{x \rightarrow 1}(f(x) - 2) = 0 \\
& \Rightarrow \lim_{x \rightarrow 1} f(x) - \lim_{x \rightarrow 1} 2 = 0 \\
& \Rightarrow \lim_{x \rightarrow 1} f(x) - 2 = 0 \\
& \therefore \lim_{x \rightarrow 1} f(x) = 2
\end{aligned}$$


---

Question 32:

$$\text{If } f(x) = \begin{cases} mx^2 + n, & x < 0 \\ nx + m, & 0 \leq x \leq 1 \\ nx^3 + m, & x > 1 \end{cases}. \text{ For what integers } m \text{ and } n \text{ does } \lim_{x \rightarrow 0} f(x) \text{ and } \lim_{x \rightarrow 1} f(x) \text{ exist?}$$

Answer:

The given function is

$$f(x) = \begin{cases} mx^2 + n, & x < 0 \\ nx + m, & 0 \leq x \leq 1 \\ nx^3 + m, & x > 1 \end{cases}$$

$$\begin{aligned}
\lim_{x \rightarrow 0^-} f(x) &= \lim_{x \rightarrow 0} (mx^2 + n) \\
&= m(0)^2 + n \\
&= n
\end{aligned}$$

$$\begin{aligned}
\lim_{x \rightarrow 0^+} f(x) &= \lim_{x \rightarrow 0} (nx + m) \\
&= n(0) + m \\
&= m.
\end{aligned}$$

Thus,  $\lim_{x \rightarrow 0} f(x)$  exists if  $m = n$ .

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1} (nx + m)$$

$$= n(1) + m$$

$$= m + n$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} (nx^3 + m)$$

$$= n(1)^3 + m$$

$$= m + n$$

$$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} f(x).$$

Thus,  $\lim_{x \rightarrow 1} f(x)$  exists for any integral value of m and n.

---

Page No 312:

Question 1:

Find the derivative of  $x^2 - 2$  at  $x = 10$ .

Answer:

Let  $f(x) = x^2 - 2$ . Accordingly,

$$\begin{aligned}f'(10) &= \lim_{h \rightarrow 0} \frac{f(10+h) - f(10)}{h} \\&= \lim_{h \rightarrow 0} \frac{[(10+h)^2 - 2] - (10^2 - 2)}{h} \\&= \lim_{h \rightarrow 0} \frac{10^2 + 2.10.h + h^2 - 2 - 10^2 + 2}{h} \\&= \lim_{h \rightarrow 0} \frac{20h + h^2}{h} \\&= \lim_{h \rightarrow 0} (20 + h) = (20 + 0) = 20\end{aligned}$$

Thus, the derivative of  $x^2 - 2$  at  $x = 10$  is 20.

---

Question 2:

Find the derivative of  $99x$  at  $x = 100$ .

Answer:

Let  $f(x) = 99x$ . Accordingly,

$$\begin{aligned}
 f'(100) &= \lim_{h \rightarrow 0} \frac{f(100+h) - f(100)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{99(100+h) - 99(100)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{99 \times 100 + 99h - 99 \times 100}{h} \\
 &= \lim_{h \rightarrow 0} \frac{99h}{h} \\
 &= \lim_{h \rightarrow 0} (99) = 99
 \end{aligned}$$

Thus, the derivative of  $99x$  at  $x = 100$  is 99.

---

**Question 3:**

Find the derivative of  $x$  at  $x = 1$ .

**Answer:**

Let  $f(x) = x$ . Accordingly,

$$\begin{aligned}
 f'(1) &= \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{(1+h) - 1}{h} \\
 &= \lim_{h \rightarrow 0} \frac{h}{h} \\
 &= \lim_{h \rightarrow 0} (1) \\
 &= 1
 \end{aligned}$$

Thus, the derivative of  $x$  at  $x = 1$  is 1.

---

**Question 4:**

Find the derivative of the following functions from first principle.

(i)  $x^3 - 27$  (ii)  $(x - 1)(x - 2)$

(iii)  $\frac{1}{x^2}$  (iv)  $\frac{x+1}{x-1}$

**Answer:**

(i) Let  $f(x) = x^3 - 27$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{[(x+h)^3 - 27] - (x^3 - 27)}{h} \\
&= \lim_{h \rightarrow 0} \frac{x^3 + h^3 + 3x^2h + 3xh^2 - x^3}{h} \\
&= \lim_{h \rightarrow 0} \frac{h^3 + 3x^2h + 3xh^2}{h} \\
&= \lim_{h \rightarrow 0} (h^2 + 3x^2 + 3xh) \\
&= 0 + 3x^2 + 0 = 3x^2
\end{aligned}$$

(ii) Let  $f(x) = (x-1)(x-2)$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{(x+h-1)(x+h-2) - (x-1)(x-2)}{h} \\
&= \lim_{h \rightarrow 0} \frac{(x^2 + hx - 2x + hx + h^2 - 2h - x - h + 2) - (x^2 - 2x - x + 2)}{h} \\
&= \lim_{h \rightarrow 0} \frac{(hx + hx + h^2 - 2h - h)}{h} \\
&= \lim_{h \rightarrow 0} \frac{2hx + h^2 - 3h}{h} \\
&= \lim_{h \rightarrow 0} (2x + h - 3) \\
&= (2x + 0 - 3) \\
&= 2x - 3
\end{aligned}$$

(iii) Let  $f(x) = \frac{1}{x^2}$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h)^2} - \frac{1}{x^2}}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{x^2 - (x+h)^2}{x^2 (x+h)^2} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{x^2 - x^2 - h^2 - 2hx}{x^2 (x+h)^2} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-h^2 - 2hx}{x^2 (x+h)^2} \right] \\
&= \lim_{h \rightarrow 0} \left[ \frac{-h - 2x}{x^2 (x+h)^2} \right] \\
&= \frac{0 - 2x}{x^2 (x+0)^2} = \frac{-2}{x^3}
\end{aligned}$$

(iv) Let  $f(x) = \frac{x+1}{x-1}$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\left( \frac{x+h+1}{x+h-1} - \frac{x+1}{x-1} \right)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{(x-1)(x+h+1) - (x+1)(x+h-1)}{(x-1)(x+h-1)} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{(x^2 + hx + x - x - h - 1) - (x^2 + hx - x + x + h - 1)}{(x-1)(x+h-1)} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2h}{(x-1)(x+h-1)} \right] \\
&= \lim_{h \rightarrow 0} \left[ \frac{-2}{(x-1)(x+h-1)} \right] \\
&= \frac{-2}{(x-1)(x-1)} = \frac{-2}{(x-1)^2}
\end{aligned}$$

Question 5:

For the function

$$f(x) = \frac{x^{100}}{100} + \frac{x^{99}}{99} + \dots + \frac{x^2}{2} + x + 1$$

Prove that  $f'(1) = 100f'(0)$

Answer:

The given function is

$$f(x) = \frac{x^{100}}{100} + \frac{x^{99}}{99} + \dots + \frac{x^2}{2} + x + 1$$

$$\frac{d}{dx} f(x) = \frac{d}{dx} \left[ \frac{x^{100}}{100} + \frac{x^{99}}{99} + \dots + \frac{x^2}{2} + x + 1 \right]$$

$$\frac{d}{dx} f(x) = \frac{d}{dx} \left( \frac{x^{100}}{100} \right) + \frac{d}{dx} \left( \frac{x^{99}}{99} \right) + \dots + \frac{d}{dx} \left( \frac{x^2}{2} \right) + \frac{d}{dx}(x) + \frac{d}{dx}(1)$$

On using theorem  $\frac{d}{dx}(x^n) = nx^{n-1}$ , we obtain

$$\begin{aligned} \frac{d}{dx} f(x) &= \frac{100x^{99}}{100} + \frac{99x^{98}}{99} + \dots + \frac{2x}{2} + 1 + 0 \\ &= x^{99} + x^{98} + \dots + x + 1 \end{aligned}$$

$$\therefore f'(x) = x^{99} + x^{98} + \dots + x + 1$$

At  $x = 0$ ,

$$f'(0) = 1$$

At  $x = 1$ ,

$$f'(1) = 1^{99} + 1^{98} + \dots + 1 + 1 = [1 + 1 + \dots + 1 + 1]_{100 \text{ terms}} = 1 \times 100 = 100$$

Thus,  $f'(1) = 100 \times f'(0)$

Page No 313:

Question 6:

Find the derivative of  $x^n + ax^{n-1} + a^2x^{n-2} + \dots + a^{n-1}x + a^n$  for some fixed real number  $a$ .

Answer:

$$\text{Let } f(x) = x^n + ax^{n-1} + a^2x^{n-2} + \dots + a^{n-1}x + a^n$$

$$\begin{aligned}\therefore f'(x) &= \frac{d}{dx} (x^n + ax^{n-1} + a^2 x^{n-2} + \dots + a^{n-1} x + a^n) \\ &= \frac{d}{dx} (x^n) + a \frac{d}{dx} (x^{n-1}) + a^2 \frac{d}{dx} (x^{n-2}) + \dots + a^{n-1} \frac{d}{dx} (x) + a^n \frac{d}{dx} (1)\end{aligned}$$

On using theorem  $\frac{d}{dx} x^n = nx^{n-1}$ , we obtain

$$\begin{aligned}f'(x) &= nx^{n-1} + a(n-1)x^{n-2} + a^2(n-2)x^{n-3} + \dots + a^{n-1} + a^n(0) \\ &= nx^{n-1} + a(n-1)x^{n-2} + a^2(n-2)x^{n-3} + \dots + a^{n-1}\end{aligned}$$


---

**Question 7:**

For some constants a and b, find the derivative of

(i)  $(x - a)(x - b)$  (ii)  $(ax^2 + b)^2$  (iii)  $\frac{x-a}{x-b}$

**Answer:**

(i) Let  $f(x) = (x - a)(x - b)$

$$\Rightarrow f(x) = x^2 - (a+b)x + ab$$

$$\begin{aligned}\therefore f'(x) &= \frac{d}{dx} (x^2 - (a+b)x + ab) \\ &= \frac{d}{dx} (x^2) - (a+b) \frac{d}{dx} (x) + \frac{d}{dx} (ab)\end{aligned}$$

On using theorem  $\frac{d}{dx} (x^n) = nx^{n-1}$ , we obtain

$$f'(x) = 2x - (a+b) + 0 = 2x - a - b$$

(ii) Let  $f(x) = (ax^2 + b)^2$

$$\Rightarrow f(x) = a^2 x^4 + 2abx^2 + b^2$$

$$\therefore f'(x) = \frac{d}{dx} (a^2 x^4 + 2abx^2 + b^2) = a^2 \frac{d}{dx} (x^4) + 2ab \frac{d}{dx} (x^2) + \frac{d}{dx} (b^2)$$

On using theorem  $\frac{d}{dx} x^n = nx^{n-1}$ , we obtain

$$\begin{aligned}f'(x) &= a^2 (4x^3) + 2ab(2x) + b^2(0) \\ &= 4a^2 x^3 + 4abx \\ &= 4ax(ax^2 + b)\end{aligned}$$

(iii) Let  $f(x) = \frac{(x-a)}{(x-b)}$

$$\Rightarrow f'(x) = \frac{d}{dx} \left( \frac{x-a}{x-b} \right)$$

By quotient rule,

$$\begin{aligned}f'(x) &= \frac{(x-b)\frac{d}{dx}(x-a) - (x-a)\frac{d}{dx}(x-b)}{(x-b)^2} \\&= \frac{(x-b)(1) - (x-a)(1)}{(x-b)^2} \\&= \frac{x-b-x+a}{(x-b)^2} \\&= \frac{a-b}{(x-b)^2}\end{aligned}$$

---

Question 8:

$$\frac{x^n - a^n}{x - a}$$

Find the derivative of  $\frac{x^n - a^n}{x - a}$  for some constant a.

Answer:

$$\begin{aligned}\text{Let } f(x) &= \frac{x^n - a^n}{x - a} \\ \Rightarrow f'(x) &= \frac{d}{dx} \left( \frac{x^n - a^n}{x - a} \right)\end{aligned}$$

By quotient rule,

$$\begin{aligned}f'(x) &= \frac{(x-a)\frac{d}{dx}(x^n - a^n) - (x^n - a^n)\frac{d}{dx}(x-a)}{(x-a)^2} \\&= \frac{(x-a)(nx^{n-1} - 0) - (x^n - a^n)}{(x-a)^2} \\&= \frac{nx^n - anx^{n-1} - x^n + a^n}{(x-a)^2}\end{aligned}$$

---

Question 9:

Find the derivative of

(i)  $2x - \frac{3}{4}$  (ii)  $(5x^3 + 3x - 1)(x - 1)$

(iii)  $x^{-3}(5 + 3x)$  (iv)  $x^5(3 - 6x^{-9})$

$$(v) x^{-4} (3 - 4x^{-5}) \quad (vi) \frac{2}{x+1} - \frac{x^2}{3x-1}$$

Answer:

$$(i) \text{ Let } f(x) = 2x - \frac{3}{4}$$

$$\begin{aligned} f'(x) &= \frac{d}{dx} \left( 2x - \frac{3}{4} \right) \\ &= 2 \frac{d}{dx}(x) - \frac{d}{dx} \left( \frac{3}{4} \right) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

$$(ii) \text{ Let } f(x) = (5x^3 + 3x - 1)(x - 1)$$

By Leibnitz product rule,

$$\begin{aligned} f'(x) &= (5x^3 + 3x - 1) \frac{d}{dx}(x - 1) + (x - 1) \frac{d}{dx}(5x^3 + 3x - 1) \\ &= (5x^3 + 3x - 1)(1) + (x - 1)(15x^2 + 3) \\ &= (5x^3 + 3x - 1) + (x - 1)(15x^2 + 3) \\ &= 5x^3 + 3x - 1 + 15x^3 + 3x - 15x^2 - 3 \\ &= 20x^3 - 15x^2 + 6x - 4 \end{aligned}$$

$$(iii) \text{ Let } f(x) = x^{-3} (5 + 3x)$$

By Leibnitz product rule,

$$\begin{aligned} f'(x) &= x^{-3} \frac{d}{dx}(5 + 3x) + (5 + 3x) \frac{d}{dx}(x^{-3}) \\ &= x^{-3} (0 + 3) + (5 + 3x)(-3x^{-3-1}) \\ &= x^{-3} (3) + (5 + 3x)(-3x^{-4}) \\ &= 3x^{-3} - 15x^{-4} - 9x^{-3} \\ &= -6x^{-3} - 15x^{-4} \\ &= -3x^{-3} \left( 2 + \frac{5}{x} \right) \\ &= \frac{-3x^{-3}}{x} (2x + 5) \\ &= \frac{-3}{x^4} (5 + 2x) \end{aligned}$$

$$(iv) \text{ Let } f(x) = x^5 (3 - 6x^{-9})$$

By Leibnitz product rule,

$$\begin{aligned}
f'(x) &= x^5 \frac{d}{dx}(3 - 6x^{-9}) + (3 - 6x^{-9}) \frac{d}{dx}(x^5) \\
&= x^5 \{0 - 6(-9)x^{-9-1}\} + (3 - 6x^{-9})(5x^4) \\
&= x^5(54x^{-10}) + 15x^4 - 30x^{-5} \\
&= 54x^{-5} + 15x^4 - 30x^{-5} \\
&= 24x^{-5} + 15x^4 \\
&= 15x^4 + \frac{24}{x^5}
\end{aligned}$$

(v) Let  $f(x) = x^{-4}(3 - 4x^{-5})$

By Leibnitz product rule,

$$\begin{aligned}
f'(x) &= x^{-4} \frac{d}{dx}(3 - 4x^{-5}) + (3 - 4x^{-5}) \frac{d}{dx}(x^{-4}) \\
&= x^{-4} \{0 - 4(-5)x^{-5-1}\} + (3 - 4x^{-5})(-4)x^{-4-1} \\
&= x^{-4}(20x^{-6}) + (3 - 4x^{-5})(-4x^{-5}) \\
&= 20x^{-10} - 12x^{-5} + 16x^{-10} \\
&= 36x^{-10} - 12x^{-5} \\
&= -\frac{12}{x^5} + \frac{36}{x^{10}}
\end{aligned}$$

(vi) Let  $f(x) = \frac{2}{x+1} - \frac{x^2}{3x-1}$

$$f'(x) = \frac{d}{dx}\left(\frac{2}{x+1}\right) - \frac{d}{dx}\left(\frac{x^2}{3x-1}\right)$$

By quotient rule,

$$\begin{aligned}
f'(x) &= \left[ \frac{(x+1)\frac{d}{dx}(2) - 2\frac{d}{dx}(x+1)}{(x+1)^2} \right] - \left[ \frac{(3x-1)\frac{d}{dx}(x^2) - x^2\frac{d}{dx}(3x-1)}{(3x-1)^2} \right] \\
&= \left[ \frac{(x+1)(0) - 2(1)}{(x+1)^2} \right] - \left[ \frac{(3x-1)(2x) - (x^2)(3)}{(3x-1)^2} \right] \\
&= \frac{-2}{(x+1)^2} - \left[ \frac{6x^2 - 2x - 3x^2}{(3x-1)^2} \right] \\
&= \frac{-2}{(x+1)^2} - \left[ \frac{3x^2 - 2x^2}{(3x-1)^2} \right] \\
&= \frac{-2}{(x+1)^2} - \frac{x(3x-2)}{(3x-1)^2}
\end{aligned}$$

---

Question 10:

Find the derivative of  $\cos x$  from first principle.

Answer:

Let  $f(x) = \cos x$ . Accordingly, from the first principle,

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos x}{h} \\&= \lim_{h \rightarrow 0} \left[ \frac{\cos x \cos h - \sin x \sin h - \cos x}{h} \right] \\&= \lim_{h \rightarrow 0} \left[ \frac{-\cos x(1 - \cos h) - \sin x \sin h}{h} \right] \\&= \lim_{h \rightarrow 0} \left[ \frac{-\cos x(1 - \cos h)}{h} - \frac{\sin x \sin h}{h} \right] \\&= -\cos x \left( \lim_{h \rightarrow 0} \frac{1 - \cos h}{h} \right) - \sin x \lim_{h \rightarrow 0} \left( \frac{\sin h}{h} \right) \\&= -\cos x(0) - \sin x(1) \quad \left[ \lim_{h \rightarrow 0} \frac{1 - \cos h}{h} = 0 \text{ and } \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1 \right] \\&= -\sin x \\∴ f'(x) &= -\sin x\end{aligned}$$

---

Question 11:

Find the derivative of the following functions:

- (i)  $\sin x \cos x$  (ii)  $\sec x$  (iii)  $5 \sec x + 4 \cos x$
- (iv)  $\operatorname{cosec} x$  (v)  $3 \cot x + 5 \operatorname{cosec} x$
- (vi)  $5 \sin x - 6 \cos x + 7$  (vii)  $2 \tan x - 7 \sec x$

Answer:

(i) Let  $f(x) = \sin x \cos x$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\sin(x+h)\cos(x+h) - \sin x \cos x}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{2h} [2\sin(x+h)\cos(x+h) - 2\sin x \cos x] \\
&= \lim_{h \rightarrow 0} \frac{1}{2h} [\sin 2(x+h) - \sin 2x] \\
&= \lim_{h \rightarrow 0} \frac{1}{2h} \left[ 2\cos \frac{2x+2h+2x}{2} \cdot \sin \frac{2x+2h-2x}{2} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \cos \frac{4x+2h}{2} \sin \frac{2h}{2} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} [\cos(2x+h) \sin h] \\
&= \lim_{h \rightarrow 0} \cos(2x+h) \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\
&= \cos(2x+0) \cdot 1 \\
&= \cos 2x
\end{aligned}$$

(ii) Let  $f(x) = \sec x$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\sec(x+h) - \sec x}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\cos(x+h)} - \frac{1}{\cos x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\cos x - \cos(x+h)}{\cos x \cos(x+h)} \right] \\
&= \frac{1}{\cos x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2 \sin\left(\frac{x+x+h}{2}\right) \sin\left(\frac{x-x-h}{2}\right)}{\cos(x+h)} \right] \\
&= \frac{1}{\cos x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2 \sin\left(\frac{2x+h}{2}\right) \sin\left(-\frac{h}{2}\right)}{\cos(x+h)} \right] \\
&= \frac{1}{\cos x} \cdot \lim_{h \rightarrow 0} \frac{\left[ \sin\left(\frac{2x+h}{2}\right) \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \right]}{\cos(x+h)} \\
&= \frac{1}{\cos x} \cdot \lim_{\frac{h}{2} \rightarrow 0} \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \cdot \lim_{h \rightarrow 0} \frac{\sin\left(\frac{2x+h}{2}\right)}{\cos(x+h)} \\
&= \frac{1}{\cos x} \cdot 1 \cdot \frac{\sin x}{\cos x} \\
&= \sec x \tan x
\end{aligned}$$

(iii) Let  $f(x) = 5 \sec x + 4 \cos x$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{5 \sec(x+h) + 4 \cos(x+h) - [5 \sec x + 4 \cos x]}{h} \\
&= 5 \lim_{h \rightarrow 0} \frac{[\sec(x+h) - \sec x]}{h} + 4 \lim_{h \rightarrow 0} \frac{[\cos(x+h) - \cos x]}{h} \\
&= 5 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\cos(x+h)} - \frac{1}{\cos x} \right] + 4 \lim_{h \rightarrow 0} \frac{1}{h} [\cos(x+h) - \cos x] \\
&= 5 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\cos x - \cos(x+h)}{\cos x \cos(x+h)} \right] + 4 \lim_{h \rightarrow 0} \frac{1}{h} [\cos x \cos h - \sin x \sin h - \cos x] \\
&= \frac{5}{\cos x} \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2 \sin\left(\frac{x+x+h}{2}\right) \sin\left(\frac{x-x-h}{2}\right)}{\cos(x+h)} \right] + 4 \lim_{h \rightarrow 0} \frac{1}{h} [-\cos x(1-\cos h) - \sin x \sin h] \\
&= \frac{5}{\cos x} \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2 \sin\left(\frac{2x+h}{2}\right) \sin\left(-\frac{h}{2}\right)}{\cos(x+h)} \right] + 4 \left[ -\cos x \lim_{h \rightarrow 0} \frac{(1-\cos h)}{h} - \sin x \lim_{h \rightarrow 0} \frac{\sin h}{h} \right] \\
&= \frac{5}{\cos x} \lim_{h \rightarrow 0} \left[ \frac{\sin\left(\frac{2x+h}{2}\right) \cdot \frac{\sin\left(\frac{h}{2}\right)}{\frac{h}{2}}}{\cos(x+h)} \right] + 4 [(-\cos x)(0) - (\sin x).1] \\
&= \frac{5}{\cos x} \left[ \lim_{h \rightarrow 0} \frac{\sin\left(\frac{2x+h}{2}\right)}{\cos(x+h)} \cdot \lim_{h \rightarrow 0} \frac{\sin\left(\frac{h}{2}\right)}{\frac{h}{2}} \right] - 4 \sin x \\
&= \frac{5}{\cos x} \cdot \frac{\sin x}{\cos x} \cdot 1 - 4 \sin x \\
&= 5 \sec x \tan x - 4 \sin x
\end{aligned}$$

(iv) Let  $f(x) = \operatorname{cosec} x$ . Accordingly, from the first principle,

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{1}{h} [\csc(x+h) - \csc x]$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\sin(x+h)} - \frac{1}{\sin x} \right]$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin x - \sin(x+h)}{\sin(x+h)\sin x} \right]$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{2\cos\left(\frac{x+x+h}{2}\right) \cdot \sin\left(\frac{x-x-h}{2}\right)}{\sin(x+h)\sin x} \right]$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{2\cos\left(\frac{2x+h}{2}\right) \sin\left(-\frac{h}{2}\right)}{\sin(x+h)\sin x} \right]$$

$$= \lim_{h \rightarrow 0} \frac{-\cos\left(\frac{2x+h}{2}\right) \cdot \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)}}{\sin(x+h)\sin x}$$

$$= \lim_{h \rightarrow 0} \left( \frac{-\cos\left(\frac{2x+h}{2}\right)}{\sin(x+h)\sin x} \right) \cdot \lim_{\frac{h}{2} \rightarrow 0} \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)}$$

$$= \left( \frac{-\cos x}{\sin x \sin x} \right) \cdot 1$$

$$= -\csc x \cot x$$

(v) Let  $f(x) = 3\cot x + 5\csc x$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{3 \cot(x+h) + 5 \operatorname{cosec}(x+h) - 3 \cot x - 5 \operatorname{cosec} x}{h} \\
&= 3 \lim_{h \rightarrow 0} \frac{1}{h} [\cot(x+h) - \cot x] + 5 \lim_{h \rightarrow 0} \frac{1}{h} [\operatorname{cosec}(x+h) - \operatorname{cosec} x]
\end{aligned} \quad \dots(1)$$

Now,  $\lim_{h \rightarrow 0} \frac{1}{h} [\cot(x+h) - \cot x]$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\cos(x+h)}{\sin(x+h)} - \frac{\cos x}{\sin x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\cos(x+h)\sin x - \cos x \sin(x+h)}{\sin x \sin(x+h)} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x-x-h)}{\sin x \sin(x+h)} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(-h)}{\sin x \sin(x+h)} \right] \\
&= - \left( \lim_{h \rightarrow 0} \frac{\sin h}{h} \right) \cdot \left( \lim_{h \rightarrow 0} \frac{1}{\sin x \cdot \sin(x+h)} \right) \\
&= -1 \cdot \frac{1}{\sin x \cdot \sin(x+0)} = \frac{-1}{\sin^2 x} = -\operatorname{cosec}^2 x
\end{aligned} \quad \dots(2)$$

$$\begin{aligned}
& \lim_{h \rightarrow 0} \frac{1}{h} [\cosec(x+h) - \cosecx] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\sin(x+h)} - \frac{1}{\sin x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin x - \sin(x+h)}{\sin(x+h)\sin x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{2 \cos\left(\frac{x+x+h}{2}\right) \cdot \sin\left(\frac{x-x-h}{2}\right)}{\sin(x+h)\sin x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{2 \cos\left(\frac{2x+h}{2}\right) \sin\left(-\frac{h}{2}\right)}{\sin(x+h)\sin x} \right] \\
&\quad - \cos\left(\frac{2x+h}{2}\right) \cdot \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \\
&= \lim_{h \rightarrow 0} \frac{-\cos\left(\frac{2x+h}{2}\right)}{\sin(x+h)\sin x} \cdot \lim_{\frac{h}{2} \rightarrow 0} \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \\
&= \left( \frac{-\cos x}{\sin x \sin x} \right) \cdot 1 \\
&= -\cosecx \cot x
\end{aligned} \tag{3}$$

From (1), (2), and (3), we obtain

$$f'(x) = -3\cosec^2 x - 5\cosec x \cot x$$

(vi) Let  $f(x) = 5\sin x - 6\cos x + 7$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} [5 \sin(x+h) - 6 \cos(x+h) + 7 - 5 \sin x + 6 \cos x - 7] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} [5 \{\sin(x+h) - \sin x\} - 6 \{\cos(x+h) - \cos x\}] \\
&= 5 \lim_{h \rightarrow 0} \frac{1}{h} [\sin(x+h) - \sin x] - 6 \lim_{h \rightarrow 0} \frac{1}{h} [\cos(x+h) - \cos x] \\
&= 5 \lim_{h \rightarrow 0} \frac{1}{h} \left[ 2 \cos\left(\frac{x+h+x}{2}\right) \sin\left(\frac{x+h-x}{2}\right) \right] - 6 \lim_{h \rightarrow 0} \frac{\cos x \cos h - \sin x \sin h - \cos x}{h} \\
&= 5 \lim_{h \rightarrow 0} \frac{1}{h} \left[ 2 \cos\left(\frac{2x+h}{2}\right) \sin\frac{h}{2} \right] - 6 \lim_{h \rightarrow 0} \left[ \frac{-\cos x(1-\cos h) - \sin x \sin h}{h} \right] \\
&= 5 \lim_{h \rightarrow 0} \left( \cos\left(\frac{2x+h}{2}\right) \frac{\sin \frac{h}{2}}{\frac{h}{2}} \right) - 6 \lim_{h \rightarrow 0} \left[ \frac{-\cos x(1-\cos h)}{h} - \frac{\sin x \sin h}{h} \right] \\
&= 5 \left[ \lim_{h \rightarrow 0} \cos\left(\frac{2x+h}{2}\right) \right] \left[ \lim_{\substack{h \rightarrow 0 \\ \frac{h}{2}}} \frac{\sin \frac{h}{2}}{\frac{h}{2}} \right] - 6 \left[ (-\cos x) \left( \lim_{h \rightarrow 0} \frac{1-\cos h}{h} \right) - \sin x \lim_{h \rightarrow 0} \left( \frac{\sin h}{h} \right) \right] \\
&= 5 \cos x \cdot 1 - 6 [(-\cos x) \cdot (0) - \sin x \cdot 1] \\
&= 5 \cos x + 6 \sin x
\end{aligned}$$

(vii) Let  $f(x) = 2 \tan x - 7 \sec x$ . Accordingly, from the first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} [2 \tan(x+h) - 7 \sec(x+h) - 2 \tan x + 7 \sec x] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} [2 \{\tan(x+h) - \tan x\} - 7 \{\sec(x+h) - \sec x\}] \\
&= 2 \lim_{h \rightarrow 0} \frac{1}{h} [\tan(x+h) - \tan x] - 7 \lim_{h \rightarrow 0} \frac{1}{h} [\sec(x+h) - \sec x] \\
&= 2 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x} \right] - 7 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\cos(x+h)} - \frac{1}{\cos x} \right] \\
&= 2 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h)\cos x - \sin x \cos(x+h)}{\cos x \cos(x+h)} \right] - 7 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\cos x - \cos(x+h)}{\cos x \cos(x+h)} \right] \\
&= 2 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h-x)}{\cos x \cos(x+h)} \right] - 7 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2 \sin\left(\frac{x+x+h}{2}\right) \sin\left(\frac{x-x-h}{2}\right)}{\cos x \cos(x+h)} \right] \\
&= 2 \lim_{h \rightarrow 0} \left[ \left( \frac{\sin h}{h} \right) \frac{1}{\cos x \cos(x+h)} \right] - 7 \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2 \sin\left(\frac{2x+h}{2}\right) \sin\left(-\frac{h}{2}\right)}{\cos x \cos(x+h)} \right] \\
&= 2 \left( \lim_{h \rightarrow 0} \frac{\sin h}{h} \right) \left( \lim_{h \rightarrow 0} \frac{1}{\cos x \cos(x+h)} \right) - 7 \left( \lim_{\frac{h}{2} \rightarrow 0} \frac{\sin \frac{h}{2}}{\frac{h}{2}} \right) \left( \lim_{h \rightarrow 0} \frac{\sin\left(\frac{2x+h}{2}\right)}{\cos x \cos(x+h)} \right) \\
&= 2 \cdot 1 \cdot \frac{1}{\cos x \cos x} - 7 \cdot 1 \left( \frac{\sin x}{\cos x \cos x} \right) \\
&= 2 \sec^2 x - 7 \sec x \tan x
\end{aligned}$$


---

Page No 317:

Question 1:

Find the derivative of the following functions from first principle:

(i)  $-x$  (ii)  $(-x)^{-1}$  (iii)  $\sin(x+1)$

(iv)  $\cos\left(x - \frac{\pi}{8}\right)$

Answer:

(i) Let  $f(x) = -x$ . Accordingly,  $f(x+h) = -(x+h)$

By first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{-(x+h) - (-x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{-x - h + x}{h} \\
&= \lim_{h \rightarrow 0} \frac{-h}{h} \\
&= \lim_{h \rightarrow 0} (-1) = -1
\end{aligned}$$

(ii) Let  $f(x) = (-x)^{-1} = \frac{1}{-x} = \frac{-1}{x}$ . Accordingly,  $f(x+h) = \frac{-1}{(x+h)}$

By first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-1}{x+h} - \left( \frac{-1}{x} \right) \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-1}{x+h} + \frac{1}{x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-x + (x+h)}{x(x+h)} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-x + x + h}{x(x+h)} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{h}{x(x+h)} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{x(x+h)} \\
&= \frac{1}{x \cdot x} = \frac{1}{x^2}
\end{aligned}$$

(iii) Let  $f(x) = \sin(x+1)$ . Accordingly,  $f(x+h) = \sin(x+h+1)$

By first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} [\sin(x+h+1) - \sin(x+1)] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ 2 \cos\left(\frac{x+h+1+x+1}{2}\right) \sin\left(\frac{x+h+1-x-1}{2}\right) \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ 2 \cos\left(\frac{2x+h+2}{2}\right) \sin\left(\frac{h}{2}\right) \right] \\
&= \lim_{h \rightarrow 0} \left[ \cos\left(\frac{2x+h+2}{2}\right) \cdot \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \right] \\
&= \lim_{h \rightarrow 0} \cos\left(\frac{2x+h+2}{2}\right) \cdot \lim_{\substack{h \rightarrow 0 \\ 2}} \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \quad \left[ \text{As } h \rightarrow 0 \Rightarrow \frac{h}{2} \rightarrow 0 \right] \\
&= \cos\left(\frac{2x+0+2}{2}\right) \cdot 1 \quad \left[ \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right] \\
&= \cos(x+1)
\end{aligned}$$

(iv) Let  $f(x) = \cos\left(x - \frac{\pi}{8}\right)$ . Accordingly,  $f(x+h) = \cos\left(x+h - \frac{\pi}{8}\right)$

By first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \cos\left(x+h - \frac{\pi}{8}\right) - \cos\left(x - \frac{\pi}{8}\right) \right]
\end{aligned}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ -2 \sin \left( \frac{x+h-\frac{\pi}{8}+x-\frac{\pi}{8}}{2} \right) \sin \left( \frac{x+h-\frac{\pi}{8}-x+\frac{\pi}{8}}{2} \right) \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ -2 \sin \left( \frac{2x+h-\frac{\pi}{4}}{2} \right) \sin \frac{h}{2} \right] \\
&= \lim_{h \rightarrow 0} \left[ -\sin \left( \frac{2x+h-\frac{\pi}{4}}{2} \right) \frac{\sin \left( \frac{h}{2} \right)}{\left( \frac{h}{2} \right)} \right] \\
&= \lim_{h \rightarrow 0} \left[ -\sin \left( \frac{2x+h-\frac{\pi}{4}}{2} \right) \right] \cdot \lim_{\frac{h}{2} \rightarrow 0} \frac{\sin \left( \frac{h}{2} \right)}{\left( \frac{h}{2} \right)} \quad \left[ \text{As } h \rightarrow 0 \Rightarrow \frac{h}{2} \rightarrow 0 \right] \\
&= -\sin \left( \frac{2x+0-\frac{\pi}{4}}{2} \right) \cdot 1 \\
&= -\sin \left( x - \frac{\pi}{8} \right)
\end{aligned}$$


---

Question 2:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):  $(x + a)$

Answer:

Let  $f(x) = x + a$ . Accordingly,  $f(x+h) = x + h + a$

By first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{x+h+a-x-a}{h} \\
&= \lim_{h \rightarrow 0} \left( \frac{h}{h} \right) \\
&= \lim_{h \rightarrow 0} (1) \\
&= 1
\end{aligned}$$


---

Question 3:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):

$$(px+q)\left(\frac{r}{x}+s\right)$$

Answer:

$$\text{Let } f(x) = (px+q)\left(\frac{r}{x}+s\right)$$

By Leibnitz product rule,

$$\begin{aligned} f'(x) &= (px+q)\left(\frac{r}{x}+s\right)' + \left(\frac{r}{x}+s\right)(px+q)' \\ &= (px+q)\left(rx^{-1}+s\right)' + \left(\frac{r}{x}+s\right)(p) \\ &= (px+q)\left(-rx^{-2}\right) + \left(\frac{r}{x}+s\right)p \\ &= (px+q)\left(\frac{-r}{x^2}\right) + \left(\frac{r}{x}+s\right)p \\ &= \frac{-pr}{x} - \frac{qr}{x^2} + \frac{pr}{x} + ps \\ &= ps - \frac{qr}{x^2} \end{aligned}$$


---

Question 4:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):  $(ax+b)(cx+d)^2$

Answer:

$$\text{Let } f(x) = (ax+b)(cx+d)^2$$

By Leibnitz product rule,

$$\begin{aligned} f'(x) &= (ax+b)\frac{d}{dx}(cx+d)^2 + (cx+d)^2\frac{d}{dx}(ax+b) \\ &= (ax+b)\frac{d}{dx}(c^2x^2 + 2cdx + d^2) + (cx+d)^2\frac{d}{dx}(ax+b) \\ &= (ax+b)\left[\frac{d}{dx}(c^2x^2) + \frac{d}{dx}(2cdx) + \frac{d}{dx}d^2\right] + (cx+d)^2\left[\frac{d}{dx}ax + \frac{d}{dx}b\right] \\ &= (ax+b)(2c^2x + 2cd) + (cx+d)^2a \\ &= 2c(ax+b)(cx+d) + a(cx+d)^2 \end{aligned}$$


---

Question 5:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

q, r ands are fixed non-zero constants and m and n are integers):  $\frac{ax+b}{cx+d}$

Answer:

Let  $f(x) = \frac{ax+b}{cx+d}$

By quotient rule,

$$\begin{aligned}f'(x) &= \frac{(cx+d)\frac{d}{dx}(ax+b) - (ax+b)\frac{d}{dx}(cx+d)}{(cx+d)^2} \\&= \frac{(cx+d)(a) - (ax+b)(c)}{(cx+d)^2} \\&= \frac{acx + ad - acx - bc}{(cx+d)^2} \\&= \frac{ad - bc}{(cx+d)^2}\end{aligned}$$

---

Question 6:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r ands are fixed non-zero constants and m and n are integers):

$$\frac{1+\frac{1}{x}}{1-\frac{1}{x}}$$

Answer:

Let  $f(x) = \frac{1+\frac{1}{x}}{1-\frac{1}{x}} = \frac{x+1}{x-1} = \frac{x+1}{x-1}$ , where  $x \neq 0$

By quotient rule,

$$\begin{aligned}
f'(x) &= \frac{(x-1)\frac{d}{dx}(x+1) - (x+1)\frac{d}{dx}(x-1)}{(x-1)^2}, \quad x \neq 0, 1 \\
&= \frac{(x-1)(1) - (x+1)(1)}{(x-1)^2}, \quad x \neq 0, 1 \\
&= \frac{x-1-x-1}{(x-1)^2}, \quad x \neq 0, 1 \\
&= \frac{-2}{(x-1)^2}, \quad x \neq 0, 1
\end{aligned}$$


---

Question 7:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

q, r ands are fixed non-zero constants and m and n are integers):  $\frac{1}{ax^2 + bx + c}$

Answer:

$$\text{Let } f(x) = \frac{1}{ax^2 + bx + c}$$

By quotient rule,

$$\begin{aligned}
f'(x) &= \frac{(ax^2 + bx + c)\frac{d}{dx}(1) - \frac{d}{dx}(ax^2 + bx + c)}{(ax^2 + bx + c)^2} \\
&= \frac{(ax^2 + bx + c)(0) - (2ax + b)}{(ax^2 + bx + c)^2} \\
&= \frac{-(2ax + b)}{(ax^2 + bx + c)^2}
\end{aligned}$$


---

Question 8:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

q, r ands are fixed non-zero constants and m and n are integers):  $\frac{ax+b}{px^2 + qx + r}$

Answer:

$$\text{Let } f(x) = \frac{ax+b}{px^2 + qx + r}$$

By quotient rule,

$$\begin{aligned}
f'(x) &= \frac{(px^2 + qx + r)\frac{d}{dx}(ax + b) - (ax + b)\frac{d}{dx}(px^2 + qx + r)}{(px^2 + qx + r)^2} \\
&= \frac{(px^2 + qx + r)(a) - (ax + b)(2px + q)}{(px^2 + qx + r)^2} \\
&= \frac{apx^2 + aqx + ar - 2apx^2 - aqx - 2bpx - bq}{(px^2 + qx + r)^2} \\
&= \frac{-apx^2 - 2bpx + ar - bq}{(px^2 + qx + r)^2}
\end{aligned}$$


---

Question 9:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

$$q, r \text{ and } m \text{ and } n \text{ are fixed non-zero constants and } m \text{ and } n \text{ are integers): } \frac{px^m + qx^n}{ax + b}$$

Answer:

$$\text{Let } f(x) = \frac{px^m + qx^n}{ax + b}$$

By quotient rule,

$$\begin{aligned}
f'(x) &= \frac{(ax + b)\frac{d}{dx}(px^m + qx^n) - (px^m + qx^n)\frac{d}{dx}(ax + b)}{(ax + b)^2} \\
&= \frac{(ax + b)(mpx^{m-1} + nqx^{n-1}) - (px^m + qx^n)(a)}{(ax + b)^2} \\
&= \frac{2apx^m + aqx^{n-1} + 2bpqx^{m-1} + bq - apx^m - aqx^{n-1}}{(ax + b)^2} \\
&= \frac{apx^m + 2bpqx^{m-1} + bq - ar}{(ax + b)^2}
\end{aligned}$$


---

Question 10:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

$$q, r \text{ and } m \text{ and } n \text{ are fixed non-zero constants and } m \text{ and } n \text{ are integers): } \frac{a}{x^4} - \frac{b}{x^2} + \cos x$$

Answer:

$$\text{Let } f(x) = \frac{a}{x^4} - \frac{b}{x^2} + \cos x$$

$$\begin{aligned}f'(x) &= \frac{d}{dx}\left(\frac{a}{x^4}\right) - \frac{d}{dx}\left(\frac{b}{x^2}\right) + \frac{d}{dx}(\cos x) \\&= a\frac{d}{dx}(x^{-4}) - b\frac{d}{dx}(x^{-2}) + \frac{d}{dx}(\cos x) \\&= a(-4x^{-5}) - b(-2x^{-3}) + (-\sin x) \quad \left[ \frac{d}{dx}(x^n) = nx^{n-1} \text{ and } \frac{d}{dx}(\cos x) = -\sin x \right] \\&= \frac{-4a}{x^5} + \frac{2b}{x^3} - \sin x\end{aligned}$$

---

Question 11:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r ands are fixed non-zero constants and m and n are integers):  $4\sqrt{x} - 2$

Answer:

$$\text{Let } f(x) = 4\sqrt{x} - 2$$

$$\begin{aligned}f'(x) &= \frac{d}{dx}(4\sqrt{x} - 2) = \frac{d}{dx}(4\sqrt{x}) - \frac{d}{dx}(2) \\&= 4\frac{d}{dx}\left(x^{\frac{1}{2}}\right) - 0 = 4\left(\frac{1}{2}x^{\frac{1}{2}-1}\right) \\&= \left(2x^{-\frac{1}{2}}\right) = \frac{2}{\sqrt{x}}\end{aligned}$$

---

Question 12:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r ands are fixed non-zero constants and m and n are integers):  $(ax + b)^n$

Answer:

$$\text{Let } f(x) = (ax + b)^n. \text{ Accordingly, } f(x+h) = \{a(x+h) + b\}^n = (ax + ah + b)^n$$

By first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{(ax+ah+b)^n - (ax+b)^n}{h} \\
&= \lim_{h \rightarrow 0} \frac{(ax+b)^n \left(1 + \frac{ah}{ax+b}\right)^n - (ax+b)^n}{h} \\
&= (ax+b)^n \lim_{h \rightarrow 0} \frac{\left(1 + \frac{ah}{ax+b}\right)^n - 1}{h} \\
&= (ax+b)^n \lim_{h \rightarrow 0} \frac{1}{n} \left[ \left\{ 1 + n \left( \frac{ah}{ax+b} \right) + \frac{n(n-1)}{2} \left( \frac{ah}{ax+b} \right)^2 + \dots \right\} - 1 \right] \\
&\quad (\text{Using binomial theorem}) \\
&= (ax+b)^n \lim_{h \rightarrow 0} \frac{1}{h} \left[ n \left( \frac{ah}{ax+b} \right) + \frac{n(n-1)a^2h^2}{2(ax+b)^2} + \dots \right] (\text{Terms containing higher degrees of } h) \\
&= (ax+b)^n \lim_{h \rightarrow 0} \left[ \frac{na}{(ax+b)} + \frac{n(n-1)a^2h}{2(ax+b)^2} + \dots \right] \\
&= (ax+b)^n \left[ \frac{na}{(ax+b)} + 0 \right] \\
&= na \frac{(ax+b)^n}{(ax+b)} \\
&= na(ax+b)^{n-1}
\end{aligned}$$


---

Question 13:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):  $(ax + b)^n (cx + d)^m$

Answer:

$$\text{Let } f(x) = (ax+b)^n (cx+d)^m$$

By Leibnitz product rule,

$$f'(x) = (ax+b)^n \frac{d}{dx}(cx+d)^m + (cx+d)^m \frac{d}{dx}(ax+b)^n \quad \dots(1)$$

$$\text{Now, let } f_1(x) = (cx+d)^m$$

$$f_1(x+h) = (cx+ch+d)^m$$

$$\begin{aligned}
f'_1(x) &= \lim_{h \rightarrow 0} \frac{f_1(x+h) - f_1(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{(cx+ch+d)^m - (cx+d)^m}{h} \\
&= (cx+d)^m \lim_{h \rightarrow 0} \frac{1}{h} \left[ \left( 1 + \frac{ch}{cx+d} \right)^m - 1 \right] \\
&= (cx+d)^m \lim_{h \rightarrow 0} \frac{1}{h} \left[ \left( 1 + \frac{mch}{(cx+d)} + \frac{m(m-1)}{2} \frac{(c^2 h^2)}{(cx+d)^2} + \dots \right) - 1 \right] \\
&= (cx+d)^m \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{mch}{(cx+d)} + \frac{m(m-1)c^2 h^2}{2(cx+d)^2} + \dots \text{ (Terms containing higher degrees of } h \text{)} \right] \\
&= (cx+d)^m \lim_{h \rightarrow 0} \left[ \frac{mc}{(cx+d)} + \frac{m(m-1)c^2 h}{2(cx+d)^2} + \dots \right] \\
&= (cx+d)^m \left[ \frac{mc}{cx+d} + 0 \right] \\
&= \frac{mc(cx+d)^{m-1}}{(cx+d)} \\
&= mc(cx+d)^{m-1}
\end{aligned}$$

$$\frac{d}{dx}(cx+d)^m = mc(cx+d)^{m-1} \quad \dots(2)$$

$$\text{Similarly, } \frac{d}{dx}(ax+b)^n = na(ax+b)^{n-1} \quad \dots(3)$$

Therefore, from (1), (2), and (3), we obtain

$$\begin{aligned}
f'(x) &= (ax+b)^n \left\{ mc(cx+d)^{m-1} \right\} + (cx+d)^m \left\{ na(ax+b)^{n-1} \right\} \\
&= (ax+b)^{n-1} (cx+d)^{m-1} [mc(ax+b) + na(cx+d)]
\end{aligned}$$

Question 14:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):  $\sin(x+a)$

Answer:

$$\text{Let } f(x) = \sin(x+a)$$

$$f(x+h) = \sin(x+h+a)$$

By first principle,

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\sin(x+h+a) - \sin(x+a)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ 2 \cos\left(\frac{x+h+a+x+a}{2}\right) \sin\left(\frac{x+h+a-x-a}{2}\right) \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ 2 \cos\left(\frac{2x+2a+h}{2}\right) \sin\left(\frac{h}{2}\right) \right] \\
&= \lim_{h \rightarrow 0} \left[ \cos\left(\frac{2x+2a+h}{2}\right) \left\{ \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \right\} \right] \\
&= \lim_{h \rightarrow 0} \cos\left(\frac{2x+2a+h}{2}\right) \lim_{\frac{h}{2} \rightarrow 0} \left\{ \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \right\} \quad \left[ \text{As } h \rightarrow 0 \Rightarrow \frac{h}{2} \rightarrow 0 \right] \\
&= \cos\left(\frac{2x+2a}{2}\right) \times 1 \quad \left[ \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right] \\
&= \cos(x+a)
\end{aligned}$$


---

Question 15:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):  $\operatorname{cosec} x \cot x$

Answer:

$$\text{Let } f(x) = \operatorname{cosec} x \cot x$$

By Leibnitz product rule,

$$f'(x) = \operatorname{cosec} x (\cot x)' + \cot x (\operatorname{cosec} x)' \quad \dots(1)$$

$$\text{Let } f_1(x) = \cot x. \text{ Accordingly, } f_1(x+h) = \cot(x+h)$$

By first principle,

$$\begin{aligned}
f_1'(x) &= \lim_{h \rightarrow 0} \frac{f_1(x+h) - f_1(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\cot(x+h) - \cot x}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{\cos(x+h)}{\sin(x+h)} - \frac{\cos x}{\sin x} \right) \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin x \cos(x+h) - \cos x \sin(x+h)}{\sin x \sin(x+h)} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x-x-h)}{\sin x \sin(x+h)} \right] \\
&= \frac{1}{\sin x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(-h)}{\sin(x+h)} \right] \\
&= \frac{-1}{\sin x} \cdot \left( \lim_{h \rightarrow 0} \frac{\sin h}{h} \right) \left( \lim_{h \rightarrow 0} \frac{1}{\sin(x+h)} \right) \\
&= \frac{-1}{\sin x} \cdot 1 \cdot \left( \frac{1}{\sin(x+0)} \right) \\
&= \frac{-1}{\sin^2 x} \\
&= -\operatorname{cosec}^2 x
\end{aligned}$$

$\therefore (\cot x)' = -\operatorname{cosec}^2 x \quad \dots (2)$

Now, let  $f_2(x) = \operatorname{cosec} x$ . Accordingly,  $f_2(x+h) = \operatorname{cosec}(x+h)$

By first principle,

$$\begin{aligned}
f_2'(x) &= \lim_{h \rightarrow 0} \frac{f_2(x+h) - f_2(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} [\operatorname{cosec}(x+h) - \operatorname{cosec} x]
\end{aligned}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\sin(x+h)} - \frac{1}{\sin x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin x - \sin(x+h)}{\sin x \sin(x+h)} \right] \\
&= \frac{1}{\sin x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{2 \cos\left(\frac{x+x+h}{2}\right) \sin\left(\frac{x-x-h}{2}\right)}{\sin(x+h)} \right] \\
&= \frac{1}{\sin x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{2 \cos\left(\frac{2x+h}{2}\right) \sin\left(\frac{-h}{2}\right)}{\sin(x+h)} \right] \\
&= \frac{1}{\sin x} \cdot \lim_{h \rightarrow 0} \left[ \frac{-\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \cdot \frac{\cos\left(\frac{2x+h}{2}\right)}{\sin(x+h)} \right] \\
&= \frac{-1}{\sin x} \cdot \lim_{h \rightarrow 0} \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \cdot \lim_{h \rightarrow 0} \frac{\cos\left(\frac{2x+h}{2}\right)}{\sin(x+h)} \\
&= \frac{-1}{\sin x} \cdot 1 \cdot \frac{\cos\left(\frac{2x+0}{2}\right)}{\sin(x+0)} \\
&= \frac{-1}{\sin x} \cdot \frac{\cos x}{\sin x} \\
&= -\operatorname{cosec} x \cdot \cot x
\end{aligned}$$

$$\therefore (\operatorname{cosec} x)' = -\operatorname{cosec} x \cdot \cot x \quad \dots (3)$$

From (1), (2), and (3), we obtain

$$\begin{aligned}
f'(x) &= \operatorname{cosec} x (-\operatorname{cosec}^2 x) + \cot x (-\operatorname{cosec} x \cot x) \\
&= -\operatorname{cosec}^3 x - \cot^2 x \operatorname{cosec} x
\end{aligned}$$

Question 16:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

q, r and s are fixed non-zero constants and m and n are integers):  $\frac{\cos x}{1+\sin x}$

Answer:

$$\text{Let } f(x) = \frac{\cos x}{1+\sin x}$$

By quotient rule,

$$\begin{aligned}f'(x) &= \frac{(1+\sin x)\frac{d}{dx}(\cos x) - (\cos x)\frac{d}{dx}(1+\sin x)}{(1+\sin x)^2} \\&= \frac{(1+\sin x)(-\sin x) - (\cos x)(\cos x)}{(1+\sin x)^2} \\&= \frac{-\sin x - \sin^2 x - \cos^2 x}{(1+\sin x)^2} \\&= \frac{-\sin x - (\sin^2 x + \cos^2 x)}{(1+\sin x)^2} \\&= \frac{-\sin x - 1}{(1+\sin x)^2} \\&= \frac{-(1+\sin x)}{(1+\sin x)^2} \\&= \frac{-1}{(1+\sin x)}\end{aligned}$$

---

Page No 318:

Question 17:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

q, r and s are fixed non-zero constants and m and n are integers):  $\frac{\sin x + \cos x}{\sin x - \cos x}$

Answer:

Let  $f(x) = \frac{\sin x + \cos x}{\sin x - \cos x}$

By quotient rule,

$$\begin{aligned}
f'(x) &= \frac{(\sin x - \cos x) \frac{d}{dx}(\sin x + \cos x) - (\sin x + \cos x) \frac{d}{dx}(\sin x - \cos x)}{(\sin x - \cos x)^2} \\
&= \frac{(\sin x - \cos x)(\cos x - \sin x) - (\sin x + \cos x)(\cos x + \sin x)}{(\sin x - \cos x)^2} \\
&= \frac{-(\sin x - \cos x)^2 - (\sin x + \cos x)^2}{(\sin x - \cos x)^2} \\
&= \frac{-[\sin^2 x + \cos^2 x - 2\sin x \cos x + \sin^2 x + \cos^2 x + 2\sin x \cos x]}{(\sin x - \cos x)^2} \\
&= \frac{-[1+1]}{(\sin x - \cos x)^2} \\
&= \frac{-2}{(\sin x - \cos x)^2}
\end{aligned}$$


---

Question 18:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

q, r and s are fixed non-zero constants and m and n are integers):  $\frac{\sec x - 1}{\sec x + 1}$

Answer:

$$\text{Let } f(x) = \frac{\sec x - 1}{\sec x + 1}$$

$$f(x) = \frac{\frac{1}{\cos x} - 1}{\frac{1}{\cos x} + 1} = \frac{1 - \cos x}{1 + \cos x}$$

By quotient rule,

$$\begin{aligned}
f'(x) &= \frac{(1+\cos x)\frac{d}{dx}(1-\cos x) - (1-\cos x)\frac{d}{dx}(1+\cos x)}{(1+\cos x)^2} \\
&= \frac{(1+\cos x)(\sin x) - (1-\cos x)(-\sin x)}{(1+\cos x)^2} \\
&= \frac{\sin x + \cos x \sin x + \sin x - \sin x \cos x}{(1+\cos x)^2} \\
&= \frac{2\sin x}{(1+\cos x)^2} \\
&= \frac{2\sin x}{\left(1 + \frac{1}{\sec x}\right)^2} = \frac{2\sin x}{\frac{(\sec x + 1)^2}{\sec^2 x}} \\
&= \frac{2\sin x \sec^2 x}{(\sec x + 1)^2} \\
&= \frac{2\sin x \sec x}{(\sec x + 1)^2} \\
&= \frac{2\sec x \tan x}{(\sec x + 1)^2}
\end{aligned}$$


---

Question 19:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):  $\sin^n x$

Answer:

Let  $y = \sin^n x$ .

Accordingly, for  $n = 1$ ,  $y = \sin x$ .

$$\therefore \frac{dy}{dx} = \cos x, \text{ i.e., } \frac{d}{dx} \sin x = \cos x$$

For  $n = 2$ ,  $y = \sin^2 x$ .

$$\begin{aligned}
\therefore \frac{dy}{dx} &= \frac{d}{dx} (\sin x \sin x) \\
&= (\sin x)' \sin x + \sin x (\sin x)' \quad [\text{By Leibnitz product rule}] \\
&= \cos x \sin x + \sin x \cos x \\
&= 2 \sin x \cos x \quad \dots(1)
\end{aligned}$$

For  $n = 3$ ,  $y = \sin^3 x$ .

$$\begin{aligned}
\therefore \frac{dy}{dx} &= \frac{d}{dx} (\sin x \sin^2 x) \\
&= (\sin x)' \sin^2 x + \sin x (\sin^2 x)' \quad [\text{By Leibnitz product rule}] \\
&= \cos x \sin^2 x + \sin x (2 \sin x \cos x) \quad [\text{Using (1)}] \\
&= \cos x \sin^2 x + 2 \sin^2 x \cos x \\
&= 3 \sin^2 x \cos x
\end{aligned}$$

We assert that  $\frac{d}{dx} (\sin^n x) = n \sin^{(n-1)} x \cos x$

Let our assertion be true for  $n = k$ .

i.e.,  $\frac{d}{dx} (\sin^k x) = k \sin^{(k-1)} x \cos x \quad \dots(2)$

Consider

$$\begin{aligned}
\frac{d}{dx} (\sin^{k+1} x) &= \frac{d}{dx} (\sin x \sin^k x) \\
&= (\sin x)' \sin^k x + \sin x (\sin^k x)' \quad [\text{By Leibnitz product rule}] \\
&= \cos x \sin^k x + \sin x (k \sin^{(k-1)} x \cos x) \quad [\text{Using (2)}] \\
&= \cos x \sin^k x + k \sin^k x \cos x \\
&= (k+1) \sin^k x \cos x
\end{aligned}$$

Thus, our assertion is true for  $n = k + 1$ .

Hence, by mathematical induction,  $\frac{d}{dx} (\sin^n x) = n \sin^{(n-1)} x \cos x$

---

Question 20:

Find the derivative of the following functions (it is to be understood that  $a, b, c, d, p,$

$q, r$  and  $s$  are fixed non-zero constants and  $m$  and  $n$  are integers):  $\frac{a+b \sin x}{c+d \cos x}$

Answer:

Let  $f(x) = \frac{a+b \sin x}{c+d \cos x}$

By quotient rule,

$$\begin{aligned}
f'(x) &= \frac{(c+d \cos x) \frac{d}{dx}(a+b \sin x) - (a+b \sin x) \frac{d}{dx}(c+d \cos x)}{(c+d \cos x)^2} \\
&= \frac{(c+d \cos x)(b \cos x) - (a+b \sin x)(-d \sin x)}{(c+d \cos x)^2} \\
&= \frac{cb \cos x + bd \cos^2 x + ad \sin x + bd \sin^2 x}{(c+d \cos x)^2} \\
&= \frac{bc \cos x + ad \sin x + bd(\cos^2 x + \sin^2 x)}{(c+d \cos x)^2} \\
&= \frac{bc \cos x + ad \sin x + bd}{(c+d \cos x)^2}
\end{aligned}$$


---

Question 21:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

$$\frac{\sin(x+a)}{\cos x} \quad q, r \text{ and } m \text{ and } n \text{ are fixed non-zero constants and } m \text{ and } n \text{ are integers:}$$

Answer:

$$\text{Let } f(x) = \frac{\sin(x+a)}{\cos x}$$

By quotient rule,

$$\begin{aligned}
f'(x) &= \frac{\cos x \frac{d}{dx}[\sin(x+a)] - \sin(x+a) \frac{d}{dx}\cos x}{\cos^2 x} \\
f'(x) &= \frac{\cos x \frac{d}{dx}[\sin(x+a)] - \sin(x+a)(-\sin x)}{\cos^2 x} \quad \dots (i)
\end{aligned}$$

Let  $g(x) = \sin(x+a)$ . Accordingly,  $g(x+h) = \sin(x+h+a)$

By first principle,

$$\begin{aligned}
g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} [\sin(x+h+a) - \sin(x+a)] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ 2 \cos\left(\frac{x+h+a+x+a}{2}\right) \sin\left(\frac{x+h+a-x-a}{2}\right) \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ 2 \cos\left(\frac{2x+2a+h}{2}\right) \sin\left(\frac{h}{2}\right) \right] \\
&= \lim_{h \rightarrow 0} \left[ \cos\left(\frac{2x+2a+h}{2}\right) \left\{ \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \right\} \right] \\
&= \lim_{h \rightarrow 0} \cos\left(\frac{2x+2a+h}{2}\right) \cdot \lim_{\frac{h}{2} \rightarrow 0} \left\{ \frac{\sin\left(\frac{h}{2}\right)}{\left(\frac{h}{2}\right)} \right\} \quad \left[ \text{As } h \rightarrow 0 \Rightarrow \frac{h}{2} \rightarrow 0 \right] \\
&= \left( \cos\frac{2x+2a}{2} \right) \times 1 \quad \left[ \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1 \right] \\
&= \cos(x+a)
\end{aligned}
\tag{... (ii)}$$

From (i) and (ii), we obtain

$$\begin{aligned}
f'(x) &= \frac{\cos x \cdot \cos(x+a) + \sin x \sin(x+a)}{\cos^2 x} \\
&= \frac{\cos(x+a-x)}{\cos^2 x} \\
&= \frac{\cos a}{\cos^2 x}
\end{aligned}$$


---

Question 22:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):  $x^4 (5 \sin x - 3 \cos x)$

Answer:

$$\text{Let } f(x) = x^4 (5 \sin x - 3 \cos x)$$

By product rule,

$$\begin{aligned}
f'(x) &= x^4 \frac{d}{dx}(5 \sin x - 3 \cos x) + (5 \sin x - 3 \cos x) \frac{d}{dx}(x^4) \\
&= x^4 \left[ 5 \frac{d}{dx}(\sin x) - 3 \frac{d}{dx}(\cos x) \right] + (5 \sin x - 3 \cos x) \frac{d}{dx}(x^4) \\
&= x^4 [5 \cos x - 3(-\sin x)] + (5 \sin x - 3 \cos x)(4x^3) \\
&= x^3 [5x \cos x + 3x \sin x + 20 \sin x - 12 \cos x]
\end{aligned}$$


---

Question 23:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r ands are fixed non-zero constants and m and n are integers):  $(x^2 + 1) \cos x$

Answer:

$$\text{Let } f(x) = (x^2 + 1) \cos x$$

By product rule,

$$\begin{aligned}
f'(x) &= (x^2 + 1) \frac{d}{dx}(\cos x) + \cos x \frac{d}{dx}(x^2 + 1) \\
&= (x^2 + 1)(-\sin x) + \cos x(2x) \\
&= -x^2 \sin x - \sin x + 2x \cos x
\end{aligned}$$


---

Question 24:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r ands are fixed non-zero constants and m and n are integers):  $(ax^2 + \sin x)(p + q \cos x)$

Answer:

$$\text{Let } f(x) = (ax^2 + \sin x)(p + q \cos x)$$

By product rule,

$$\begin{aligned}
f'(x) &= (ax^2 + \sin x) \frac{d}{dx}(p + q \cos x) + (p + q \cos x) \frac{d}{dx}(ax^2 + \sin x) \\
&= (ax^2 + \sin x)(-q \sin x) + (p + q \cos x)(2ax + \cos x) \\
&= -q \sin x(ax^2 + \sin x) + (p + q \cos x)(2ax + \cos x)
\end{aligned}$$


---

Question 25:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r ands are fixed non-zero constants and m and n are integers):  $(x + \cos x)(x - \tan x)$

Answer:

$$\text{Let } f(x) = (x + \cos x)(x - \tan x)$$

By product rule,

$$\begin{aligned}f'(x) &= (x + \cos x) \frac{d}{dx}(x - \tan x) + (x - \tan x) \frac{d}{dx}(x + \cos x) \\&= (x + \cos x) \left[ \frac{d}{dx}(x) - \frac{d}{dx}(\tan x) \right] + (x - \tan x)(1 - \sin x) \\&= (x + \cos x) \left[ 1 - \frac{d}{dx} \tan x \right] + (x - \tan x)(1 - \sin x) \quad \dots (\text{i})\end{aligned}$$

$$\text{Let } g(x) = \tan x. \text{ Accordingly, } g(x+h) = \tan(x+h)$$

By first principle,

$$\begin{aligned}g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\&= \lim_{h \rightarrow 0} \left( \frac{\tan(x+h) - \tan x}{h} \right) \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x} \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h)\cos x - \sin x\cos(x+h)}{\cos(x+h)\cos x} \right] \\&= \frac{1}{\cos x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h-x)}{\cos(x+h)} \right] \\&= \frac{1}{\cos x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin h}{\cos(x+h)} \right] \\&= \frac{1}{\cos x} \cdot \left( \lim_{h \rightarrow 0} \frac{\sin h}{h} \right) \cdot \left( \lim_{h \rightarrow 0} \frac{1}{\cos(x+h)} \right) \\&= \frac{1}{\cos x} \cdot 1 \cdot \frac{1}{\cos(x+0)} \\&= \frac{1}{\cos^2 x} \\&= \sec^2 x \quad \dots (\text{ii})\end{aligned}$$

Therefore, from (i) and (ii), we obtain

$$\begin{aligned}f'(x) &= (x + \cos x)(1 - \sec^2 x) + (x - \tan x)(1 - \sin x) \\&= (x + \cos x)(-\tan^2 x) + (x - \tan x)(1 - \sin x) \\&= -\tan^2 x(x + \cos x) + (x - \tan x)(1 - \sin x)\end{aligned}$$

Question 26:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r ands are fixed non-zero constants and m and n are integers):  $\frac{4x+5\sin x}{3x+7\cos x}$

Answer:

Let  $f(x) = \frac{4x+5\sin x}{3x+7\cos x}$

By quotient rule,

$$\begin{aligned} f'(x) &= \frac{(3x+7\cos x)\frac{d}{dx}(4x+5\sin x) - (4x+5\sin x)\frac{d}{dx}(3x+7\cos x)}{(3x+7\cos x)^2} \\ &= \frac{(3x+7\cos x)\left[4\frac{d}{dx}(x)+5\frac{d}{dx}(\sin x)\right] - (4x+5\sin x)\left[3\frac{d}{dx}x+7\frac{d}{dx}\cos x\right]}{(3x+7\cos x)^2} \\ &= \frac{(3x+7\cos x)(4+5\cos x) - (4x+5\sin x)(3-7\sin x)}{(3x+7\cos x)^2} \\ &= \frac{12x+15x\cos x+28\cos x+35\cos^2 x - 12x+28x\sin x-15\sin x+35\sin^2 x}{(3x+7\cos x)^2} \\ &= \frac{15x\cos x+28\cos x+28x\sin x-15\sin x+35(\cos^2 x+\sin^2 x)}{(3x+7\cos x)^2} \\ &= \frac{35+15x\cos x+28\cos x+28x\sin x-15\sin x}{(3x+7\cos x)^2} \end{aligned}$$

Question 27:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r ands are fixed non-zero constants and m and n are integers):

Let  $f(x) = \frac{x^2 \cos\left(\frac{\pi}{4}\right)}{\sin x}$

Answer:

Let  $f(x) = \frac{x^2 \cos\left(\frac{\pi}{4}\right)}{\sin x}$

By quotient rule,

$$\begin{aligned}
f'(x) &= \cos \frac{\pi}{4} \cdot \left[ \frac{\sin x \frac{d}{dx}(x^2) - x^2 \frac{d}{dx}(\sin x)}{\sin^2 x} \right] \\
&= \cos \frac{\pi}{4} \cdot \left[ \frac{\sin x \cdot 2x - x^2 \cos x}{\sin^2 x} \right] \\
&= \frac{x \cos \frac{\pi}{4} [2 \sin x - x \cos x]}{\sin^2 x}
\end{aligned}$$


---

Question 28:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and m and n are integers):  $\frac{x}{1 + \tan x}$

Answer:

$$\text{Let } f(x) = \frac{x}{1 + \tan x}$$

$$\begin{aligned}
f'(x) &= \frac{(1 + \tan x) \frac{d}{dx}(x) - x \frac{d}{dx}(1 + \tan x)}{(1 + \tan x)^2} \\
f'(x) &= \frac{(1 + \tan x) - x \cdot \frac{d}{dx}(1 + \tan x)}{(1 + \tan x)^2} \quad \dots (i)
\end{aligned}$$

Let  $g(x) = 1 + \tan x$ . Accordingly,  $g(x+h) = 1 + \tan(x+h)$ .

By first principle,

$$\begin{aligned}
g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\
&= \lim_{h \rightarrow 0} \left[ \frac{1 + \tan(x+h) - 1 - \tan x}{h} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h)\cos x - \sin x \cos(x+h)}{\cos(x+h)\cos x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h-x)}{\cos(x+h)\cos x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin h}{\cos(x+h)\cos x} \right] \\
&= \left( \lim_{h \rightarrow 0} \frac{\sin h}{h} \right) \cdot \left( \lim_{h \rightarrow 0} \frac{1}{\cos(x+h)\cos x} \right) \\
&= 1 \times \frac{1}{\cos^2 x} = \sec^2 x \\
\Rightarrow \frac{d}{dx}(1 + \tan x) &= \sec^2 x \quad \dots \text{(ii)}
\end{aligned}$$

From (i) and (ii), we obtain

$$f'(x) = \frac{1 + \tan x - x \sec^2 x}{(1 + \tan x)^2}$$


---

Question 29:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p, q, r and s are fixed non-zero constants and m and n are integers):  $(x + \sec x)(x - \tan x)$

Answer:

$$\text{Let } f(x) = (x + \sec x)(x - \tan x)$$

By product rule,

$$\begin{aligned}
f'(x) &= (x + \sec x) \frac{d}{dx}(x - \tan x) + (x - \tan x) \frac{d}{dx}(x + \sec x) \\
&= (x + \sec x) \left[ \frac{d}{dx}(x) - \frac{d}{dx} \tan x \right] + (x - \tan x) \left[ \frac{d}{dx}(x) + \frac{d}{dx} \sec x \right] \\
&= (x + \sec x) \left[ 1 - \frac{d}{dx} \tan x \right] + (x - \tan x) \left[ 1 + \frac{d}{dx} \sec x \right]
\end{aligned} \quad \dots \text{(i)}$$

Let  $f_1(x) = \tan x$ ,  $f_2(x) = \sec x$

Accordingly,  $f_1(x+h) = \tan(x+h)$  and  $f_2(x+h) = \sec(x+h)$

$$\begin{aligned}
f'_1(x) &= \lim_{h \rightarrow 0} \left( \frac{f_1(x+h) - f_1(x)}{h} \right) \\
&= \lim_{h \rightarrow 0} \left( \frac{\tan(x+h) - \tan x}{h} \right) \\
&= \lim_{h \rightarrow 0} \left[ \frac{\tan(x+h) - \tan x}{h} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h)}{\cos(x+h)} - \frac{\sin x}{\cos x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h)\cos x - \sin x\cos(x+h)}{\cos(x+h)\cos x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin(x+h-x)}{\cos(x+h)\cos x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\sin h}{\cos(x+h)\cos x} \right] \\
&= \left( \lim_{h \rightarrow 0} \frac{\sin h}{h} \right) \cdot \left( \lim_{h \rightarrow 0} \frac{1}{\cos(x+h)\cos x} \right) \\
&= 1 \times \frac{1}{\cos^2 x} = \sec^2 x
\end{aligned}$$

$$\Rightarrow \frac{d}{dx} \tan x = \sec^2 x \quad \dots \text{(ii)}$$

$$\begin{aligned}
f'_2(x) &= \lim_{h \rightarrow 0} \left( \frac{f_2(x+h) - f_2(x)}{h} \right) \\
&= \lim_{h \rightarrow 0} \left( \frac{\sec(x+h) - \sec x}{h} \right) \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\cos(x+h)} - \frac{1}{\cos x} \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{\cos x - \cos(x+h)}{\cos(x+h)\cos x} \right] \\
&= \frac{1}{\cos x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2 \sin\left(\frac{x+x+h}{2}\right) \cdot \sin\left(\frac{x-x-h}{2}\right)}{\cos(x+h)} \right] \\
&= \frac{1}{\cos x} \cdot \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{-2 \sin\left(\frac{2x+h}{2}\right) \cdot \sin\left(\frac{-h}{2}\right)}{\cos(x+h)} \right] \\
&= \frac{1}{\cos x} \cdot \lim_{h \rightarrow 0} \left[ \frac{\sin\left(\frac{2x+h}{2}\right) \left\{ \frac{\sin\left(\frac{h}{2}\right)}{\frac{h}{2}} \right\}}{\cos(x+h)} \right] \\
&= \sec x \cdot \frac{\left\{ \lim_{h \rightarrow 0} \sin\left(\frac{2x+h}{2}\right) \right\} \left\{ \lim_{\frac{h}{2} \rightarrow 0} \frac{\sin\left(\frac{h}{2}\right)}{\frac{h}{2}} \right\}}{\lim_{h \rightarrow 0} \cos(x+h)} \\
&= \sec x \cdot \frac{\sin x \cdot 1}{\cos x} \\
&\Rightarrow \frac{d}{dx} \sec x = \sec x \tan x
\end{aligned}
\quad . \quad .. \text{ (iii)}$$

From (i), (ii), and (iii), we obtain

$$f'(x) = (x + \sec x)(1 - \sec^2 x) + (x - \tan x)(1 + \sec x \tan x)$$

Question 30:

Find the derivative of the following functions (it is to be understood that a, b, c, d, p,

q, r and s are fixed non-zero constants and m and n are integers):  $\frac{x}{\sin^n x}$

Answer:

Let  $f(x) = \frac{x}{\sin^n x}$

By quotient rule,

$$f'(x) = \frac{\sin^n x \frac{d}{dx}x - x \frac{d}{dx}\sin^n x}{\sin^{2n} x}$$

It can be easily shown that  $\frac{d}{dx}\sin^n x = n\sin^{n-1} x \cos x$

Therefore,

$$\begin{aligned} f'(x) &= \frac{\sin^n x \frac{d}{dx}x - x \frac{d}{dx}\sin^n x}{\sin^{2n} x} \\ &= \frac{\sin^n x \cdot 1 - x(n\sin^{n-1} x \cos x)}{\sin^{2n} x} \\ &= \frac{\sin^{n-1} x (\sin x - nx \cos x)}{\sin^{2n} x} \\ &= \frac{\sin x - nx \cos x}{\sin^{n+1} x} \end{aligned}$$

---