

Short Answer Type Questions

Q1. If $A = \{-1, 2, 3\}$ and $B = \{1, 3\}$, then determine

- (i) $A \times B$ (ii) $B \times C$ (c) $B \times B$ (iv) $A \times A$

Sol: We have $A = \{-1, 2, 3\}$ and $B = \{1, 3\}$

- (i) $A \times B = \{(-1, 1), (-1, 3), (2, 1), (2, 3), (3, 1), (3, 3)\}$
(ii) $B \times A = \{(1, -1), (1, 2), (1, 3), (3, -1), (3, 2), (3, 3)\}$
(iii) $B \times B = \{(1, 1), (1, 3), (3, 1), (3, 3)\}$
(iv) $A \times A = \{(-1, -1), (-1, 2), (-1, 3), (2, -1), (2, 2), (2, 3), (3, -1), (3, 2), (3, 3)\}$

Q2. If $P = \{x : x < 3, x \in N\}$, $Q = \{x : x \leq 2, x \in W\}$. Find $(P \cup Q) \times (P \cap Q)$, where W is the set of whole numbers.

Sol: We have, $P = \{x : x < 3, x \in N\} = \{1, 2\}$

And $Q = \{x : x \leq 2, x \in W\} = \{0, 1, 2\}$

$P \cup Q = \{0, 1, 2\}$ and $P \cap Q = \{1, 2\}$

$$(P \cup Q) \times (P \cap Q) = \{0, 1, 2\} \times \{1, 2\} \\ = \{(0, 1), (0, 2), (1, 1), (1, 2), (2, 1), (2, 2)\}$$

Q3. If $A = \{x : x \in W, x < 2\}$, $B = \{x : x \in N, 1 < x < 5\}$, $C = \{3, 5\}$. Find

- (i) $A \times (B \cap C)$ (ii) $A \times (B \cup C)$

Sol: We have, $A = \{x : x \in W, x < 2\} = \{0, 1\}$;

$B = \{x : x \in N, 1 < x < 5\} = \{2, 3, 4\}$; and $C = \{3, 5\}$

(i) $B \cap C = \{3\}$

$$A \times (B \cap C) = \{0, 1\} \times \{3\} = \{(0, 3), (1, 3)\}$$

(ii) $(B \cup C) = \{2, 3, 4, 5\}$

$$A \times (B \cup C) = \{0, 1\} \times \{2, 3, 4, 5\}$$

$$= \{(0, 2), (0, 3), (0, 4), (0, 5), (1, 2), (1, 3), (1, 4), (1, 5)\}$$

Q4. In each of the following cases, find a and b . (i) $(2a + b, a - b) = (8, 3)$ (ii) $\{a/4, a - 2b\} = \{0, 6 + b\}$

Sol: (i) We have, $(2a + b, a - b) = (8, 3)$

$$\Rightarrow 2a + b = 8 \text{ and } a - b = 3$$

On solving, we get $a = 11/3$ and $b = 2/3$

(ii) We have, $\left(\frac{a}{4}, a - 2b\right) = (0, 6 + b)$

$$\Rightarrow \frac{a}{4} = 0 \Rightarrow a = 0$$

$$\text{and } a - 2b = 6 + b$$

$$\Rightarrow 0 - 2b = 6 + b$$

$$\Rightarrow b = -2$$

$$\therefore a = 0, b = -2$$

Q5. Given $A = \{1, 2, 3, 4, 5\}$, $S = \{(x, y) : x \in A, y \in A\}$. Find the ordered pairs which satisfy the conditions given below

$$x+y=5 \quad (\text{ii}) \quad x+y<5 \quad (\text{iii}) \quad x+y>8$$

Sol: We have, $A = \{1, 2, 3, 4, 5\}$, $S = \{(x, y) : x \in A, y \in A\}$

(i) The set of ordered pairs satisfying $x + y = 5$ is $\{(1, 4), (2, 3), (3, 2), (4, 1)\}$

(ii) The set of ordered pairs satisfying $x+y < 5$ is $\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (3, 1)\}$

(iii) The set of ordered pairs satisfying $x+y > 8$ is $\{(4, 5), (5, 4), (5, 5)\}$.

Q6. Given $R = \{(x, y) : x, y \in W, x^2 + y^2 = 25\}$. Find the domain and range of R

Sol: We have, $R = \{(x, y) : x, y \in W, x^2 + y^2 = 25\}$

$$= \{(0, 5), (3, 4), (4, 3), (5, 0)\}$$

Domain of R = Set of first element of ordered pairs in $R = \{0, 3, 4, 5\}$

Range of R = Set of second element of ordered pairs in $R = \{5, 4, 3, 0\}$

Q7. If $R_1 = \{(x, y) | y = 2x + 7, \text{ where } x \in R \text{ and } -5 \leq x \leq 5\}$ is a relation. Then find the domain and range of R_1 .

Sol: We have, $R_1 = \{(x, y) | y = 2x + 7, \text{ where } x \in R \text{ and } -5 \leq x \leq 5\}$

Domain of $R_1 = \{-5 \leq x \leq 5, x \in R\} = [-5, 5]$

$$x \in [-5, 5]$$

$$\Rightarrow 2x \in [-10, 10]$$

$$\Rightarrow 2x + 7 \in [-3, 17]$$

Range is $[-3, 17]$

Q8. If $R_2 = \{(x, y) | x \text{ and } y \text{ are integers and } x^2 + y^2 = 64\}$ is a relation. Then find R_2

Sol: We have, $R_2 = \{(x, y) | x \text{ and } y \text{ are integers and } x^2 + y^2 = 64\}$

Clearly, $x^2 = 0$ and $y^2 = 64$ or $x^2 = 64$ and $y^2 = 0$

$x = 0$ and $y = \pm 8$

or $x = \pm 8$ and $y = 0$

$$R_2 = \{(0, 8), (0, -8), (8, 0), (-8, 0)\}$$

Q9. If $R_3 = \{(x, |x|) \mid x \text{ is a real number}\}$ is a relation. Then find domain and range

Sol: We have, $R_3 = \{(x, |x|) \mid x \text{ is a real number}\}$

Clearly, domain of $R_3 = \mathbb{R}$

Now, $x \in \mathbb{R}$ and $|x| \geq 0$.

Range of R_3 is $[0, \infty)$

Q10. Is the given relation a function? Give reasons for your answer.

(i) $h = \{(4, 6), (3, 9), (-11, 6), (3, 11)\}$

(ii) $f = \{(x, x) \mid x \text{ is a real number}\}$

(iii) $g = \{(n, 1/n) \mid n \text{ is a positive integer}\}$

(iv) $s = \{(n, n^2) \mid n \text{ is a positive integer}\}$

(v) $t = \{(x, 3) \mid x \text{ is a real number}\}$

Sol: (i) We have, $h = \{(4, 6), (3, 9), (-11, 6), (3, 11)\}$.

Since pre-image 3 has two images 9 and 11, it is not a function.

(ii) We have, $f = \{(x, x) \mid x \text{ is a real number}\}$

Since every element in the domain has unique image, it is a function.

(iii) We have, $g = \{(n, 1/n) \mid n \text{ is a positive integer}\}$

For n , it is a positive integer and $1/n$ is unique and distinct. Therefore, every element in the domain has unique image. So, it is a function.

(iv) We have, $s = \{(n, n^2) \mid n \text{ is a positive integer}\}$

Since the square of any positive integer is unique, every element in the domain has unique image. Hence, it is a function.

(v) We have, $t = \{(x, 3) \mid x \text{ is a real number}\}$.

Since every element in the domain has the image 3, it is a constant function.

Q11. If f and g are real functions defined by $f(x) = x^2 + 7$ and $g(x) = 3x + 5$, find each of the following

(i) $f(3) + g(-5)$

(ii) $f(1/2) \times g(14)$

(iii) $f(-2) + g(-1)$

(iv) $f(t) - f(-2)$

(v) $\frac{f(t) - f(5)}{t - 5}$, if $t \neq 5$

Sol. Given that, f and g are real functions defined by $f(x) = x^2 + 7$ and $g(x) = 3x + 5$.

$$(i) f(3) = (3)^2 + 7 = 9 + 7 = 16$$

$$\text{and } g(-5) = 3(-5) + 5 = -15 + 5 = -10$$

$$\therefore f(3) + g(-5) = 16 - 10 = 6$$

$$(ii) f(1/2) = (1/2)^2 + 7 = (1/4) + 7 = 29/4$$

$$\text{and } g(14) = 3(14) + 5 = 42 + 5 = 47$$

$$\therefore f(1/2) \times g(14) = (29/4) \times 47 = 1363/4$$

$$(iii) f(-2) = (-2)^2 + 7 = 4 + 7 = 11$$

$$\text{and } g(-1) = 3(-1) + 5 = -3 + 5 = 2$$

$$\therefore f(-2) + g(-1) = 11 + 2 = 13$$

$$(iv) f(t) = t^2 + 7 \text{ and } f(-2) = (-2)^2 + 7 = 4 + 7 = 11$$

$$\therefore f(t) - f(-2) = t^2 + 7 - 11 = t^2 - 4$$

$$(v) f(t) = t^2 + 7 \text{ and } f(5) = 5^2 + 7 = 25 + 7 = 32$$

$$\therefore \frac{f(t) - f(5)}{t - 5}, \text{ if } t \neq 5$$

$$= \frac{t^2 + 7 - 32}{t - 5}$$

$$= \frac{t^2 - 25}{t - 5} = \frac{(t - 5)(t + 5)}{(t - 5)} = t + 5 \quad [\because t \neq 5]$$

Q12. Let f and g be real functions defined by $f(x) = 2x + 1$ and $g(x) = 4x - 7$.

(i) For what real numbers x , $f(x) = g(x)$?

(ii) For what real numbers x , $f(x) < g(x)$?

Sol: We have, $f(x) = 2x + 1$ and $g(x) = 4x - 7$

(i) Now $f(x) = g(x)$

$$\Rightarrow 2x + 1 = 4x - 7$$

$$\Rightarrow 2x = 8 \Rightarrow x = 4$$

(ii) $f(x) < g(x)$

$$\Rightarrow 2x + 1 < 4x - 7$$

$$\Rightarrow 8 < 2x$$

$$\Rightarrow x > 4$$

Q13. If f and g are two real valued functions defined as $f(x) = 2x + 1$, $g(x) = x^2 + 1$, then find.

- (i) $f+g$ (ii) $f-g$ (iii) fg (iv) $\frac{f}{g}$

Sol. We have, $f(x) = 2x + 1$ and $g(x) = x^2 + 1$

$$\begin{aligned} \text{(i)} \quad (f+g)(x) &= f(x) + g(x) \\ &= 2x + 1 + x^2 + 1 = x^2 + 2x + 2 \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad (f-g)(x) &= f(x) - g(x) \\ &= (2x + 1) - (x^2 + 1) = 2x + 1 - x^2 - 1 = 2x - x^2 \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad (fg)(x) &= f(x) \cdot g(x) \\ &= (2x + 1)(x^2 + 1) = 2x^3 + 2x + x^2 + 1 = 2x^3 + x^2 + 2x + 1 \end{aligned}$$

$$\text{(iv)} \quad \frac{f}{g}(x) = \frac{f(x)}{g(x)} = \frac{2x + 1}{x^2 + 1}$$

Q14. Express the following functions as set of ordered pairs and determine their range.

$f: X \rightarrow R, f(x) = x^3 + 1$, where $X = \{-1, 0, 3, 9, 7\}$

Sol: We have, $f: X \rightarrow R, f(x) = x^3 + 1$.

Where $X = \{-1, 0, 3, 9, 7\}$

$$\text{Now } f(-1) = (-1)^3 + 1 = -1 + 1 = 0$$

$$f(0) = (0)^3 + 0 = 0 + 0 = 0$$

$$f(3) = (3)^3 + 1 = 27 + 1 = 28$$

$$f(9) = (9)^3 + 1 = 729 + 1 = 730$$

$$f(7) = (7)^3 + 1 = 343 + 1 = 344$$

$$f = \{(-1, 0), (0, 1), (3, 28), (9, 730), (7, 344)\}$$

$$\text{Range of } f = \{0, 1, 28, 730, 344\}$$

Q15. Find the values of x for which the functions $f(x) = 3x^2 - 1$ and $g(x) = 3+x$ are equal.

Sol: $f(x) = g(x)$

$$\Rightarrow 3x^2 - 1 = 3 + x \Rightarrow 3x^2 - x - 4 = 0 \Rightarrow (3x - 4)(x + 1) = 0$$

$$x = -1, 4/3$$

Q16. Is $g = \{(1, 1), (2, 3), (3, 5), (4, 7)\}$ a function? Justify. If this is described by the relation, $g(x) = x +$, then what values should be assigned to and ?

Sol: We have, $g = \{(1, 1), (2, 3), (3, 5), (4, 7)\}$

Since, every element has unique image under g . So, g is a function.

Now, $g(x) = x +$ For $(1, 1)$, $g(1) = a(1) + P$

$$\Rightarrow 1 = + \quad (\text{i})$$

For (2, 3), $g(2) = (2) +$

$$\Rightarrow 3 = 2 + \quad (\text{ii})$$

On solving Eqs. (i) and (ii), we get $= 2, = -1$

$$f(x) = 2x - 1$$

Also, (3, 5) and (4, 7) satisfy the above function.

Q17. Find the domain of each of the following functions given by

$$(i) f(x) = \frac{1}{\sqrt{1 - \cos x}}$$

$$(ii) f(x) = \frac{1}{\sqrt{x + |x|}}$$

$$(iii) f(x) = x|x|$$

$$(iv) f(x) = \frac{x^3 - x + 3}{x^2 - 1}$$

$$(v) f(x) = \frac{3x}{28 - x}$$

Sol. (i) We have, $f(x) = \frac{1}{\sqrt{1 - \cos x}}$

$$\text{Now } -1 \leq \cos x \leq 1$$

$$\Rightarrow -1 \leq -\cos x \leq 1$$

$$\Rightarrow 0 \leq 1 - \cos x \leq 1$$

So, $f(x)$ is defined, if $1 - \cos x \neq 0$

$$\therefore \cos x \neq 1$$

$$\therefore x \neq 2n\pi, n \in Z$$

\therefore Domain of f is $R - \{2n\pi : n \in Z\}$

(ii) We have, $f(x) = \frac{1}{\sqrt{x+|x|}}$

$$\text{If } x > 0, x + |x| = x + x = 2x > 0$$

$$\text{If } x < 0, x + |x| = x - x = 0$$

Clearly, $x = 0$ is not possible.

$$\therefore \text{Domain of } f = R^+$$

(iii) We have, $f(x) = x|x|$

We know that ' x ' and ' $|x|$ ' are defined for all real values.

Clearly, $f(x)$ is defined for all $x \in R$.

$$\therefore \text{Domain of } f = R$$

(iv) We have, $f(x) = \frac{x^3 - x + 3}{x^2 - 1}$

$f(x)$ is not defined, if $x^2 - 1 = 0$

$$\Rightarrow (x-1)(x+1) = 0$$

$$\Rightarrow x = -1, 1$$

$$\therefore \text{Domain of } f = R - \{-1, 1\}$$

(v) We have, $f(x) = \frac{3x}{28-x}$

Clearly, $f(x)$ is not defined, if $28 - x = 0$

$$\Rightarrow x \neq 28$$

$$\therefore \text{Domain of } f = R - \{28\}$$

Q18. Find the range of the following functions given by

(i) $f(x) = \frac{3}{2 - x^2}$

(ii) $f(x) = 1 - |x-2|$

(iii) $f(x) = |x-3|$

(iv) $f(x) = 1 + 3 \cos 2x$

Sol. (i) We have, $f(x) = \frac{3}{2-x^2} = y$ (let)

$$\Rightarrow 2 - x^2 = \frac{3}{y} \Rightarrow x^2 = 2 - \frac{3}{y}$$

Since $x^2 \geq 0$, $2 - \frac{3}{y} \geq 0$

$$\Rightarrow \frac{2y - 3}{y} \geq 0$$

$\Rightarrow 2y - 3 \geq 0$ and $y > 0$ or $2y - 3 \leq 0$ and $y < 0$

$\Rightarrow y \geq 3/2$ or $y < 0$

$\Rightarrow y \in (-\infty, 0) \cup [3/2, \infty)$

\therefore Range of f is $(-\infty, 0) \cup [3/2, \infty)$

(ii) We know that, $|x - 2| \geq 0$

$$\Rightarrow -|x - 2| \leq 0$$

$$\Rightarrow 1 - |x - 2| \leq 1$$

$$\Rightarrow f(x) \leq 1$$

\therefore Range of f is $(-\infty, 1]$

(iii) We know that, $|x - 3| \geq 0$

$$\Rightarrow f(x) \geq 0$$

\therefore Range of $f = [0, \infty)$

(iv) We know that, $-1 \leq \cos 2x \leq 1$

$$\Rightarrow -3 \leq 3 \cos 2x \leq 3$$

$$\Rightarrow -2 \leq 1 + 3 \cos 2x \leq 4$$

$$\Rightarrow -2 \leq f(x) \leq 4$$

\therefore Range of $f = [-2, 4]$

Q19. Redefine the function $f(x) = |x-2| + |2+x|$, $-3 \leq x \leq 3$

$$\text{Sol. } f(x) = \begin{cases} -(x-2) - (2+x), & -3 \leq x < -2 \\ -(x-2) + (2+x), & -2 \leq x < 2 \\ (x-2) + (2+x), & 2 \leq x \leq 3 \end{cases}$$

$$= \begin{cases} -2x, & -3 \leq x < -2 \\ 4, & -2 \leq x < 2 \\ 2x, & 2 \leq x \leq 3 \end{cases}$$

When $-3 \leq x \leq -2$, $4 \leq -2x \leq 6$

When $2 \leq x \leq 3$, $4 \leq 2x \leq 6$

Thus range is $[4, 6]$.

20. If $f(x) = \frac{x}{x+1}$, then show that

$$\text{(i)} \quad f\left(\frac{1}{x}\right) = -f(x) \quad \text{(ii)} \quad f\left(-\frac{1}{x}\right) = \frac{-1}{f(x)}$$

Sol. We have, $f(x) = \frac{x-1}{x+1}$

$$\text{(i)} \quad f\left(\frac{1}{x}\right) = \frac{\frac{1}{x}-1}{\frac{1}{x}+1} = \frac{\frac{1-x}{x}}{\frac{1+x}{x}} = \frac{1-x}{1+x} = -f(x)$$

$$\text{(ii)} \quad f\left(-\frac{1}{x}\right) = \frac{-\frac{1}{x}-1}{-\frac{1}{x}+1} = \frac{-1-x}{-1+x} = \frac{1+x}{1-x} = \frac{-1}{f(x)}$$

$$\therefore \quad f\left(-\frac{1}{x}\right) = -\frac{1}{f(x)}$$

Q21. Let $f(x) = \sqrt{x}$ and $g(x) = x$ be two functions defined in the domain $R^+ \cup \{0\}$. Find

- (i) $(f+g)(x)$
- (ii) $(f-g)(x)$
- (iii) $(fg)(x)$
- (iv) $f/g(x)$

Sol. We have, $f(x) = \sqrt{x}$ and $g(x) = x$ be two function defined in the domain $R^+ \cup \{0\}$

$$(i) (f+g)(x) = f(x) + g(x) = \sqrt{x} + x$$

$$(ii) (f-g)(x) = f(x) - g(x) = \sqrt{x} - x$$

$$(iii) (fg)(x) = f(x).g(x) = \sqrt{x} \cdot x = x^{\frac{3}{2}}$$

$$(iv) \left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} = \frac{\sqrt{x}}{x} = \frac{1}{\sqrt{x}}$$

22. Find the domain and Range of the function $f(x) = \frac{1}{\sqrt{x-5}}$.

Sol. We have, $f(x) = \frac{1}{\sqrt{x-5}}$

Clearly, $f(x)$ is defined, if $x-5 > 0 \Rightarrow x > 5$

Thus, domain of f is $(5, \infty)$.

For $x-5 > 0$, $\sqrt{x-5} > 0$

$$\therefore \frac{1}{\sqrt{x-5}} > 0$$

Hence, range of f is $(0, \infty)$

Q23. If $f(x) = y = ax - b / cx - a$ then prove that $f(y) = x$

Sol. We have, $f(x) = y = \frac{ax - b}{cx - a}$

$$\begin{aligned}\therefore f(y) &= \frac{ay - b}{cy - a} = \frac{a\left(\frac{ax - b}{cx - a}\right) - b}{c\left(\frac{ax - b}{cx - a}\right) - a} \\ &= \frac{a(ax - b) - b(cx - a)}{c(ax - b) - a(cx - a)} \\ &= \frac{a^2x - ab - bcx + ab}{acx - bc - acx + a^2} = \frac{a^2x - bcx}{a^2 - bc} = \frac{x(a^2 - bc)}{(a^2 - bc)} \\ \therefore f(y) &= x\end{aligned}$$

Objective Type Questions

Q24. Let $n(A) = m$, and $n(B) = n$. Then the total number of non-empty relations that can be defined from A to B is

- (a) m^n
- (b) $n^m - 1$
- (c) $mn - 1$
- (d) $2^{mn} - 1$

Sol: (d) We have, $n(A) = m$ and $n(B) = n$

$$n(A \times B) = n(A) \cdot n(B) = mn$$

Total number of relation from A to B = Number of subsets of $A \times B = 2^{mn}$

So, total number of non-empty relations = $2^{mn} - 1$

Q25. If $[x]^2 - 5[x] + 6 = 0$, where $[.]$ denote the greatest integer function, then

- (a) $x \in [3, 4]$
- (b) $x \in (2, 3]$
- (c) $x \in [2, 3]$
- (d) $x \in [2, 4)$

Sol: (d) We have $[x]^2 - 5[x] + 6 = 0 \Rightarrow [(x - 3)([x] - 2)] = 0$

$$\Rightarrow [x] = 2, 3.$$

$$\text{For } [x] = 2, x \in [2, 3)$$

$$\text{For } [x] = 3, x \in [3, 4)$$

$$x \in [2, 3) \cup [3, 4)$$

$$\text{Or } x \in [2, 4)$$

26. Range of $f(x) = \frac{1}{1 - 2 \cos x}$ is

(a) $\left[\frac{1}{3}, 1 \right]$

(b) $\left[-1, \frac{1}{3} \right]$

(c) $(-\infty, -1] \cup \left[\frac{1}{3}, \infty \right)$

(d) $\left[-\frac{1}{3}, 1 \right]$

Sol. (c) We know that, $-1 \leq \cos x \leq 1$

$$\Rightarrow -1 \leq -\cos x \leq 1$$

$$\Rightarrow -2 \leq -2 \cos x \leq 2$$

$$\Rightarrow -1 \leq 1 - 2 \cos x \leq 3$$

Now $f(x) = \frac{1}{1 - 2 \cos x}$ is defined if

$$-1 \leq 1 - 2 \cos x < 0 \text{ or } 0 < 1 - 2 \cos x \leq 3$$

$$\Rightarrow -1 \geq \frac{1}{1 - 2 \cos x} > -\infty \text{ or } \infty > \frac{1}{1 - 2 \cos x} \geq \frac{1}{3}$$

$$\Rightarrow \frac{1}{1 - 2 \cos x} \in (-\infty, -1] \cup \left[\frac{1}{3}, \infty \right)$$

27. Let $f(x) = \sqrt{1+x^2}$, then

- (a) $f(xy) = f(x) \times f(y)$ (b) $f(xy) \geq f(x) \times f(y)$
(c) $f(xy) \leq f(x) \times f(y)$ (d) None of these

Sol. (c) We have, $f(x) = \sqrt{1+x^2}$

$$f(xy) = \sqrt{1+x^2y^2}$$

$$f(x) \cdot f(y) = \sqrt{1+x^2} \cdot \sqrt{1+y^2} = \sqrt{(1+x^2)(1+y^2)} = \sqrt{1+x^2+y^2+x^2y^2}$$

$$\text{Now } \sqrt{1+x^2y^2} \leq \sqrt{1+x^2+y^2+x^2y^2}$$

$$\Rightarrow f(xy) \leq f(x) \times f(y)$$

28. Domain of $\sqrt{a^2 - x^2}$ ($a > 0$) is

- (a) $(-a, a)$ (b) $[-a, a]$ (c) $[0, a]$ (d) $(-a, 0]$

Sol. (b) We have $f(x) = \sqrt{a^2 - x^2}$

Clearly $f(x)$ is defined, if $a^2 - x^2 \geq 0$

$$\Rightarrow x^2 \leq a^2$$

$$\Rightarrow -a \leq x \leq a$$

$[\because a > 0]$

\therefore Domain of f is $[-a, a]$

Q29. If $f(x) = ax + b$, where a and b are integers, $f(-1) = -5$ and $f(3) = 3$, then a and b are equal to

- (a) $a = -3, b = -1$
(b) $a = 2, b = -3$
(c) $a = 0, b = 2$
(d) $a = 2, b = 3$

Sol. (b) We have, $f(x) = ax + b$

$$\therefore f(-1) = a(-1) + b$$

$$\Rightarrow -5 = -a + b \quad (\text{i})$$

$$\text{Also, } f(3) = a(3) + b$$

$$\Rightarrow 3 = 3a + b \quad (\text{ii})$$

On solving Eqs. (i) and (ii), we get

$$a = 2 \text{ and } b = -3$$

30. The domain of the function f defined by $f(x) = \sqrt{4-x} + \frac{1}{\sqrt{x^2-1}}$ is equal to

- (a) $(-\infty, -1) \cup (1, 4]$ (b) $(-\infty, -1] \cup (1, 4]$
(c) $(-\infty, -1) \cup [1, 4]$ (d) $(-\infty, -1) \cup [1, 4)$

Sol. (a) We have, $f(x) = \sqrt{4-x} + \frac{1}{\sqrt{x^2-1}}$

$f(x)$ is defined if $4 - x \geq 0$ and $x^2 - 1 > 0$

$$\Rightarrow x - 4 \leq 0 \text{ and } (x + 1)(x - 1) > 0$$

$$\Rightarrow x \leq 4 \text{ and } (x < -1 \text{ or } x > 1)$$

\therefore Domain of $f = (-\infty, -1) \cup (1, 4]$

31. The domain and range of the real function f defined by $f(x) = \frac{4-x}{x-4}$ is

- (a) Domain = R , Range = $\{-1, 1\}$
 - (b) Domain = $R - \{1\}$, Range = R
 - (c) Domain = $R - \{4\}$, Range = $\{-1\}$
 - (d) Domain = $R - \{-4\}$, Range = $\{-1, 1\}$

Sol. (c) We have, $f(x) = \frac{4-x}{x-4} = -1$, for $x \neq 4$

32. The domain and range of real function f defined by $f(x) = \sqrt{x-1}$ is given by

- (a) Domain = $(1, \infty)$, Range = $(0, \infty)$ (b) Domain = $[1, \infty)$, Range = $(0, \infty)$
(c) Domain = $[1, \infty)$, Range = $[0, \infty)$ (d) Domain = $[1, \infty)$, Range = $[0, \infty)$

Sol. (d) We have, $f(x) = \sqrt{x-1}$

Clearly, $f(x)$ is defined if $x - 1 \geq 0$

$$\Rightarrow x \geq 1$$

$$\therefore \text{Domain of } f = [1, \infty)$$

Now for $x \geq 1$, $x - 1 \geq 0$

$$\Rightarrow \sqrt{x-1} \geq 1$$

$$\Rightarrow \text{Range of } f = [0, \infty)$$

33. The domain of the function f given by $f(x) = \frac{x^2 + 2x + 1}{x^2 - x - 6}$ is

- (a) $R - \{3, -2\}$ (b) $R - \{-3, 2\}$ (c) $R - [3, -2]$ (d) $R - (3, -2)$

Sol. (a) We have, $f(x) = \frac{x^2 + 2x + 1}{x^2 - x - 6}$

$f(x)$ is not defined, if $x^2 - x - 6 = 0$

$$\Rightarrow (x - 3)(x + 2) = 0$$

$$\therefore x = -2, 3$$

$$\therefore \text{Domain of } f = R - \{-2, 3\}$$

- 34.** The domain and range of the function f given by $f(x) = 2 - |x - 5|$ is
 (a) Domain = $R+$, Range = $(-\infty, 1]$ (b) Domain = R , Range = $(-\infty, 2]$
 (c) Domain = R , Range = $(-\infty, 2)$ (d) Domain = $R+$, Range = $(-\infty, 2]$

Sol. (b) We have, $f(x) = 2 - |x - 5|$

Clearly, $f(x)$ is defined for all $x \in R$.

$$\therefore \text{Domain of } f = R$$

$$\text{Now, } |x - 5| \geq 0, \forall x \in R$$

$$\Rightarrow -|x - 5| \leq 0$$

$$\Rightarrow 2 - |x - 5| \leq 2$$

$$\therefore f(x) \leq 2$$

$$\therefore \text{Range of } f = (-\infty, 2]$$

- 35.** The domain for which the functions defined by $f(x) = 3x^2 - 1$ and $g(x) = 3 + x$ are equal is

- (a) $\left\{-1, \frac{4}{3}\right\}$ (b) $\left[-1, \frac{4}{3}\right]$ (c) $\left(-1, -\frac{4}{3}\right)$ (d) $\left[-1, -\frac{4}{3}\right)$

Sol. (a) We have, $f(x) = 3x^2 - 1$ and $g(x) = 3 + x$

$$f(x) = g(x)$$

$$\Rightarrow 3x^2 - 1 = 3 + x \Rightarrow 3x^2 - x - 4 = 0 \Rightarrow (3x - 4)(x + 1) = 0$$

$$\therefore x = -1, \frac{4}{3}$$

Fill in the Blanks Type Questions

Q36. Let f and g be two real functions given by $f = \{(0, 1), (2, 0), (3, -4), (4, 2), (5, 1)\}$

$g = \{(1, 0), (2, 2), (3, -1), (4, 4), (5, 3)\}$ then the domain of $f \times g$ is given by _____.

Sol: We have, $f = \{(0, 1), (2, 0), (3, -4), (4, 2), (5, 1)\}$ and $g = \{(1, 0), (2, 2), (3, 1), (4, 4), (5, 3)\}$

Domain of $f = \{0, 2, 3, 4, 5\}$

And Domain of $g = \{1, 2, 3, 4, 5\}$

Domain of $(f \times g) = (\text{Domain of } f) \cap (\text{Domain of } g) = \{2, 3, 4, 5\}$

Matching Column Type Questions

Q37. Let $f = \{(2,4), (5,6), (8,-1), (10,-3)\}$ and $g = \{(2,5), (7,1), (8,4), (10,13), (11,5)\}$ be two real functions.
Then match the following:

Column I		Column II	
(a)	$f - g$	(i)	$\left\{ \left(2, \frac{4}{5}\right), \left(8, \frac{-1}{4}\right), \left(10, \frac{-3}{13}\right) \right\}$
(b)	$f + g$	(ii)	$\{(2, 20), (8, -4), (10, -39)\}$
(c)	$f \times g$	(iii)	$\{(2, -1), (8, -5), (10, -16)\}$
(d)	$\frac{f}{g}$	(iv)	$\{(2, 9), (8, 3), (10, 10)\}$

Sol. Domain of $f(x)$ is $\{2, 5, 8, 10\}$.

Domain of $g(x)$ is $\{2, 7, 8, 10, 11\}$.

Thus, domain of $f \pm g, f \times g$ and f/g is $\{2, 8, 10\}$.

For function $y = f(x)$, we have $f(2) = 4, f(8) = -1$ and $f(10) = -3$

For function $y = g(x)$, we have $g(2) = 5, g(8) = 4$ and $g(10) = 13$

$$(f - g)(2) = f(2) - g(2) = 4 - 5 = -1$$

$$(f - g)(8) = f(8) - g(8) = -1 - 4 = -5$$

$$(f - g)(10) = f(10) - g(10) = -3 - 13 = -16$$

Thus, $(f - g)(x) = \{(2, -1), (8, -5), (10, -16)\}$

$$(f + g)(2) = f(2) + g(2) = 4 + 5 = 9$$

$$(f + g)(8) = f(8) + g(8) = -1 + 4 = 3$$

$$(f + g)(10) = f(10) + g(10) = -3 + 13 = 10$$

Thus, $(f + g)(x) = \{(2, 9), (8, 3), (10, 10)\}$

$$(f \cdot g)(2) = f(2) \cdot g(2) = 4 \cdot 5 = 20$$

$$(f \cdot g)(8) = f(8) \cdot g(8) = (-1) \cdot 4 = -4$$

$$(f \cdot g)(10) = f(10) \cdot g(10) = (-3) \cdot 13 = -39$$

Thus $(f \cdot g)(x) = \{(2, 20), (8, -4), (10, -39)\}$

$$(f/g)(2) = f(2)/g(2) = 4/5 = 4/5$$

$$(f/g)(8) = f(8)/g(8) = (-1)/4 = -1/4$$

$$(f/g)(10) = f(10)/g(10) = (-3)/13 = -3/13$$

Thus $(f/g)(x) = \{(2, 4/5), (8, -1/4), (10, -3/13)\}$

So, correct matching is: (a) – (iii), (b) – (iv), (c) – (ii) and (d) – (i)

True/False Type Questions

Q38. The ordered pair (5,2) belongs to the relation $R = \{(x,y) : y = x - 5, x, y \in \mathbb{Z}\}$

Sol: False

We have, $R = \{(x, y) : y = x - 5, x, y \in \mathbb{Z}\}$

When $x = 5$, then $y = 5 - 5 = 0$ Hence, (5, 2) does not belong to R.

Q39. If $P = \{1, 2\}$, then $P \times P \times P = \{(1, 1, 1), (2, 2, 2), (1, 2, 2), (2, 1, 1)\}$

Sol: False

We have, $P = \{1, 2\}$ and $n(P) = 2$

$$n(P \times P \times P) = n(P) \times n(P) \times n(P) = 2 \times 2 \times 2$$

= 8 But given $P \times P \times P$ has 4 elements.

Q40. If $A = \{1, 2, 3\}$, $B = \{3, 4\}$ and $C = \{4, 5, 6\}$, then $(A \times B) \cup (A \times C) = \{(1, 3), (1, 4), (1, 5), (1, 6), (2, 3), (2, 4), (2, 5), (2, 6), (3, 3), (3, 4), (3, 5), (3, 6)\}$.

Sol: True

We have. $A = \{1, 2, 3\}$, $B = \{3, 4\}$ and $C = \{4, 5, 6\}$

$A \times B = \{(1, 3), (1, 4), (2, 3), (2, 4), (3, 3), (3, 4)\}$

And $A \times C = \{(1, 4), (1, 5), (1, 6), (2, 4), (2, 5), (2, 6), (3, 4), (3, 5), (3, 6)\}$

$(A \times B) \cup (A \times C) = \{(1, 3), (1, 4), (1, 5), (1, 6), (2, 3), (2, 4), (2, 5), (2, 6), (3, 3), (3, 4), (3, 5), (3, 6)\}$

41. If $(x - 2, y + 5) = \left(-2, \frac{1}{3}\right)$ are two equal ordered pairs, then $x = 4$, $y = \frac{-14}{3}$.

Sol. False

We have, $(x - 2, y + 5) = \left(-2, \frac{1}{3}\right)$

$$\Rightarrow x - 2 = -2, y + 5 = \frac{1}{3}$$

$$\Rightarrow x = 0, y = \frac{-14}{3}$$

Q42. If $A \times B = \{(a, x), (a, y), (b, x), (b, y)\}$, then $M = \{a, b\}$, $B = \{x, y\}$.

Sol: True

We have, $A \times B = \{\{a, x\}, \{a, y\}, \{b, x\}, \{b, y\}\}$

$A = \text{Set of first element of ordered pairs in } A \times B = \{a, b\}$

$B = \text{Set of second element of ordered pairs in } A \times B = \{x, y\}$