

Unit 7 (Algebraic Expression, Identities & Factorisation)

Multiple Choice Questions

Question. 1 The product of a monomial and a binomial is a

- (a) monomial (b) binomial
(c) trinomial (d) None of these

Solution. (b) Monomial consists of only single term and binomial contains two terms. So, the multiplication of a binomial by a monomial will always produce a binomial, whose first term is the product of monomial and the binomial's first term and second term is the product of monomial and the binomial's second term.

Question. 2 In a polynomial, the exponents of the variables are always (a) integers (b) positive integers (c) non-negative integers (d) non-positive integers

Solution. (c) In a polynomial, the exponents of the variables are either positive integers or 0. Constant term C can be written as Cx^0 . We do not consider the expressions as a polynomial which consist of the variables having negative/fractional exponent.

Question. 3 Which of the following is correct?

- (a) $(a - b)^2 = a^2 + 2ab - b^2$ (b) $(a - b)^2 = a^2 - 2ab + b^2$
(c) $(a - b)^2 = a^2 - b^2$ (d) $(a + b)^2 = a^2 + 2ab - b^2$

Solution.

(b) We have,

$$(a - b)^2 = (a - b)(a - b)$$

$$= a(a - b) - b(a - b)$$

$$= a \cdot a - a \cdot b - b \cdot a + b \cdot b$$

$$= a^2 - ab - ab + b^2$$

$$= a^2 - 2ab + b^2$$

$$[\because a \cdot b = b \cdot a]$$

$$\text{and } (a + b)^2 = (a + b)(a + b)$$

$$= a \cdot a + a \cdot b + b \cdot a + b \cdot b$$

$$= a^2 + 2ab + b^2$$

Question. 4 The sum of $-7pq$ and $2pq$ is

(a) $-9pq$ (b) $9pq$

(c) $5pq$ (d) $-5pq$

Solution.

(d) Given, monomials are $-7pq$ and $2pq$.

$$\therefore \text{Their sum} = -7pq + 2pq = (-7 + 2) pq$$

[both monomials consist of like terms, so adding their numerical coefficient]

$$= -5pq$$

Question. 5 If we subtract $-3x^2y^2$ from x^2y^2 , then we get

(a) $-4x^2y^2$

(b) $-2x^2y^2$

(c) $2x^2y^2$

(d) $4x^2y^2$

Solution.

(d) Given, monomials are $-3x^2y^2$ and x^2y^2 . Now, we have to subtract the first one from the second one.

$$\text{i.e. } x^2y^2 - (-3x^2y^2) = x^2y^2 - (-3)x^2y^2$$

$$= x^2y^2 + 3x^2y^2$$

$$= (1 + 3)x^2y^2$$

$$= 4x^2y^2$$

Question. 6 Like term as $4m^3n^2$ is

(a) $4m^2n^2$ (b) $-6m^3n^2$

(c) $6pm^3n^2$ (d) $4m^3n$

Solution. (b) We know that, the like terms contain the same literal factor. So, the like term as $4m^3n^2$, is $-6m^3n^2$, as it contains the same literal factor m^3n^2 .

Question. 7 Which of the following is a binomial?

(a) $7 \times a + a$

(b) $6a^2 + 7b + 2c$

(c) $4a \times 3b \times 2c$

(d) $6(a^2 + b)$

Solution.

(d) Binomials are algebraic expressions consisting of two unlike terms.

$$(a) 7 \times a + a = 7a + a = 8a$$

[monomial]

$$(b) 6a^2 + 7b + 2c$$

[trinomial]

$$(c) 4a \times 3b \times 2c = 24abc$$

[monomial]

$$(d) 6(a^2 + b) = 6a^2 + 6b$$

[binomial]

Question. 8 Sum of $a - b + ab$, $b + c - bc$ and $c - a - ac$ is

(a) $2c + ab - ac - bc$

(b) $2c - ab - ac - bc$

(c) $2c + ab + ac + bc$

(d) $2c - ab + ac + bc$

Solution.

(a) Required sum = $(a - b + ab) + (b + c - bc) + (c - a - ac)$
 $= a - b + ab + b + c - bc + c - a - ac$
 $= 2c + ab - ac - bc$ [adding the like terms and retaining others]

Question. 9 Product of the monomials $4p, -7q^3, -7pq$ is

- (a) $196 p^2q^4$ (b) $196 pq^4$
 (c) $-196 p^2q^4$ (d) $196 p^2q^3$

Solution.

(a) Required product = $4p \times (-7q^3) \times (-7pq)$
 $= 4 \times (-7) \times (-7) p \times q^3 \times pq$ [multiplying the numerical coefficients]
 $= 196 p^2q^4$ [multiplying the literal factors having same variables]

Question. 10 Area of a rectangle with length $4ab$ and breadth $6b^2$ is

- (a) $24a^2b^2$ (b) $24 ab^3$
 (c) $24 ab^2$ (d) $24ab$

Solution.

(b) We know that, area of a rectangle = Length \times Breadth = $4ab \times 6b^2$
 This is the product of two monomials.
 \therefore Area of rectangle = $(4 \times 6) ab \times b^2$
 $= 24 ab^3$

Question. 11 Volume of a rectangular box (cuboid) with length = $2ab$, breadth = $3ac$ and height = $2ac$ is

- (a) $12 a^3bc^2$ (b) $12 a^3bc$
 (c) $12 a^2bc$ (d) $2 ab + 3 ac + 2ac$

Solution.

(a) We know that, volume of a cuboid = Length \times Breadth \times Height = $2ab \times 3ac \times 2ac$
 $= (2 \times 3 \times 2) ab \times ac \times ac = 12 a \times a \times a \times b \times c \times c = 12a^3bc^2$

Question. 12 Product of $6a^2 - 7b + 5ab$ and $2ab$ is

- (a) $12a^3b - 14ab^2 + 10ab$ (b) $12a^3b - 14ab^2 + 10a^2b^2$
 (c) $6a^2 - 7b + 7ab$ (d) $12a^2b - 7ab^2 + 10ab$

Solution.

(b) Required product = $2ab \times (6a^2 - 7b + 5ab)$
 This is the product of a trinomial by a monomial, so we multiply monomial with each term of the trinomial.
 $\therefore 2ab \times (6a^2 - 7b + 5ab) = 2ab \times 6a^2 + 2ab(-7b) + 2ab \times 5ab$
 $= 12a^3b - 14ab^2 + 10a^2b^2$

Question. 13 Square of $3x - 4y$ is

- (a) $9x^2 - 16y^2$ (b) $6x^2 - 8y^2$
 (c) $9x^2 + 16y^2 + 24xy$ (d) $9x^2 + 16y^2 - 24xy$

Solution.

(d) Square of $(3x - 4y)$ will be $(3x - 4y)^2$.

Comparing $(3x - 4y)^2$ with $(a - b)^2$, we get $a = 3x$ and $b = 4y$.

Now, using the identity $(a - b)^2 = a^2 - 2ab + b^2$

$$\begin{aligned}(3x - 4y)^2 &= (3x)^2 - 2 \cdot 3x \cdot 4y + (4y)^2 \\ &= 9x^2 - 24xy + 16y^2\end{aligned}$$

Question. 14 Which of the following are like terms?

(a) $5xyz^2, -3xy^2z$

(b) $-5xyz^2, 7xyz^2$

(c) $5xyz^2, 5x^3yz$

(d) $5xyz^2, x^2y^2z^2$

Solution.

(b) We know that, the terms having same algebraic (literal) factors are called like terms.

(a) $5xyz^2, -3xy^2z$ [unlike terms]

(b) $-5xyz^2, 7xyz^2$ [like terms]

(c) $5xyz^2, 5x^2yz$ [unlike terms]

(d) $5xyz^2, x^2y^2z^2$ [unlike terms]

Question. 15 Coefficient of y in the term of $-y^3$ is

(a)-1 (b)-3 (c) -1^3 (d) 1^3

Solution.

(c) We can write $\frac{-y}{3}$ as $-\frac{1}{3} \times y$.

So, the coefficient of y is $-\frac{1}{3}$.

Question. 16 $a^2 - b^2$ is equal to

(a) $(a - b)^2$

(b) $(a - b)(a - b)$

(c) $(a + b)(a - b)$

(d) $(a + b)(a + b)$

Solution.

(c) (a) $(a - b)^2 = a^2 - 2ab + b^2$

(b) $(a - b)(a - b) = (a - b)^2 = a^2 - 2ab + b^2$

(c) $(a + b)(a - b) = a(a - b) + b(a - b) = a^2 - ab + ba - b^2 = a^2 - b^2$ [$\because ab = ba$]

(d) $(a + b)(a + b) = (a + b)^2 = a^2 + 2ab + b^2$

Question. 17 Common factor Of $17abc, 34ab^2, 51a^2b$ is

(a) $17abc$ (b) $17ab$ (c) $17ac$ (d) $17a^2b^2c$

Solution.

(b) Given, $17abc = 17 \times a \times b \times c$

$$34ab^2 = 2 \times 17 \times a \times b \times b$$

$$51a^2b = 3 \times 17 \times a \times a \times b$$

Now, collecting the common factors, we get $17 \times a \times b = 17ab$

Question. 18 Square of $9x - 7xy$ is

(a) $81x^2 + 49x^2y^2$

(b) $81x^2 - 49x^2y^2$

(c) $81x^2 + 49x^2y^2 - 126x^2y$

(d) $81x^2 + 49x^2y^2 - 63x^2y$

Solution.

(c) Square of $(9x - 7xy) = (9x - 7xy)^2$

Comparing with $(a - b)^2$, we get $a = 9x$ and $b = 7xy$

$$(9x - 7xy)^2 = (9x)^2 - 2 \cdot 9x \cdot 7xy + (7xy)^2 \quad [\text{using the identity, } (a - b)^2 = a^2 - 2ab + b^2]$$

$$= 81x^2 - 126x^2y + 49x^2y^2$$

$$= 81x^2 + 49x^2y^2 - 126x^2y$$

Question. 19 Factorised form of $23xy - 46x + 54y - 108$ is

- (a) $(23x + 54)(y - 2)$ (b) $(23x + 54y)(y - 2)$
 (c) $(23xy + 54y)(-46x - 108)$ (d) $(23x + 54)(y + 2)$

Solution.

(a) We have, $23xy - 46x + 54y - 108 = 23xy - 2 \times 23x + 54y - 2 \times 54$
 $= 23x(y - 2) + 54(y - 2)$
 [taking common out in I and II expressions]
 $= (y - 2)(23x + 54)$ [taking $(y - 2)$ common]
 $= (23x + 54)(y - 2)$

Question. 20 Factorised form of $r^2 - 10r + 21$ is

- (a) $(r-1)(r-4)$ (b) $(r-7)(r-3)$ (c) $(r-7)(r+3)$ (d) $(r+7)(r+3)$

Solution.

(b) We have, $r^2 - 10r + 21$
 $= r^2 - 7r - 3r + 21 = r(r - 7) - 3(r - 7)$
 [by splitting the middle term, so that the product of their numerical coefficients is equal constant term]
 $= (r - 7)(r - 3)$ [$\because x^2 + (a + b)x + ab = (x + a)(x + b)$]

Question. 21 Factorised form of $p^2 - 17p - 38$ is

- (a) $(p - 19)(p + 2)$ (b) $(p - 19)(p - 2)$ (c) $(p + 19)(p + 2)$ (d) $(p + 19)(p - 2)$

Solution.

(a) We have, $p^2 - 17p - 38 = p^2 - 19p + 2p - 38$
 [by splitting the middle term, so that the product of their numerical coefficients is equal constant term]
 $= p(p - 19) + 2(p - 19) = (p - 19)(p + 2)$ [$\because x^2 + (a + b)x + ab = (x + a)(x + b)$]

Question. 22 On dividing $57 p^2 qr$ by $114pq$, we get

- (a) $\frac{1}{4} pr$ (b) $\frac{3}{4} pr$ (c) $\frac{1}{2} pr$ (d) $2pr$

Solution.

(c) Required value = $\frac{57 p^2 qr}{114 pq} = \frac{57 \times p \times p \times q \times r}{114 \times p \times q} = \frac{57}{114} pr = \frac{1}{2} pr$

Question. 23 On dividing $p(4p^2 - 16)$ by $4p(p - 2)$, we get

- (a) $2p + 4$ (b) $2p - 4$ (c) $p + 2$ (d) $p - 2$

Solution.

(c) We have,
 $\frac{p(4p^2 - 16)}{4p(p - 2)} = \frac{p[(2p)^2 - 4^2]}{4p(p - 2)}$
 $= \frac{(2p - 4)(2p + 4)}{4(p - 2)}$ [$\because a^2 - b^2 = (a + b)(a - b)$]
 $= \frac{2(p - 2) \cdot 2(p + 2)}{4(p - 2)} = \frac{4(p - 2)(p + 2)}{4(p - 2)} = p + 2$

Question. 24 The common factor of $3ab$ and $2cd$ is

- (a) 1 (b) -1 (c) a (d) c

Solution. (a) We have, monomials $3ab$ and $2cd$ Now, $3ab = 3 \times a \times b$ $2cd = 2 \times c \times d$
 Observing the monomials, we see that, there is no common factor (neither numerical nor literal) between them except 1.

Question. 25 An irreducible factor of $24x^2y^2$ is

- (a) a^2 (b) y^2 (c) x (d) $24x$

Solution. (c) A factor is said to be irreducible, if it cannot be factorised further.

We have, $24x^2y^2 = 2 \times 2 \times 2 \times 3 \times x \times x \times y \times y$ Hence, an irreducible factor of $24x^2y^2$ is x .

Question. 26 Number of factors of $(a + b)^2$ is

(a) 4 (b) 3 (c) 2 (d) 1

Solution. (c) We can write $(a + b)^2$ as, $(a + b)(a + b)$ and this cannot be factorised further.

Hence, number of factors of $(a + b)^2$ is 2.

Question. 27 The factorised form of $3x - 24$ is

(a) $3x \times 24$ (b) $3(x - 8)$ (c) $24(x - 3)$ (d) $3(x-12)$

Solution. (b) We have,

$3x - 24 = 3 \times x - 3 \times 8 = 3(x - 8)$ [taking 3 as common]

Question. 28 The factors of $x^2 - 4$ are

(a) $(x - 2), (x - 2)$ (b) $(x + 2), (x - 2)$

(c) $(x + 2), (x + 2)$ (d) $(x - 4), (x - 4)$

Solution.

(b) We have,

$$x^2 - 4 = x^2 - 2^2 = (x + 2)(x - 2)$$

$$[\because a^2 - b^2 = (a + b)(a - b)]$$

Hence, $(x + 2), (x - 2)$ are factors of $x^2 - 4$.

Question. 29 The value of $(-27x^2y) \div (-9xy)$ is

(a) $3xy$ (b) $-3xy$ (c) $-3x$ (d) $3x$

Solution.

(d) We have,

$$\frac{(-27x^2y) \div (-9xy)}{(-9xy)} = \frac{-27x^2y}{-9xy} = \frac{27 \times x \times x \times y}{9 \times x \times y} = \frac{27}{9}x = 3x$$

Question. 30 The value of $(2x^2 - 4) \div (2)$ is

(a) $2x^2 + 2$

(b) $x^2 + 2$

(c) $x^2 + 4$

(d) $2x^2 + 4$

Solution.

(b) We have,

$$\begin{aligned} \frac{(2x^2 - 4) \div (2)}{(2)} &= \frac{2x^2 - 4}{2} = \frac{2(x^2 - 2)}{2} \\ &= x^2 - 2 \end{aligned}$$

[taking 2 as common]

Question. 31 The value of $(3x^3 - 9x^2 + 27x) \div 3x$ is

(a) $x^2 + 9 + 27x$

(b) $3x^2 + 3x^2 + 27x$

(c) $3x^2 + 9x^2 + 9$

(d) $x^2 + 3x + 9$

Solution.

(d) We have,

$$(3x^3 - 9x^2 + 27x) \div 3x = \frac{3x^3 - 9x^2 + 27x}{3x} = \frac{3x^3}{3x} + \frac{9x^2}{3x} + \frac{27x}{3x} = x^2 + 3x + 9$$

Question. 32 The value of $(a + b)^2 + (a - b)^2$ is

(a) $2a + 2b$

(b) $2a - 2b$

(c) $2a^2 + 2b^2$

(d) $2a^2 - 2b^2$

Solution.

(c) We have,

$$\begin{aligned}(a + b)^2 + (a - b)^2 &= (a^2 + b^2 + 2ab) + (a^2 + b^2 - 2ab) \\ &= (a^2 + a^2) + (b^2 + b^2) + (2ab - 2ab) \quad [\because (a + b)^2 = a^2 + b^2 + 2ab \text{ and } (a - b)^2 = a^2 + b^2 - 2ab] \\ &= 2a^2 + 2b^2 \quad [\text{combining the like terms}]\end{aligned}$$

Question. 33 The value of $(a + b)^2 - (a - b)^2$ is

- (a) $4ab$ (b) $-4ab$ (c) $2a^2 + 2b^2$ (d) $2a^2 - 2b^2$

Solution.

(a) We have,

$$\begin{aligned}(a + b)^2 - (a - b)^2 &= a^2 + b^2 + 2ab - (a^2 + b^2 - 2ab) \\ &= a^2 + b^2 + 2ab - a^2 - b^2 + 2ab = a^2 - a^2 + b^2 - b^2 + 2ab + 2ab = 2ab + 2ab = 4ab \\ & \quad [\text{combining the like terms}]\end{aligned}$$

Fill in the Blanks

In questions 34 to 58, fill in the blanks to make the statements true.

Question. 34 The product of two terms with like signs is a term.

Solution. Positive

If both the like terms are either positive or negative, then the resultant term will always be positive.

Question. 35 The product of two terms with unlike signs is a term.

Solution. Negative

As the product of a positive term and a negative term is always negative.

Question. 36 $a(b + c) = a \times \text{---} + a \times \text{---}$

Solution. b,c

we have, $a(b+c)=a \times b + a \times c$ [using left distributive law]

Question. 37 $(a-b) \text{---} = a^2 - 2ab + b^2$

Solution.

(a - b)

$$\begin{aligned}\text{We know that, } (a - b)(a - b) &= (a - b)^2 \\ &= a^2 - 2ab + b^2 \quad [\because (a - b)^2 = a^2 - 2ab + b^2]\end{aligned}$$

Question. 38 $a^2 - b^2 = (a+b) \text{---}$

Solution.

(a - b)

$$\text{We have, } a^2 - b^2 = (a + b)(a - b) \quad [\because a^2 - b^2 = (a + b)(a - b)]$$

Alternative Method

$$\begin{aligned}\text{Let } (a^2 - b^2) &= (a + b)x \\ \Rightarrow x &= \frac{a^2 - b^2}{a + b} = \frac{(a + b)(a - b)}{a + b} = a - b\end{aligned}$$

Question. 39 $(a - b)^2 + \text{---} = a^2 - b^2$

Solution.

$$2ab - 2b^2$$

$$\text{Let } (a - b)^2 + x = a^2 - b^2$$

$$\Rightarrow a^2 + b^2 - 2ab + x = a^2 - b^2 \quad [\because (a - b)^2 = a^2 + b^2 - 2ab]$$

$$\Rightarrow x = a^2 - b^2 - (a^2 + b^2 - 2ab) = a^2 - b^2 - a^2 - b^2 + 2ab = 2ab - 2b^2$$

Question. 40 $(a + b)^2 - 2ab = \text{-----} + \text{-----}$.

Solution.

$$a^2 + b^2$$

We have,

$$(a + b)^2 - 2ab = a^2 + b^2 + 2ab - 2ab \quad [\because (a + b)^2 = a^2 + b^2 + 2ab]$$

$$= a^2 + b^2$$

Question. 41 $(x+a)(x+b) = x^2 + (a+b)x + \text{-----}$.

Solution.

$$ab$$

We have,

$$(x + a)(x + b) = x^2 + bx + ax + ab$$

$$= x^2 + (a + b)x + ab$$

Question. 42 The product of two polynomials is a -----.

Solution. Polynomial

As the product of two polynomials is again a polynomial.

Question. 43 Common factor of $ax^2 + bx$ is -----.

Solution.

$$x$$

$$\text{We have, } ax^2 + bx = x(ax + b) \quad [\text{taking } x \text{ as common}]$$

Question. 44 Factorised form of $18mn + 10mnp$ is -----.

Solution.

$$2mn(9 + 5p)$$

$$\text{We have, } 18mn + 10mnp = 2 \times 9 \times m \times n + 2 \times 5 \times m \times n \times p$$

$$= 2mn(9 + 5p) \quad [\text{taking } 2mn \text{ as common}]$$

Question. 45 Factorised form of $4y^2 - 12y + 9$ is -----.

Solution.

$$(2y - 3)(2y - 3)$$

Let

$$4y^2 - 12y + 9 = (2y)^2 - 2 \times 2y \times 3 + 3^2$$

$$= (2y - 3)^2 \quad [\because (a - b)^2 = a^2 - 2ab + b^2]$$

$$= (2y - 3)(2y - 3)$$

Question. 46 $38x^3y^2z \div 19xy^2$ is equal to -----.

Solution.

$$2x^2z$$

$$\text{We have, } 38x^3y^2z + 19xy^2$$

$$\text{i.e., } \frac{38x^3y^2z}{19xy^2} = \frac{38 \times x \times x \times x \times y \times y \times z}{19 \times x \times y \times y} = \frac{38}{19} x^2z = 2x^2z$$

Question. 47 Volume of a rectangular box with length $2x$, breadth $3y$ and height $4z$ is ---.

Solution. 24 xyz

We know that, the volume of a rectangular box,

$$V = \text{Length} \times \text{Breadth} \times \text{Height} = 2x \times 3y \times 4z = (2 \times 3 \times 4) xyz = 24 xyz$$

Question. 48 $67^2 - 37^2 = (67 - 37) \times \text{-----} = \text{-----}$.

Solution.

67 + 37, 3120

$$\begin{aligned} \text{We have, } 67^2 - 37^2 &= (67 - 37) (67 + 37) \\ &= 30 \times 104 = 3120 \end{aligned}$$

$$[\because a^2 - b^2 = (a - b) (a + b)]$$

Question. 49 $103^2 - 102^2 = \text{-----} \times (103 - 102) = \text{-----}$.

Solution.

(103 + 102), 205

We have,

$$\begin{aligned} 103^2 - 102^2 &= (103 + 102)(103 - 102) \\ &= 205 \times 1 = 205 \end{aligned}$$

$$[\because a^2 - b^2 = (a + b) (a - b)]$$

Question. 50 Area of a rectangular plot with sides $4y^2$ and $3y^2$ is-----.

Solution.

$12x^2y^2$

We know that, area of rectangle = Length \times Breadth

$$\therefore \text{Area of a rectangular plot} = 4x^2 \times 3y^2 = (4 \times 3)x^2y^2 = 12x^2y^2$$

Question. 51 Volume of a rectangular box with $l = b = h = 2x$ is -----.

Solution.

$8x^3$

$$\begin{aligned} \text{Volume of a rectangular box} &= l \times b \times h = 2x \times 2x \times 2x \\ &= (2 \times 2 \times 2) x^3 \\ &= 8x^3 \end{aligned}$$

Question. 52 The numerical coefficient in $-37abc$ is-----.

Solution. -37

The constant term (with their sign) involved in term of an algebraic expression is called the numerical coefficient of that term.

Question. 53 Number of terms in the expression a^2 and $+bc \times d$ is -.

Solution.

$$\text{We have, } a^2 + bc \times d = a^2 + bcd$$

\therefore The number of terms in this expression is 2 as bcd is treated as a single term.

Question. 54 The sum of areas of two squares with sides $4a$ and $4b$ is-----.

Solution.

$16(a^2 + b^2)$

$$\therefore \text{Area of a square} = (\text{Side})^2$$

$$\therefore \text{Area of the square whose one side is } 4a = (4a)^2 = 16a^2$$

$$\text{Area of the square with side } 4b = (4b)^2 = 16b^2$$

$$\therefore \text{Sum of areas} = 16a^2 + 16b^2 = 16(a^2 + b^2)$$

Question. 55 The common factor method of factorisation for a polynomial is based on

-----property.

Solution.Distributive

In this method, we regroup the terms in such a way, so that each term in the group contains a common literal or number or both.

Question. 56 The side of the square of area $9y^2$ is-----.

Solution.

3y

Given, area of a square = $9y^2$

We know that, the area of a square with side $a = a^2$

$$\therefore a^2 = 9y^2$$

$$\Rightarrow a^2 = (3y)^2$$

$$\Rightarrow a = 3y \quad \text{[taking square root both sides]}$$

Question. 57 On simplification, $\frac{3x+3}{3}$ =-----.

Solution.

x + 1

$$\text{We have, } \frac{3x+3}{3} = \frac{3x}{3} + \frac{3}{3} = x + 1$$

Question. 58 The factorisation of $2x + 4y$ is-----.

Solution. $2(x + 2y)$

We have, $2x + 4y = 2x + 2 \times 2y = 2(x + 2y)$

True/False

In questions 59 to 80, state whether the statements are True or False

Question. 59 $(a + b)^2 = a^2 + b^2$.

Solution.

False

$$\text{We have, } (a + b)^2 = a^2 + b^2 + 2ab \quad \text{[an algebraic identity]}$$

Question. 60 $(a - b)^2 = a^2 - b^2$.

Solution.

False

$$\text{We have, } (a - b)^2 = a^2 + b^2 - 2ab \quad \text{[an algebraic identity]}$$

Question. 61 $(a+b)(a-b)=a^2 - b^2$

Solution.

True

$$\begin{aligned} \text{We know that, } (a + b)(a - b) &= a \times a - a \times b + b \times a - b \times b \\ &= a^2 - b^2 \\ &= a^2 - ab + ba - b^2 \end{aligned}$$

Question. 62 The product of two negative terms is a negative term.

Solution.False

Since, the product of two negative terms is always a positive term, i.e. $(-) \times (-) = (+)$.

Question. 63 The product of one negative and one positive term is a negative term.

Solution.True

When we multiply a negative term by a positive term, the resultant will be a negative term, i.e.

$(-) \times (+) = (-)$.

Question. 64 The numerical coefficient of the term $-6x^2y^2$ is -6.

Solution. True

Since, the constant term (i.e. a number) present in the expression $-6x^2y^2$ is -6.

Question. 65 $p^2q+q^2r+r^2q$ is a binomial.

Solution. False

Since, the given expression contains three unlike terms, so it is a trinomial.

Question. 66 The factors of $a^2 - 2ab + b^2$ are $(a + b)$ and $(a + b)$.

Solution.

False

We have, $a^2 - 2ab + b^2 = (a - b)^2 = (a - b)(a - b)$ [an algebraic identity]

Question. 67 h is a factor of $2\pi(h + r)$.

Solution.

False

h is not a factor of $2\pi(h + r)$.

This expression has only two factors 2π and $(h + r)$.

Question. 68 Some of the factors of $\frac{n^2}{2} + \frac{n}{2}$ are $\frac{1}{2}n$ and $(n+1)$.

Solution.

True

We have, $\frac{n^2}{2} + \frac{n}{2} = \frac{n^2 + n}{2} = \frac{1}{2}n(n + 1)$

\therefore The factors are $\frac{1}{2}n$ and $(n + 1)$.

Question. 69 An equation is true for all values of its variables.

Solution. False

As equation is true only for some values of its variables, e.g. $2x - 4 = 0$ is true, only for $x = 2$.

Question. 70 $x^2 + (a+b)x + ab = (a+b)(x + ab)$

Solution.

False

As we know that,

$$x^2 + (a + b)x + ab = (x + a)(x + b)$$

Question. 71 Common factors of $11pq^3$, $121p^2q^3$, $1331p^3q$ is $11p^2q^2$

Solution.

False

We have,

$$11pq^3 = 11 \times p \times q \times q$$

$$121p^2q^3 = 11 \times 11 \times p \times p \times q \times q \times q$$

$$1331p^3q = 11 \times 11 \times 11 \times p \times p \times p \times q$$

\therefore Common factor = $11 \times p \times q = 11pq$

Question. 72 Common factors of $121a^2b^2 + 4ab^2 - 32$ is 4.

Solution.

True

As we have,

$$12a^2b^2 + 4ab^2 - 32 = 2 \times 2 \times 3 \times a \times a \times b \times b + 2 \times 2 \times a \times b \times b - 2^2 \times 2^3 \\ = 4(3a^2b^2 + ab^2 - 8)$$

Thus, the common factor is 4.

Question. 73 Factorisation of $-3a^2+3ab+3ac$ is $3a(-a-b-c)$.

Solution.

False

We have,

$$-3a^2 + 3ab + 3ac = 3a(-a + b + c)$$

Question. 74 Factorised form of $p^2+30p+216$ is $(p+18)(p-12)$.

Solution.

False

We have,

$$p^2 + 30p + 216 = p^2 + (12 + 18)p + 216 \\ = p^2 + 12p + 18p + 216 \quad \text{[by splitting the middle term]} \\ = p(p + 12) + 18(p + 12) \\ = (p + 18)(p + 12)$$

Question. 75 The difference of the squares of two consecutive numbers is their sum.

Solution.

True

Let n and $n + 1$ be any two consecutive numbers, then their sum $= n + n + 1 = 2n + 1$

Now, the difference of their squares,

$$(n + 1)^2 - n^2 = n^2 + 1 + 2n - n^2 \\ = 2n + 1 \quad \text{[}\therefore (a + b)^2 = a^2 + 2ab + b^2\text{]}$$

Question. 76 $abc + bca + cab$ is a monomial.

Solution. True

The given expression seems to be a trinomial but it is not as it contains three like terms which can be added to form a monomial, i.e. $abc + abc + abc = 3abc$

Question. 77 On dividing $\frac{p}{3}$ by $\frac{3}{p}$, the quotient is 9

Solution.

False

$$\text{We have, } \frac{p}{3} \div \frac{3}{p} = \frac{p}{3} \times \frac{p}{3} = \frac{1}{9}p^2$$

$$\left[\therefore \text{reciprocal of } \frac{3}{p} \text{ is } \frac{p}{3} \right]$$

Hence, the quotient is $\frac{1}{9}p^2$.

Question. 78 The value of p for $51^2-49^2=100p$ is 2.

Solution.

True

$$\text{We have, } 51^2 - 49^2 = 100p$$

$$\Rightarrow (51 + 49)(51 - 49) = 100p$$

$$\Rightarrow 100 \times 2 = 100p$$

$$\Rightarrow p = 2$$

$$\left[\therefore a^2 - b^2 = (a + b)(a - b) \right]$$

Question. 79 $(9x - 51) \div 9$ is $x - 51$.

Solution.

False

$$\text{We have, } (9x - 51) \div 9 = \frac{9x - 51}{9} = \frac{9x}{9} - \frac{51}{9} = x - \frac{51}{9}$$

Question. 80 The value of $(a+1)(a-1)(a^2+1)$ is a^4-1 .

Solution.

True

$$\text{We have, } (a + 1)(a - 1)(a^2 + 1) = (a^2 - 1)(a^2 + 1)$$

[using the identity, $(a + b)(a - b) = a^2 - b^2$ in first two factors]

$$= (a^2)^2 - 1^2 \quad \text{[again using the same identity]}$$

$$= a^4 - 1$$

Question. 81 Add:

(i) $7a^2bc, -3abc^2, 3a^2bc, 2abc^2$

(ii) $9ax + 3by - cz, -5by + ax + 3cz$

(iii) $xy^2z^2 + 3x^2y^2z - 4x^2yz^2, -9x^2y^2z + 3xy^2z^2 + x^2yz^2$

(iv) $5x^2 - 3xy + 4y^2 - 9, 7y^2 + 5xy - 2x^2 + 13$

(v) $2p^4 - 3p^3 + p^2 - 5p + 7, -3p^4 - 7p^3 - 3p^2 - p - 12$

(vi) $3a(a - b + c), 2b(a - b + c)$

(vii) $3a(2b + 5c), 3c(2a + 2b)$

Solution.

(i) We have,

$$\begin{aligned} 7a^2bc + (-3abc^2) + 3a^2bc + 2abc^2 &= 7a^2bc - 3abc^2 + 3a^2bc + 2abc^2 \\ &= (7a^2bc + 3a^2bc) + (-3abc^2 + 2abc^2) \quad \text{[grouping like terms]} \\ &= 10a^2bc + (-abc^2) \\ &= 10a^2bc - abc^2 \end{aligned}$$

(ii) We have,

$$\begin{aligned} (9ax + 3by - cz) + (-5by + ax + 3cz) &= 9ax + 3by - cz - 5by + ax + 3cz \\ &= (9ax + ax) + (3by - 5by) + (-cz + 3cz) \quad \text{[grouping like terms]} \\ &= 10ax - 2by + 2cz \end{aligned}$$

(iii) We have,

$$\begin{aligned}(xy^2z^2 + 3x^2y^2z - 4x^2yz^2) + (-9x^2y^2z + 3xy^2z^2 + x^2yz^2) \\= xy^2z^2 + 3x^2y^2z - 4x^2yz^2 - 9x^2y^2z + 3xy^2z^2 + x^2yz^2 \\= (xy^2z^2 + 3xy^2z^2) + (3x^2y^2z - 9x^2y^2z) + (-4x^2yz^2 + x^2yz^2) \\= 4xy^2z^2 - 6x^2y^2z - 3x^2yz^2\end{aligned}$$

[grouping like terms]

(iv) We have,

$$\begin{aligned}(5x^2 - 3xy + 4y^2 - 9) + (7y^2 + 5xy - 2x^2 + 13) \\= 5x^2 - 3xy + 4y^2 - 9 + 7y^2 + 5xy - 2x^2 + 13 \\= (5x^2 - 2x^2) + (-3xy + 5xy) + (4y^2 + 7y^2) + (-9 + 13) \\= 3x^2 + 2xy + 11y^2 + 4\end{aligned}$$

[grouping like terms]

(v) We have,

$$\begin{aligned}(2p^4 - 3p^3 + p^2 - 5p + 7) + (-3p^4 - 7p^3 - 3p^2 - p - 12) \\= 2p^4 - 3p^3 + p^2 - 5p + 7 - 3p^4 - 7p^3 - 3p^2 - p - 12 \\= (2p^4 - 3p^4) + (-3p^3 - 7p^3) + (p^2 - 3p^2) + (-5p - p) + (7 - 12) \\= -p^4 - 10p^3 - 2p^2 - 6p - 5\end{aligned}$$

[grouping like terms]

(vi) We have,

$$\begin{aligned}3a(a - b + c) + 2b(a - b + c) \\= (3a^2 - 3ab + 3ac) + (2ab - 2b^2 + 2bc) \\= 3a^2 - 3ab + 2ab + 3ac + 2bc - 2b^2 = 3a^2 - ab + 3ac + 2bc - 2b^2\end{aligned}$$

[grouping like terms]

(vii) We have,

$$\begin{aligned}3a(2b + 5c) + 3c(2a + 2b) \\= (6ab + 15ac) + (6ac + 6bc) = 6ab + 15ac + 6ac + 6bc \\= 6ab + 21ac + 6bc\end{aligned}$$

[grouping like terms]

Question. 82 Subtract

- (i) $5a^2b^2c^2$ from $-7a^2b^2c^2$
- (ii) $6x^2 - 4xy + 5y^2$ from $8y^2 + 6xy - 3x^2$
- (iii) $2ab^2c^2 + 4a^2b^2c - 5a^2bc^2$ from $-10a^2b^2c + 4ab^2c^2 + 2a^2bc^2$
- (iv) $3t^4 - 4t^3 + 2t^2 - 6t + 6$ from $-4t^4 + 8t^3 - 4t^2 - 2t + 11$
- (v) $2ab + 5bc - 7ac$ from $5ab - 2bc - 2ac + 10abc$
- (vi) $7p(3q + 7p)$ from $8p(2p - 7q)$
- (vii) $-3p^2 + 3pq + 3px$ from $3p(-p - a - r)$

Solution.

(i) We have, $5a^2b^2c^2$ and $-7a^2b^2c^2$

$$\begin{aligned}\text{The required difference is given by } -7a^2b^2c^2 - 5a^2b^2c^2 \\= (-7 - 5)a^2b^2c^2 = -12a^2b^2c^2\end{aligned}$$

(ii) We have, $6x^2 - 4xy + 5y^2$ and $8y^2 + 6xy - 3x^2$

$$\begin{aligned} \text{The required difference is given by } & (8y^2 + 6xy - 3x^2) - (6x^2 - 4xy + 5y^2) \\ & = 8y^2 + 6xy - 3x^2 - 6x^2 + 4xy - 5y^2 \\ & = (8y^2 - 5y^2) + (6xy + 4xy) - (3x^2 + 6x^2) = 3y^2 + 10xy - 9x^2 \end{aligned}$$

(iii) We have,

$$2ab^2c^2 + 4a^2b^2c - 5a^2bc^2 \text{ and } -10a^2b^2c + 4ab^2c^2 + 2a^2bc^2$$

The required difference is given by

$$\begin{aligned} & (-10a^2b^2c + 4ab^2c^2 + 2a^2bc^2) - (2ab^2c^2 + 4a^2b^2c - 5a^2bc^2) \\ & = -10a^2b^2c + 4ab^2c^2 + 2a^2bc^2 - 2ab^2c^2 - 4a^2b^2c + 5a^2bc^2 \\ & = (-10a^2b^2c - 4a^2b^2c) + (4ab^2c^2 - 2ab^2c^2) + (2a^2bc^2 + 5a^2bc^2) \\ & \hspace{15em} \text{[grouping like terms]} \\ & = -14a^2b^2c + 2ab^2c^2 + 7a^2bc^2 \end{aligned}$$

(iv) We have, $3t^4 - 4t^3 + 2t^2 - 6t + 6$ and $-4t^4 + 8t^3 - 4t^2 - 2t + 11$

The required difference is given by

$$\begin{aligned} & (-4t^4 + 8t^3 - 4t^2 - 2t + 11) - (3t^4 - 4t^3 + 2t^2 - 6t + 6) \\ & = -4t^4 + 8t^3 - 4t^2 - 2t + 11 - 3t^4 + 4t^3 - 2t^2 + 6t - 6 \\ & = (-4t^4 - 3t^4) + (8t^3 + 4t^3) + (-4t^2 - 2t^2) + (-2t + 6t) + (11 - 6) \\ & \hspace{15em} \text{[grouping like terms]} \\ & = -7t^4 + 12t^3 - 6t^2 + 4t + 5 \end{aligned}$$

(v) We have, $2ab + 5bc - 7ac$ and $5ab - 2bc - 2ac + 10abc$

$$\begin{aligned} \text{The required difference is given by } & (5ab - 2bc - 2ac + 10abc) - (2ab + 5bc - 7ac) \\ & = 5ab - 2bc - 2ac + 10abc - 2ab - 5bc + 7ac \\ & = (5ab - 2ab) + (-2bc - 5bc) + (-2ac + 7ac) + 10abc \\ & \hspace{15em} \text{[grouping like terms]} \\ & = 3ab - 7bc + 5ac + 10abc \end{aligned}$$

(vi) We have, $7p(3q - 7p)$ and $8p(2p - 7q)$

$$\begin{aligned} \text{The required difference is given by } & 8p(2p - 7q) - 7p(3q + 7p) \\ & = 16p^2 - 56pq - 21pq - 49p^2 = (16p^2 - 49p^2) + (-56pq - 21pq) \\ & \hspace{15em} \text{[grouping like terms]} \\ & = -33p^2 - 77pq \end{aligned}$$

(vii) We have, $-3p^2 + 3pq + 3px$ and $3p(-p - a - r)$

The required difference is given by

$$\begin{aligned} & 3p(-p - a - r) - (-3p^2 + 3pq + 3px) = -3p^2 - 3ap - 3pr + 3p^2 - 3pq - 3px \\ & = (-3p^2 + 3p^2) - 3ap - 3pr - 3pq - 3px = -3ap - 3pr - 3pq - 3px \\ & \hspace{15em} \text{[grouping like terms]} \end{aligned}$$

Question. 83 Multiply the following:

- | | |
|--------------------------------|-------------------------------------|
| (i) $-7pq^2r^3$, $-13p^3q^2r$ | (ii) $3x^2y^2z^2$, $17xyz$ |
| (iii) $15xy^2$, $17yz^2$ | (iv) $-5a^2bc$, $11ab$, $13abc^2$ |
| (v) $-3x^2y$, $(5y - xy)$ | (vi) abc , $(bc + ca)$ |

- (vii) $7pqr, (p - q + r)$ (viii) $x^2y^2z^2, (xy - yz + zx)$
 (xi) $(p + 6), (q - 7)$ (x) $6mn, 0mn$
 (xii) a, a^5, a^6 (xiii) $-7st, -1, -13st^2$
 (xiv) $-\frac{100}{9}rs; \frac{3}{4}r^3s^2$
 (xv) $(a^2 - b^2), (a^2 + b^2)$ (xvi) $(ab + c), (ab + c)$
 (xvii) $(pq - 2r), (pq - 2r)$ (xviii) $\left(\frac{3}{4}x - \frac{4}{3}y\right), \left(\frac{2}{3}x + \frac{3}{2}y\right)$
 (xix) $\frac{3}{2}p^2 + \frac{2}{3}q^2, (2p^2 - 3q^2)$ (xx) $(x^2 - 5x + 6), (2x + 7)$
 (xxi) $(3x^2 + 4x - 8), (2x^2 - 4x + 3)$
 (xxii) $(2x - 2y - 3), (x + y + 5)$

Solution.

(i) We have, $-7pq^2r^3$ and $-13p^3q^2r$

$$\therefore (-7pq^2r^3) \times (-13p^3q^2r) = (-7) \times (-13) p^4q^4r^4 = 91p^4q^4r^4$$

(ii) We have, $3x^2y^2z^2$ and $17xyz$

$$\therefore 3x^2y^2z^2 \times 17xyz = (3 \times 17)x^3y^3z^3 = 51x^3y^3z^3$$

(iii) We have, $15xy^2$ and $17yz^2$

$$15xy^2 \times 17yz^2 = (15 \times 17)xy^2 \times yz^2 = 255xy^3z^2$$

(iv) We have, $-5a^2bc$, $11ab$ and $13abc^2$

$$\therefore -5a^2bc \times 11ab \times 13abc^2 = (-5 \times 11 \times 13)a^2bc \times ab \times abc^2 = -715a^4b^3c^3$$

(v) We have, $-3x^2y$ and $(5y - xy)$

$$\therefore -3x^2y \times (5y - xy) = -3x^2y \times 5y + 3x^2y \times xy = -15x^2y^2 + 3x^3y^2$$

(vi) We have, abc and $(bc + ca)$

$$\therefore abc \times (bc + ca) = abc \times bc + abc \times ca = ab^2c^2 + a^2bc^2$$

(vii) We have, $7pqr$ and $(p - q + r)$

$$\therefore 7pqr \times (p - q + r) = 7pqr \times p - 7pqr \times q + 7pqr \times r = 7p^2qr - 7pq^2r + 7pqr^2$$

(viii) We have, $x^2y^2z^2$ and $(xy - yz - zx)$

$$\begin{aligned} \therefore x^2y^2z^2 \times (xy - yz + zx) &= x^2y^2z^2 \times xy - x^2y^2z^2 \times yz + x^2y^2z^2 \times zx \\ &= x^3y^3z^2 - x^2y^3z^3 + x^3y^2z^3 \end{aligned}$$

(ix) We have, $(p + 6)$ and $(q - 7)$

$$\therefore (p + 6) \times (q - 7) = p(q - 7) + 6(q - 7) = pq - 7p + 6q - 42$$

(x) We have, $6mn$ and $0mn$

$$\therefore 6mn \times 0mn = (6 \times 0)mn \times mn = 0 \times m^2n^2 = 0$$

(xi) We have, a , a^5 and a^6

$$\therefore a \times a^5 \times a^6 = a^{1+5+6} = a^{12}$$

(xii) We have, $-7st$, -1 and $-13st^2$

$$\therefore -7st \times (-1) \times (-13st^2) = [-7 \times (-1) \times (-13)]st \times (st^2) = -91s^2t^3$$

(xiii) We have, b^3 , $3b^2$ and $7ab^5$

$$\therefore b^3 \times 3b^2 \times 7ab^5 = (1 \times 3 \times 7)b^3 \times b^2 \times ab^5 = 21ab^{10}$$

(xiv) We have, $\frac{-100}{9}rs$ and $\frac{3}{4}r^3s^2$

$$\therefore \frac{-100}{9}rs \times \frac{3}{4}r^3s^2 = \left(\frac{-100}{9} \times \frac{3}{4}\right)rs \times r^3s^2 = \frac{-25}{3} \times r^4s^3$$

(xv) We have, $(a^2 - b^2)$ and $(a^2 + b^2)$

$$\therefore (a^2 - b^2)(a^2 + b^2) = a^2(a^2 + b^2) - b^2(a^2 + b^2) = a^4 + a^2b^2 - b^2a^2 - b^4 = a^4 - b^4$$

(xvi) We have, $(ab + c)$ and $(ab + c)$

$$\begin{aligned}\therefore (ab + c)(ab + c) &= ab(ab + c) + c(ab + c) \\ &= a^2b^2 + abc + cab + c^2 = a^2b^2 + 2abc + c^2\end{aligned}$$

(xvii) We have, $(pq - 2r)$ and $(pq - 2r)$

$$\begin{aligned}\therefore (pq - 2r)(pq - 2r) &= pq(pq - 2r) - 2r(pq - 2r) \\ &= p^2q^2 - 2pqr - 2rpq + 4r^2 = p^2q^2 - 4pqr + 4r^2\end{aligned}$$

(xviii) We have, $\left(\frac{3}{4}x - \frac{4}{3}y\right)$ and $\left(\frac{2}{3}x + \frac{3}{2}y\right)$

$$\begin{aligned}\therefore \left(\frac{3}{4}x - \frac{4}{3}y\right)\left(\frac{2}{3}x + \frac{3}{2}y\right) &= \frac{3}{4}x\left(\frac{2}{3}x + \frac{3}{2}y\right) - \frac{4}{3}y\left(\frac{2}{3}x + \frac{3}{2}y\right) \\ &= \frac{3}{4} \times \frac{2}{3}x^2 + \frac{3}{4} \times \frac{3}{2}xy - \frac{4}{3} \times \frac{2}{3}yx - \frac{4}{3} \times \frac{3}{2}y^2 \\ &= \frac{1}{2}x^2 + \frac{9}{8}xy - \frac{8}{9}xy - 2y^2 \\ &= \frac{1}{2}x^2 + \left(\frac{9}{8} - \frac{8}{9}\right)xy - 2y^2 \\ &= \frac{1}{2}x^2 + \left(\frac{81 - 64}{72}\right)xy - 2y^2 \\ &= \frac{1}{2}x^2 + \frac{17}{72}xy - 2y^2\end{aligned}$$

(xix) We have, $\left(\frac{3}{2}p^2 + \frac{2}{3}q^2\right)$ and $(2p^2 - 3q^2)$

$$\begin{aligned}\therefore \left(\frac{3}{2}p^2 + \frac{2}{3}q^2\right)(2p^2 - 3q^2) &= \frac{3}{2}p^2(2p^2 - 3q^2) + \frac{2}{3}q^2(2p^2 - 3q^2) \\ &= \frac{3}{2}p^2 \times 2p^2 - \frac{9}{2}p^2q^2 + \frac{4}{3}q^2p^2 - 2q^4 \\ &= 3p^4 + \left(\frac{4}{3} - \frac{9}{2}\right)p^2q^2 - 2q^4 \\ &= 3p^4 + \left(\frac{8 - 27}{6}\right)p^2q^2 - 2q^4 \\ &= 3p^4 - \frac{19}{6}p^2q^2 - 2q^4\end{aligned}$$

(xx) We have, $(x^2 - 5x + 6)$ and $(2x + 7)$

$$\begin{aligned}\therefore (x^2 - 5x + 6)(2x + 7) &= x^2(2x + 7) - 5x(2x + 7) + 6(2x + 7) \\ &= 2x^3 + 7x^2 - 10x^2 - 35x + 12x + 42 \\ &= 2x^3 - 3x^2 - 23x + 42\end{aligned}$$

(xix) We have, $(3x^2 + 4x - 8)$ and $(2x^2 - 4x + 3)$

$$\begin{aligned}\therefore (3x^2 + 4x - 8)(2x^2 - 4x + 3) &= 3x^2(2x^2 - 4x + 3) + 4x(2x^2 - 4x + 3) - 8(2x^2 - 4x + 3) \\ &= 6x^4 - 12x^3 + 9x^2 + 8x^3 - 16x^2 + 12x - 16x^2 + 32x - 24 \\ &= 6x^4 - 12x^3 + 8x^3 + 9x^2 - 16x^2 - 16x^2 + 12x + 32x - 24 \\ & \hspace{15em} \text{[grouping like terms]} \\ &= 6x^4 - 4x^3 - 23x^2 + 44x - 24\end{aligned}$$

(xxii) We have, $(2x - 2y - 3)$ and $(x + y + 5)$

$$\begin{aligned}\therefore (2x - 2y - 3)(x + y + 5) &= 2x(x + y + 5) - 2y(x + y + 5) - 3(x + y + 5) \\ &= 2x^2 + 2xy + 10x - 2yx - 2y^2 - 10y - 3x - 3y - 15 \\ &= 2x^2 + 2xy - 2yx + 10x - 3x - 2y^2 - 10y - 3y - 15 \\ & \hspace{15em} \text{[grouping like terms]} \\ &= 2x^2 + 7x - 13y - 2y^2 - 15\end{aligned}$$

Question. 84 Simplify

(i) $(3x + 2y)^2 + (3x - 2y)^2$

(ii) $(3x + 2y)^2 - (3x - 2y)^2$

(iii) $\left(\frac{7}{9}a + \frac{9}{7}b\right)^2 - ab$

(iv) $\left(\frac{3}{4}x - \frac{4}{3}y\right)^2 + 2xy$

(v) $(1.5p + 1.2q)^2 - (1.5p - 1.2q)^2$

(vi) $(2.5m + 1.5q)^2 + (2.5m - 1.5q)^2$

(vii) $(x^2 - 4) + (x^2 + 4) + 16$

(viii) $(ab - c)^2 + 2abc$

(ix) $(a - b)(a^2 + b^2 + ab) - (a + b)(a^2 + b^2 - ab)$

(x) $(b^2 - 49)(b + 7) + 343$

(xi) $(4.5a + 1.5b)^2 + (4.5b + 1.5a)^2$

(xii) $(pq - qr)^2 + 4pq^2r$

(xiii) $(s^2t + tq^2)^2 - (2stq)^2$

Solution.

(i) We have,

$$\begin{aligned}(3x + 2y)^2 + (3x - 2y)^2 &= (3x)^2 + (2y)^2 + 2 \times 3x \times 2y + (3x)^2 + (2y)^2 - 2 \times 3x \times 2y \\ & \hspace{15em} \text{[using the identities, } (a + b)^2 = a^2 + b^2 + 2ab \\ & \hspace{15em} \text{and } (a - b)^2 = a^2 + b^2 - 2ab]\end{aligned}$$

$$\begin{aligned}
 &= 9x^2 + 4y^2 + 12xy + 9x^2 + 4y^2 - 12xy \\
 &= (9x^2 + 9x^2) + (4y^2 + 4y^2) + 12xy - 12xy \\
 &= 18x^2 + 8y^2
 \end{aligned}$$

(ii) We have,

$$\begin{aligned}
 &(3x + 2y)^2 - (3x - 2y)^2 \\
 &= [(3x + 2y) + (3x - 2y)][(3x + 2y) - (3x - 2y)] \\
 &\quad \text{[using the identity, } a^2 - b^2 = (a + b)(a - b)\text{]} \\
 &= (3x + 2y + 3x - 2y)(3x + 2y - 3x + 2y) = 6x \times 4y = (6 \times 4) \times xy = 24xy
 \end{aligned}$$

(iii) We have, $\left(\frac{7}{9}a + \frac{9}{7}b\right)^2 - ab$

$$\begin{aligned}
 &= \left(\frac{7}{9}a\right)^2 + \left(\frac{9}{7}b\right)^2 + 2 \times \frac{7}{9}a \times \frac{9}{7}b - ab \\
 &\quad \text{[using the identity, } (a + b)^2 = a^2 + b^2 + 2ab\text{]} \\
 &= \frac{49}{81}a^2 + \frac{81}{49}b^2 + 2ab - ab \\
 &= \frac{49}{81}a^2 + ab + \frac{81}{49}b^2
 \end{aligned}$$

(iv) We have,

$$\begin{aligned}
 &\left(\frac{3}{4}x - \frac{4}{3}y\right)^2 + 2xy \\
 &= \left(\frac{3}{4}x\right)^2 + \left(\frac{4}{3}y\right)^2 - 2 \times \frac{3}{4}x \times \frac{4}{3}y + 2xy \\
 &\quad \text{[using the identity, } (a - b)^2 = a^2 + b^2 - 2ab\text{]} \\
 &= \frac{9}{16}x^2 + \frac{16}{9}y^2 - 2xy + 2xy = \frac{9}{16}x^2 + \frac{16}{9}y^2
 \end{aligned}$$

(v) We have,

$$\begin{aligned}
 &(1.5p + 1.2q)^2 - (1.5p - 1.2q)^2 \\
 &= [(1.5p + 1.2q) + (1.5p - 1.2q)][(1.5p + 1.2q) - (1.5p - 1.2q)] \\
 &\quad \text{[using the identity, } a^2 - b^2 = (a + b)(a - b)\text{]} \\
 &= [(1.5p + 1.5p) + (1.2q - 1.2q)][(1.5p - 1.5p) + (1.2q + 1.2q)] \\
 &= 3p \times 2.4q = 7.2pq
 \end{aligned}$$

(vi) We have,

$$\begin{aligned}
 &(2.5m + 1.5q)^2 + (2.5m - 1.5q)^2 \\
 &= (2.5m)^2 + (1.5q)^2 + 2 \times 2.5m \times 1.5q + (2.5m)^2 + (1.5q)^2 - 2 \times (2.5m) \times (1.5q) \\
 &\quad \text{[using the identities, } (a + b)^2 = a^2 + b^2 + 2ab \text{ and } (a - b)^2 = a^2 + b^2 - 2ab\text{]} \\
 &= 6.25m^2 + 2.25q^2 + 6.25m^2 + 2.25q^2 \\
 &= (6.25 + 6.25)m^2 + (2.25 + 2.25)q^2 \\
 &= 12.5m^2 + 4.5q^2
 \end{aligned}$$

(vii) We have,

$$\begin{aligned}
 &(x^2 - 4) + (x^2 + 4) + 16 \\
 &= x^2 - 4 + x^2 + 4 + 16 = 2x^2 + 16
 \end{aligned}$$

(viii) $(ab - c)^2 + 2abc$
 $= (ab)^2 + c^2 - 2abc + 2abc$ [using the identity, $(a - b)^2 = a^2 + b^2 - 2ab$]
 $= a^2b^2 + c^2$

(ix) $(a - b)(a^2 + b^2 + ab) - (a + b)(a^2 + b^2 - ab)$
 $= a(a^2 + b^2 + ab) - b(a^2 + b^2 + ab) - a(a^2 + b^2 - ab) - b(a^2 + b^2 - ab)$
 $= a^3 + ab^2 + a^2b - ba^2 - b^3 - ab^2 - a^3 - ab^2 + a^2b - ba^2 - b^3 + ab^2$
 $= (a^3 - a^3) + (-b^3 - b^3) + (ab^2 - ab^2) + (a^2b - a^2b + a^2b - a^2b)$
 $= 0 - 2b^3 + 0 + 0 + 0$
 $= -2b^3$

(x) We have,
 $(b^2 - 49)(b + 7) + 343$
 $= b^2(b + 7) - 49(b + 7) + 343$
 $= b^3 + 7b^2 - 49b - 343 + 343$
 $= b^3 - 49b + 7b^2$

(xi) We have, $(4.5a + 1.5b)^2 + (4.5b + 1.5a)^2$
 $= (4.5a)^2 + (1.5b)^2 + 2 \times 4.5a \times 1.5b + (4.5b)^2 + (1.5a)^2 + 2 \times 4.5b \times 1.5a$
[using the identity, $(a + b)^2 = a^2 + b^2 + 2ab$]
 $= 20.25a^2 + 2.25b^2 + 13.5ab + 20.25b^2 + 2.25a^2 + 13.5ab$
 $= 40.5a^2 + 4.5b^2 + 27ab$

(xii) We have, $(pq - qr)^2 + 4pq^2r$
 $= p^2q^2 + q^2r^2 - 2pqr^2 + 4pq^2r$
[using the identity, $(a - b)^2 = a^2 + b^2 - 2ab$]
 $= p^2q^2 + q^2r^2 + 2pq^2r$

(xiii) We have, $(s^2t + tq^2)^2 - (2stq)^2$
 $= (s^2t)^2 + (tq^2)^2 + 2 \times s^2t \times tq^2 - 4s^2t^2q^2$
[using the identity, $(a + b)^2 = a^2 + b^2 + 2ab$]
 $= s^4t^2 + t^2q^4 + 2s^2t^2q^2 - 4s^2t^2q^2$
 $= s^4t^2 + t^2q^4 - 2s^2t^2q^2$

Question. 85 Expand the following, using suitable identities.

(i) $(xy + yz)^2$ (ii) $(x^2y - xy^2)^2$
(iii) $\left(\frac{4}{5}a + \frac{5}{4}b\right)^2$ (iv) $\left(\frac{2}{3}x - \frac{3}{2}y\right)^2$
(v) $\left(\frac{4}{5}p + \frac{5}{3}q\right)^2$ (vi) $(x + 3)(x + 7)$
(vii) $(2x + 9)(2x - 7)$ (viii) $\left(\frac{4x}{5} + \frac{y}{4}\right)\left(\frac{4x}{5} + \frac{3y}{4}\right)$
(ix) $\left(\frac{2x}{3} - \frac{2}{3}\right)\left(\frac{2x}{3} + \frac{2a}{3}\right)$ (x) $(2x - 5y)(2x - 5y)$
(xi) $\left(\frac{2a}{3} + \frac{b}{3}\right)\left(\frac{2a}{3} - \frac{b}{3}\right)$ (xii) $(x^2 + y^2)(x^2 - y^2)$
(xiii) $(a^2 + b^2)^2$ (xiv) $(7x + 5)^2$
(xv) $(0.9p - 0.5q)^2$

Solution.

(i) We have,

$$\begin{aligned}(xy + yz)^2 &= (xy)^2 + (yz)^2 + 2 \times xy \times yz \\ &= x^2y^2 + y^2z^2 + 2xy^2z\end{aligned}$$

[using the identity, $(a + b)^2 = a^2 + b^2 + 2ab$]

(ii) We have,

$$\begin{aligned}(x^2y - xy^2)^2 &= (x^2y)^2 + (xy^2)^2 - 2x^2y \times xy^2 \\ &= x^4y^2 + x^2y^4 - 2x^3y^3\end{aligned}$$

[using the identity, $(a - b)^2 = a^2 + b^2 - 2ab$]

(iii) We have,

$$\begin{aligned}\left(\frac{4}{5}a + \frac{5}{4}b\right)^2 &= \left(\frac{4}{5}a\right)^2 + \left(\frac{5}{4}b\right)^2 + 2 \times \frac{4}{5}a \times \frac{5}{4}b \\ &= \frac{16}{25}a^2 + \frac{25}{16}b^2 + 2ab\end{aligned}$$

[using the identity, $(a + b)^2 = a^2 + b^2 + 2ab$]

(iv) We have,

$$\begin{aligned}\left(\frac{2}{4}x - \frac{3}{2}y\right)^2 &= \left(\frac{2}{3}x\right)^2 + \left(\frac{3}{2}y\right)^2 - 2 \times \frac{2}{3}x \times \frac{3}{2}y \\ &= \frac{4}{9}x^2 + \frac{9}{4}y^2 - 2xy\end{aligned}$$

[using the identity, $(a - b)^2 = a^2 + b^2 - 2ab$]

(v) We have,

$$\begin{aligned}\left(\frac{4}{5}p + \frac{5}{3}q\right)^2 &= \left(\frac{4}{5}p\right)^2 + \left(\frac{5}{3}q\right)^2 + 2 \times \frac{4}{5}p \times \frac{5}{3}q \\ &= \frac{16}{25}p^2 + \frac{25}{9}q^2 + \frac{8}{3}pq\end{aligned}$$

[using the identity, $(a + b)^2 = a^2 + b^2 + 2ab$]

(vi) We have,

$$\begin{aligned}(x + 3)(x + 7) &= x^2 + (3 + 7)x + 3 \times 7 \\ &= x^2 + 10x + 21\end{aligned}$$

[using the identity, $(x + a)(x + b) = x^2 + (a + b)x + ab$]

(vii) We have,

$$\begin{aligned}(2x + 9)(2x - 7) &= (2x + 9)(2x + (-7)) \\ &= (2x)^2 + [9 + (-7)]2x + 9 \times (-7) \\ &= 4x^2 + 4x - 63\end{aligned}$$

[using the identity, $(x + a)(x + b) = x^2 + (a + b)x + ab$]

(vii) We have,

$$\begin{aligned}\left(\frac{4x}{5} + \frac{y}{4}\right)\left(\frac{4x}{5} + \frac{3y}{4}\right) &= \left(\frac{4x}{5}\right)^2 + \left(\frac{y}{4} + \frac{3y}{4}\right)\frac{4x}{5} + \frac{y}{4} \times \frac{3y}{4} \\ & \quad \text{[using the identity, } (x+a)(x+b) = x^2 + (a+b)x + ab\text{]} \\ &= \frac{16}{25}x^2 + \frac{4xy}{5} + \frac{3y^2}{16}\end{aligned}$$

(ix) We have,

$$\begin{aligned}\left(\frac{2x}{3} - \frac{2}{3}\right)\left(\frac{2x}{3} + \frac{2a}{3}\right) &= \left(\frac{2x}{3}\right)^2 + \left(\frac{-2}{3} + \frac{2a}{3}\right)\frac{2x}{3} + \left(\frac{-2}{3} \times \frac{2a}{3}\right) \\ & \quad \text{[using the identity, } (x+a)(x+b) = x^2 + (a+b)x + ab\text{]} \\ &= \frac{4x^2}{9} + \frac{2a-2}{3} \times \frac{2}{3}x - \frac{4}{9}a = \frac{4x^2}{9} + \frac{4}{9}(a-1)x - \frac{4}{9}a\end{aligned}$$

(x) We have,

$$\begin{aligned}(2x-5y)(2x-5y) &= (2x-5y)^2 \\ &= (4x)^2 + (5y)^2 - 2 \times 2x \times 5y \\ & \quad \text{[using the identity, } (a-b)^2 = a^2 + b^2 - 2ab\text{]} \\ &= 16x^2 + 25y^2 - 20xy\end{aligned}$$

(xi) We have,

$$\begin{aligned}\left(\frac{2a}{3} + \frac{b}{3}\right)\left(\frac{2a}{3} - \frac{b}{3}\right) &= \left(\frac{2a}{3}\right)^2 - \left(\frac{b}{3}\right)^2 \\ & \quad \text{[using the identity, } (a+b)(a-b) = a^2 - b^2\text{]} \\ &= \frac{4}{9}a^2 - \frac{1}{9}b^2\end{aligned}$$

(xii) We have,

$$\begin{aligned}(x^2 + y^2)(x^2 - y^2) &= (x^2)^2 - (y^2)^2 \\ & \quad \text{[using the identity, } (a+b)(a-b) = a^2 - b^2\text{]} \\ &= x^4 - y^4\end{aligned}$$

(xiii) We have,

$$\begin{aligned}(a^2 + b^2)^2 &= (a^2)^2 + (b^2)^2 + 2a^2 \times b^2 \\ &= a^4 + b^4 + 2a^2b^2 \\ & \quad \text{[using the identity, } (a+b)^2 = a^2 + b^2 + 2ab\text{]}\end{aligned}$$

(xiv) We have,

$$\begin{aligned}(7x + 5)^2 &= (7x)^2 + 5^2 + 2 \times 7x \times 5 \\ &= 49x^2 + 25 + 70x \\ & \quad \text{[using the identity, } (a+b)^2 = a^2 + b^2 + 2ab\text{]}\end{aligned}$$

(xv) We have,

$$\begin{aligned}(0.9p - 0.5q)^2 &= (0.9p)^2 + (0.5q)^2 - 2 \times 0.9p \times 0.5q \\ & \quad \text{[using the identity, } (a-b)^2 = a^2 + b^2 - 2ab\text{]} \\ &= 0.81p^2 + 0.25q^2 - 0.9pq\end{aligned}$$

Question. 86 Using suitable identities, evaluate the following:

- | | |
|------------------------------|-----------------------------|
| (i) $(52)^2$ | (ii) $(49)^2$ |
| (iii) $(103)^2$ | (iv) $(98)^2$ |
| (v) $(1005)^2$ | (vi) $(995)^2$ |
| (vii) 47×53 | (viii) 52×53 |
| (ix) 105×95 | (x) 104×97 |
| (xi) 101×103 | (xii) 98×103 |
| (xiii) $(9.9)^2$ | (xiv) 9.8×10.2 |
| (xv) 10.1×10.2 | (xvi) $(35.4)^2 - (14.6)^2$ |
| (xvii) $(69.3)^2 - (30.7)^2$ | (xviii) $(9.7)^2 - (0.3)^2$ |
| (xix) $(132)^2 - (68)^2$ | (xx) $(339)^2 - (161)^2$ |
| (xxi) $(729)^2 - (271)^2$ | |

Solution.

(i) We have,

$$\begin{aligned}
 (52)^2 &= (50 + 2)^2 \\
 &= (50)^2 + (2)^2 + 2 \times 50 \times 2 \quad [\text{using the identity, } (a + b)^2 = a^2 + b^2 + 2ab] \\
 &= 2500 + 4 + 200 \\
 &= 2704
 \end{aligned}$$

(ii) We have,

$$\begin{aligned}
 (49)^2 &= (50 - 1)^2 \\
 &= (50)^2 + 1^2 - 2 \times 50 \times 1 \quad [\text{using the identity, } (a - b)^2 = a^2 + b^2 - 2ab] \\
 &= 2500 + 1 - 100 \\
 &= 2401
 \end{aligned}$$

(iii) We have,

$$\begin{aligned}
 (103)^2 &= (100 + 3)^2 \\
 &= (100)^2 + 3^2 + 2 \times 100 \times 3 \\
 &= 10000 + 9 + 600 \\
 &= 10609 \quad [\text{using the identity, } (a + b)^2 = a^2 + b^2 + 2ab]
 \end{aligned}$$

(iv) We have,

$$\begin{aligned}
 (98)^2 &= (100 - 2)^2 \\
 &= (100)^2 + (2)^2 - 2 \times 100 \times 2 \\
 &= 10000 + 4 - 400 \\
 &= 9604 \quad [\text{using the identity, } (a - b)^2 = a^2 + b^2 - 2ab]
 \end{aligned}$$

(v) We have,

$$\begin{aligned}
 (1005)^2 &= (1000 + 5)^2 \\
 &= (1000)^2 + 5^2 + 2 \times 1000 \times 5 \\
 &= 1000000 + 25 + 10000 \\
 &= 1010025 \quad [\text{using the identity, } (a + b)^2 = a^2 + b^2 + 2ab]
 \end{aligned}$$

(vi) We have,

$$\begin{aligned}(995)^2 &= (1000 - 5)^2 \\ &= (1000)^2 + (5)^2 - 2 \times 1000 \times 5 \\ &= 1000000 + 25 - 10000 \\ &= 990025 \quad \text{[using the identity, } (a - b)^2 = a^2 + b^2 - 2ab\text{]}\end{aligned}$$

(vii) We have,

$$\begin{aligned}47 \times 53 &= (50 - 3)(50 + 3) \\ &= (50)^2 - (3)^2 \quad \text{[using the identity, } (a - b)(a + b) = a^2 - b^2\text{]} \\ &= 2500 - 9 \\ &= 2491\end{aligned}$$

viii) We have,

$$\begin{aligned}52 \times 53 &= (50 + 2)(50 + 3) \\ &= (50)^2 + (2 + 3)50 + 2 \times 3 \\ & \quad \text{[using the identity, } (x + a)(x + b) = x^2 + (a + b)x + ab\text{]} \\ &= 2500 + 250 + 6 = 2756\end{aligned}$$

(ix) We have,

$$\begin{aligned}105 \times 95 &= (100 + 5)(100 - 5) \\ &= (100)^2 - (5)^2 \quad \text{[using the identity, } (a + b)(a - b) = a^2 - b^2\text{]} \\ &= 10000 - 25 \\ &= 9975\end{aligned}$$

(x) We have,

$$\begin{aligned}104 \times 97 &= (100 + 4)(100 - 3) \\ &= (100)^2 + (4 - 3)100 + 4 \times (-3) \\ &= 10000 + 100 - 12 \\ &= 10088 \quad \text{[using the identity, } (x + a)(x + b) = x^2 + (a + b)x + ab\text{]}\end{aligned}$$

(xi) We have,

$$\begin{aligned}101 \times 103 &= (100 + 1)(100 + 3) \\ &= (100)^2 + (1 + 3)100 + 3 \times 1 \\ &= 10000 + 400 + 3 \\ &= 10403 \quad \text{[using the identity, } (x + a)(x + b) = x^2 + (a + b)x + ab\text{]}\end{aligned}$$

(xii) We have,

$$\begin{aligned}98 \times 103 &= (100 - 2)(100 + 3) \\ &= (100)^2 + (-2 + 3)100 + (-2) \times 3 \\ &= 10000 + 100 - 6 \\ &= 10094 \quad \text{[using the identity, } (x + a)(x + b) = x^2 + (a + b)x + ab\text{]}\end{aligned}$$

(xiii) We have,

$$\begin{aligned}(9.9)^2 &= (10 - 0.1)^2 \\ &= 10^2 + (0.1)^2 - 2 \times 10 \times 0.1 \\ &= 100 + 0.01 - 2 = 98.01 \\ & \quad \text{[using the identity, } (a - b)^2 = a^2 + b^2 - 2ab\text{]}\end{aligned}$$

(xiv) We have,

$$\begin{aligned}9.8 \times 10.2 &= (10 - 0.2)(10 + 0.2) \\ &= 10^2 - (0.2)^2 \\ &= 100 - 0.04 \\ &= 100 - 0.04 \\ &= 99.96\end{aligned}$$

[using the identity, $(a + b)(a - b) = a^2 - b^2$]

(xv) We have,

$$\begin{aligned}10.1 \times 10.2 &= (10 + 0.1)(10 + 0.2) \\ &= (10)^2 + (0.1 + 0.2)10 + (0.1)(0.2) \\ &= 100 + 0.3 \times 10 + 0.02 \\ &= 103.02\end{aligned}$$

[using the identity, $(x + a)(x + b) = x^2 + (a + b)x + ab$]

(xvi) We have,

$$\begin{aligned}(35.4)^2 - (14.6)^2 &= (35.4 + 14.6)(35.4 - 14.6) \\ &= 50 \times 20.8 \\ &= 1040\end{aligned}$$

[using the identity, $(a + b)(a - b) = a^2 - b^2$]

(xvii) We have,

$$\begin{aligned}(69.3)^2 - (30.7)^2 &= (69.3 + 30.7)(69.3 - 30.7) \\ &= 100 \times 38.6 \\ &= 3860\end{aligned}$$

[using the identity, $(a + b)(a - b) = a^2 - b^2$]

(xviii) We have,

$$\begin{aligned}(9.7)^2 - (0.3)^2 &= (9.7 + 0.3)(9.7 - 0.3) \\ &= 10 \times 9.4 \\ &= 94\end{aligned}$$

[using the identity, $a^2 - b^2 = (a + b)(a - b)$]

(xix) We have,

$$\begin{aligned}(132)^2 - (68)^2 &= (132 + 68)(132 - 68) \\ &= 200 \times 64 \\ &= 12800\end{aligned}$$

[using the identity, $a^2 - b^2 = (a + b)(a - b)$]

(xx) We have,

$$\begin{aligned}(339)^2 - (161)^2 &= (339 + 161)(339 - 161) \\ &= 500 \times 178 \\ &= 89000\end{aligned}$$

[using the identity, $a^2 - b^2 = (a + b)(a - b)$]

(xxi) We have,

$$\begin{aligned}(729)^2 - (271)^2 &= (729 + 271)(729 - 271) \\ &= 1000 \times 458 \\ &= 458000\end{aligned}$$

[using the identity, $a^2 - b^2 = (a + b)(a - b)$]

Question. 87 Write the greatest common factor in each of the following terms.

(i) $-18a^2, 108a$

(ii) $3x^2y, 18xy^2, -6xy$

(iii) $2xy, -y^2, 2x^2y$

(iv) $l^2m^2n, lm^2n^2, l^2mn^2$

(v) $21pqr, -7p^2q^2r^2, 49p^2qr$

(vi) $qrx, pryz, rxyz$

(vii) $3x^3y^2z, -6xy^3z^2, 12x^2yz^3$

(viii) $63p^2a^2r^2s, -9pq^2r^2s^2, 15p^2qr^2s^2, -60p^2a^2rs^2$

(ix) $13x^2y, 169xy$

(x) $11x^2, 12y^2$

Solution.

(i) We have,

$$-18a^2 = -18 \times a \times a$$

$$108a = 18 \times 10 \times a$$

∴ The greatest common factor i.e. GCF is $18a$.

(ii) We have,

$$3x^2y = 3 \times x \times x \times y$$

$$18xy^2 = 3 \times 6 \times x \times y \times y$$

$$-6xy = -1 \times 3 \times 2 \times x \times y$$

∴ GCF = $3xy$

(iii) We have,

$$2xy = 2 \times x \times y$$

$$-y^2 = -1 \times y \times y$$

$$2x^2y = 2 \times x \times x \times y$$

∴ GCF = y

(iv) We have,

$$l^2m^2n = l \times l \times m \times m \times n$$

$$lm^2n^2 = l \times m \times m \times n \times n$$

$$l^2mn^2 = l \times l \times m \times n \times n$$

∴ GCF = lmn

(v) We have,

$$21pqr = 7 \times 3 \times p \times q \times r$$

$$-7p^2q^2r^2 = -7 \times p \times p \times q \times q \times r \times r$$

$$49p^2qr = 7 \times 7 \times p \times p \times q \times r$$

∴ GCF = $7pqr$

(vi) We have,

$$qrx = q \times r \times x$$

$$pry = p \times r \times y$$

$$rxz = r \times x \times z$$

∴ GCF = rx

(vii) We have,

$$3x^3y^2z = 3 \times x \times x \times x \times y \times y \times z$$

$$-6xy^3z^2 = -3 \times 2 \times x \times y \times y \times y \times z \times z$$

$$12x^2yz^3 = 3 \times 4 \times x \times x \times y \times z \times z \times z$$

∴ GCF = $3xyz$

(viii) We have,

$$63p^2a^2r^2s = 3 \times 3 \times 7 \times p \times p \times a \times a \times r \times r \times s$$

$$-9pq^2r^2s^2 = -3 \times 3 \times p \times q \times q \times r \times r \times s \times s$$

$$15p^2qr^2s^2 = 3 \times 5 \times p \times p \times q \times r \times r \times s \times s$$

$$-80p^2a^2rs^2 = -2 \times 2 \times 3 \times 5 \times p \times p \times a \times a \times r \times s \times s$$

∴ GCF = $3prs$

(ix) We have,

$$13x^2y = 13 \times x \times x \times y$$

$$169xy = 13 \times 13 \times x \times y$$

∴ GCF = $13xy$

(x) We have, $11x^2$, $12y^2$

The GCF of 11, 12 is 1.

Also, there is no common factor between x^2 and y^2 .

Hence, the GCF of $11x^2$ and $12y^2$ is 1.

Question. 88 Factorise the following expressions.

- (i) $6ab + 12bc$
(ii) $-xy - ay$
(iii) $ax^3 - bx^2 + cx$
(iv) $l^2m^2n - lm^2n^2 - l^2mn^2$
(v) $3pqr - 6p^2q^2r^2 - 15r^2$
(vi) $x^3y^2 + x^2y^3 - xy^4 + xy$
(vii) $4xy^2 - 10x^2y + 16x^2y^2 + 2xy$
(viii) $2a^3 - 3a^2b + 5ab^2 - ab$
(ix) $63p^2q^2r^2s - 9pq^2r^2s^2 + 15p^2qr^2s^2 - 60p^2q^2rs^2$
(x) $24x^2yz^3 - 6xy^3z^2 + 15x^2y^2z - 5xyz$
(xi) $a^3 + a^2 + a + 1$
(xii) $lx + my + mx + ly$
(xiii) $a^3x - x^4 + a^2x^2 - ax^3$
(xiv) $2x^2 - 2y + 4xy - x$
(xv) $y^2 + 8zx - 2xy - 4yz$
(xvi) $ax^2y - bxyz - ax^2z + bxy^2$
(xvii) $a^2b + a^2c + ab + ac + b^2c + c^2b$
(xviii) $2ax^2 + 4axy + 3bx^2 + 2ay^2 + 6bxy + 3by^2$

Solution.

(i) We have,

$$6ab + 12bc = 6ab + 6 \times 2 \times bc = 6b(a + 2c)$$

(ii) We have,

$$-xy - ay = -y(x + a)$$

(iii) We have,

$$ax^3 - bx^2 + cx = x(ax^2 - bx + c)$$

(iv) We have,

$$l^2m^2n - lm^2n^2 - l^2mn^2 = lmn(lm - mn - ln)$$

(v) We have, $3pqr - 6p^2q^2r^2 - 15r^2$

$$= 3pqr - 3 \times 2p^2q^2r^2 - 3 \times 5r^2 = 3r(pq - 2p^2q^2r - 5r)$$

(vi) We have, $x^3y^2 + x^2y^3 - xy^4 + xy$

$$= xy(x^2y + xy^2 - y^3 + 1)$$

(vii) We have, $4xy^2 - 10x^2y + 16x^2y^2 + 2xy$

$$= 2 \times 2xy^2 - 2 \times 5 \times x^2y + 2 \times 8 \times x^2y^2 + 2xy \\ = 2xy(2y - 5x + 8xy + 1)$$

(viii) We have, $2a^3 - 3a^2b + 5ab^2 - ab$

$$= a(2a^2 - 3ab + 5b^2 - b)$$

(ix) We have, $63p^2q^2r^2s - 9pq^2r^2s^2 + 15p^2qr^2s^2 - 60p^2q^2rs^2$

$$= 3 \times 21p^2q^2r^2s - 3 \times 3pq^2r^2s^2 + 3 \times 5p^2qr^2s^2 - 3 \times 20p^2q^2rs^2 \\ = 3pqrs(21pqr - 3qrs + 5prs - 20pqs)$$

(x) We have, $24x^2yz^3 - 6xy^3z^2 + 15x^2y^2z - 5xyz$

$$= xyz(24xz^2 - 6y^2z + 15xy - 5)$$

$$(x) \text{ We have, } 24x^2yz^3 - 6xy^3z^2 + 15x^2y^2z - 5xyz$$

$$= xyz(24xz^2 - 6y^2z + 15xy - 5)$$

$$(xi) \text{ We have, } a^3 + a^2 + a + 1$$

$$= a^2(a+1) + 1(a+1) = (a+1)(a^2+1)$$

$$(xii) \text{ We have, } lx + my + mx + ly$$

$$= lx + mx + my + ly = x(l+m) + y(m+l) = (l+m)(x+y)$$

$$(xiii) \text{ We have, } a^3x - x^4 + a^2x^2 - ax^3$$

$$= x(a^3 - x^3 + a^2x - ax^2) = x(a^3 + a^2x - x^3 - ax^2)$$

$$= x[a^2(a+x) - x^2(x+a)]$$

$$= x[(x+a)(a^2 - x^2)] = x(a^2 - x^2)(a+x)$$

$$(xiv) \text{ We have, } 2x^2 - 2y + 4xy - x$$

$$= 2x^2 - x - 2y + 4xy = x(2x-1) - 2y(1-2x)$$

$$= x(2x-1) + 2y(2x-1) = (2x-1)(x+2y)$$

$$(xv) \text{ We have, } y^2 + 8zx - 2xy - 4yz$$

$$= y^2 - 2xy + 8zx - 4yz = y(y-2x) - 4z(y-2x)$$

$$= (y-2x)(y-4z)$$

$$(xvi) \text{ We have, } ax^2y - bxyz - ax^2z + bxy^2$$

$$= x(axy - byz - axz + by^2)$$

$$= x(axy - axz - byz + by^2)$$

$$= x[ax(y-z) + by(-z+y)]$$

$$= x[(ax+by)(y-z)]$$

$$(xvii) \text{ We have, } a^2b + a^2c + ab + ac + b^2c + c^2b$$

$$= (a^2b + ab + b^2c) + (a^2c + ac + c^2b)$$

$$= b(a^2 + a + bc) + c(a^2 + a + bc)$$

$$= (a^2 + a + bc)(b+c)$$

$$(xviii) \text{ We have, } 2ax^2 + 4axy + 3bx^2 + 2ay^2 + 6bxy + 3by^2$$

$$= (2ax^2 + 2ay^2 + 4axy) + (3bx^2 + 3by^2 + 6bxy)$$

$$= 2a(x^2 + y^2 + 2xy) + 3b(x^2 + y^2 + 2xy)$$

$$= (2a+3b)(x+y)^2$$

Question. 89 Factorise the following, using the identity, $a^2 + 2ab - b^2 = (a+b)^2$

$$(i) x^2 + 6x + 9$$

$$(ii) x^2 + 12x + 36$$

$$(iii) x^2 + 14x + 49$$

$$(iv) x^2 + 2x + 1$$

$$(v) 4x^2 + 4x + 1$$

$$(vi) a^2x^2 + 2ax + 1$$

$$(vii) a^2x^2 + 2abx + b^2$$

$$(viii) a^2x^2 + 2abxy + b^2y^2$$

$$(ix) 4x^2 + 12x + 9$$

$$(x) 16x^2 + 40x + 25$$

$$(xi) 9x^2 + 24x + 16$$

$$(xii) 9x^2 + 30x + 25$$

$$(xiii) 2x^3 + 24x^2 + 72x$$

$$(xiv) a^2x^3 + 2abx^2 + b^2x$$

$$(xv) 4x^4 + 12x^3 + 9x^2$$

$$(xvi) \frac{x^2}{4} + 2x + 4$$

$$(xvii) 9x^2 + 2xy + \frac{y^2}{9}$$

Solution.

(i) We have,

$$\begin{aligned}x^2 + 6x + 9 &= x^2 + 2 \cdot 3 \cdot x + 3^2 \\ &= (x + 3)^2 && [\because a^2 + 2ab + b^2 = (a + b)^2] \\ &= (x + 3)(x + 3)\end{aligned}$$

(ii) We have, $x^2 + 12x + 36$

$$\begin{aligned}&= x^2 + 2 \cdot 6 \cdot x + 6^2 \\ &= (x + 6)^2 && [\because a^2 + 2ab + b^2 = (a + b)^2] \\ &= (x + 6)(x + 6)\end{aligned}$$

(iii) We have, $x^2 + 14x + 49$

$$= x^2 + 2 \cdot 7 \cdot x + 7^2 = (x + 7)^2 = (x + 7)(x + 7)$$

(iv) We have, $x^2 + 2x + 1$

$$= x^2 + 2 \cdot 1 \cdot x + 1^2 = (x + 1)^2 = (x + 1)(x + 1)$$

(v) We have, $4x^2 + 4x + 1$

$$= (2x)^2 + 2 \cdot 2x \cdot 1 + 1^2 = (2x + 1)^2 = (2x + 1)(2x + 1)$$

(vi) We have, $a^2x^2 + 2ax + 1$

$$= (ax)^2 + 2 \cdot ax \cdot 1 + (1)^2 = (ax + 1)^2 = (ax + 1)(ax + 1)$$

(vii) We have, $a^2x^2 + 2abx + b^2$

$$= (ax)^2 + 2 \cdot ax \cdot b + b^2 = (ax + b)^2 = (ax + b)(ax + b)$$

(viii) We have, $a^2x^2 + 2abxy + b^2y^2$

$$= (ax)^2 + 2 \cdot ax \cdot by + (by)^2 = (ax + by)^2 = (ax + by)(ax + by)$$

(ix) We have, $4x^2 + 12x + 9$

$$= (2x)^2 + 2 \cdot 2x \cdot 3 + 3^2 = (2x + 3)^2 = (2x + 3)(2x + 3)$$

(x) We have, $16x^2 + 40x + 25$

$$= (4x)^2 + 2 \cdot 4x \cdot 5 + 5^2 = (4x + 5)^2 = (4x + 5)(4x + 5)$$

(xi) We have, $9x^2 + 24x + 16$

$$= (3x)^2 + 2 \cdot 3x \cdot 4 + 4^2 = (3x + 4)^2 = (3x + 4)(3x + 4)$$

(xii) We have, $9x^2 + 30x + 25$

$$= (3x)^2 + 2 \cdot 3x \cdot 5 + 5^2 = (3x + 5)^2 = (3x + 5)(3x + 5)$$

(xiii) We have, $2x^3 + 24x^2 + 72x$

$$\begin{aligned}&= 2x(x^2 + 12x + 36) = 2x(x^2 + 2 \cdot 6 \cdot x + 6^2) \\ &= 2x(x + 6)^2 = 2x(x + 6)(x + 6)\end{aligned}$$

(xiv) We have, $a^2x^3 + 2abx^2 + b^2x$

$$\begin{aligned}&= x(a^2x^2 + 2abx + b^2) = x[(ax)^2 + 2 \cdot ax \cdot b + b^2] \\ &= x(ax + b)^2 = x(ax + b)(ax + b)\end{aligned}$$

(xv) We have, $4x^4 + 12x^3 + 9x^2$

$$\begin{aligned}&= x^2(4x^2 + 12x + 9) = x^2[(2x)^2 + 2 \cdot 2x \cdot 3 + 3^2] \\ &= x^2(2x + 3)^2 = x^2(2x + 3)(2x + 3)\end{aligned}$$

(xvi) We have, $\frac{x^2}{4} + 2x + 4$

$$= \left(\frac{x}{2}\right)^2 + 2 \cdot \frac{x}{2} \cdot 2 + 2^2 = \left(\frac{x}{2} + 2\right)^2 = \left(\frac{x}{2} + 2\right)\left(\frac{x}{2} + 2\right)$$

(xvii) We have, $9x^2 + 2xy + \frac{y^2}{9}$

$$= (3x)^2 + 2 \cdot 3x \cdot \frac{y}{3} + \left(\frac{y}{3}\right)^2 = \left(3x + \frac{y}{3}\right)^2 = \left(3x + \frac{y}{3}\right)\left(3x + \frac{y}{3}\right)$$

Question. 90 Factorise the following, using the identity, $a^2 - 2ab + b^2 = (a - b)^2$

(i) $x^2 - 8x + 16$	(ii) $x^2 - 10x + 25$
(iii) $y^2 - 14y + 49$	(iv) $p^2 - 2p + 1$
(v) $4a^2 - 4ab + b^2$	(vi) $p^2y^2 - 2py + 1$
(vii) $a^2y^2 - 2aby + b^2$	(viii) $9x^2 - 12x + 4$
(ix) $4y^2 - 12y + 9$	(x) $\frac{x^2}{4} - 2x + 4$
(xi) $a^2y^3 - 2aby^2 + b^2y$	(xii) $9y^2 - 4xy + \frac{4x^2}{9}$

Solution.

(i) We have,

$$\begin{aligned} x^2 - 8x + 16 &= x^2 - 2 \cdot x \cdot 4 + 4^2 \\ &= (x - 4)^2 && [\because a^2 - 2ab + b^2 = (a - b)^2] \\ &= (x - 4)(x - 4) \end{aligned}$$

(ii) We have,

$$\begin{aligned} x^2 - 10x + 25 &= x^2 - 2 \cdot x \cdot 5 + 5^2 \\ &= (x - 5)^2 = (x - 5)(x - 5) \end{aligned}$$

(iii) We have,

$$\begin{aligned} y^2 - 14y + 49 &= y^2 - 2 \cdot y \cdot 7 + 7^2 \\ &= (y - 7)^2 = (y - 7)(y - 7) \end{aligned}$$

(iv) We have,

$$\begin{aligned} p^2 - 2p + 1 &= p^2 - 2 \cdot p \cdot 1 + 1^2 \\ &= (p - 1)^2 = (p - 1)(p - 1) \end{aligned}$$

(v) We have,

$$\begin{aligned} 4a^2 - 4ab + b^2 &= (2a)^2 - 2 \cdot 2a \cdot b + b^2 \\ &= (2a - b)^2 = (2a - b)(2a - b) \end{aligned}$$

(vi) We have,

$$\begin{aligned} p^2y^2 - 2py + 1 &= (py)^2 - 2 \cdot py \cdot 1 + 1^2 \\ &= (py - 1)^2 = (py - 1)(py - 1) \end{aligned}$$

(vii) We have,

$$\begin{aligned} a^2y^2 - 2aby + b^2 &= (ay)^2 - 2 \cdot ay \cdot b + b^2 \\ &= (ay - b)^2 = (ay - b)(ay - b) \end{aligned}$$

(viii) We have,

$$\begin{aligned} 9x^2 - 12x + 4 &= (3x)^2 - 2 \cdot 3x \cdot 2 + 2^2 \\ &= (3x - 2)^2 = (3x - 2)(3x - 2) \end{aligned}$$

(ix) We have,

$$\begin{aligned} 4y^2 - 12y + 9 &= (2y)^2 - 2 \cdot 2y \cdot 3 + 3^2 \\ &= (2y - 3)^2 = (2y - 3)(2y - 3) \end{aligned}$$

(x) We have,

$$\begin{aligned} \frac{x^2}{4} - 2x + 4 &= \left(\frac{x}{2}\right)^2 - 2 \cdot \frac{x}{2} \cdot 2 + 2^2 \\ &= \left(\frac{x}{2} - 2\right)^2 = \left(\frac{x}{2} - 2\right)\left(\frac{x}{2} - 2\right) \end{aligned}$$

(xi) We have,

$$\begin{aligned} a^2y^3 - 2aby^2 + b^2y &= y(a^2y^2 - 2aby + b^2) = y[(ay)^2 - 2 \cdot ay \cdot b + b^2] \\ &= y(ay - b)^2 = y(ay - b)(ay - b) \end{aligned}$$

(xii) We have,

$$\begin{aligned} 9y^2 - 4xy + \frac{4x^2}{9} &= (3y)^2 - 2 \cdot 3y \cdot \frac{2x}{3} + \left(\frac{2x}{3}\right)^2 \\ &= \left(3y - \frac{2x}{3}\right)^2 = \left(3y - \frac{2x}{3}\right)\left(3y - \frac{2x}{3}\right) \end{aligned}$$

Question. 91 Factorise the following

(i) $x^2 + 15x + 26$

(ii) $x^2 + 9x + 20$

(iii) $y^2 + 18y + 65$

(iv) $p^2 + 14p + 13$

(v) $y^2 + 4y - 21$

(vi) $y^2 - 2y - 15$

(vii) $18 + 11x + x^2$

(viii) $x^2 - 10x + 21$

(ix) $x^2 - 17x + 60$

(x) $x^2 + 4x - 77$

(xi) $y^2 + 7y + 12$

(xii) $p^2 - 13p - 30$

(xiii) $p^2 - 16p - 80$

Solution.

(i) We have, $x^2 + 15x + 26$

$$= x^2 + 2x + 13x + 2 \times 13 = x(x + 2) + 13(x + 2) = (x + 2)(x + 13)$$

(ii) We have, $x^2 + 9x + 20$

$$= x^2 + 5x + 4x + 5 \times 4 = x(x + 5) + 4(x + 5) = (x + 5)(x + 4)$$

(iii) We have, $y^2 + 18y + 65$

$$= y^2 + 13y + 5y + 5 \times 13 = y(y + 13) + 5(y + 13) = (y + 13)(y + 5)$$

(iv) We have, $p^2 + 14p + 13$

$$= p^2 + 13p + p + 13 \times 1 = p(p + 13) + 1(p + 13) = (p + 13)(p + 1)$$

(v) We have, $y^2 + 4y - 21$

$$= y^2 + (7 - 3)y - 21 = y^2 + 7y - 3y - 21 = y(y + 7) - 3(y + 7) = (y + 7)(y - 3)$$

(vi) We have, $y^2 - 2y - 15$

$$= y^2 + (3 - 5)y - 15 = y^2 + 3y - 5y - 15 = y(y + 3) - 5(y + 3) = (y + 3)(y - 5)$$

(vii) We have, $18 + 11x + x^2$

$$= x^2 + 11x + 18 = x^2 + (9 + 2)x + 18 = x^2 + 9x + 2x + 18$$

$$= x(x + 9) + 2(x + 9) = (x + 9)(x + 2)$$

(viii) We have, $x^2 - 10x + 21$

$$= x^2 - (7 + 3)x + 21 = x^2 - 7x - 3x + 21 = x(x - 7) - 3(x - 7)$$

$$= (x - 7)(x - 3)$$

(ix) We have, $x^2 - 17x + 60$

$$= x^2 - (12 + 5)x + 60 = x^2 - 12x - 5x + 60 = x(x - 12) - 5(x - 12)$$

$$= (x - 12)(x - 5)$$

(x) We have, $x^2 + 4x - 77$

$$x^2 + (11 - 7)x - 77 = x^2 + 11x - 7x - 77 = x(x + 11) - 7(x + 11)$$

$$= (x + 11)(x - 7)$$

(xi) We have, $y^2 + 7y + 12$

$$y^2 + (4 + 3)y + 12 = y^2 + 4y + 3y + 12 = y(y + 4) + 3(y + 4) = (y + 4)(y + 3)$$

(xii) We have, $p^2 - 13p - 30$

$$= p^2 - (15 - 2)p - 30 = p^2 - 15p + 2p - 30 = p(p - 15) + 2(p - 15)$$

$$= (p - 15)(p + 2)$$

(xiii) We have, $p^2 - 16p - 80$

$$= p^2 - (20 - 4)p - 80 = p^2 - 20p + 4p - 80 = p(p - 20) + 4(p - 20)$$

$$= (p - 20)(p + 4)$$

Question. 92 Factorise the following using the identity, $a^2 - b^2 = (a+b)(a-b)$.

- (i) $x^2 - 9$ (ii) $4x^2 - 25y^2$
 (iii) $4x^2 - 49y^2$ (iv) $3a^2b^3 - 27a^4b$
 (v) $28ay^2 - 175ax^2$ (vi) $9x^2 - 1$
 (vii) $25ax^2 - 25a$ (viii) $\frac{x^2}{9} - \frac{y^2}{25}$
 (ix) $\frac{2p^2}{25} - 32q^2$ (x) $49x^2 - 36y^2$
 (xi) $y^3 - \frac{y}{9}$ (xii) $\frac{x^2}{25} - 625$
 (xiii) $\frac{x^2}{8} - \frac{y^2}{18}$ (xiv) $\frac{4x^2}{9} - \frac{9y^2}{16}$
 (xv) $\frac{x^3y}{9} - \frac{xy^3}{16}$ (xvi) $1331x^3y - 11y^3x$
 (xvii) $\frac{1}{36}a^2b^2 - \frac{16}{49}b^2c^2$ (xviii) $a^4 - (a-b)^4$
 (xix) $x^4 - 1$ (xx) $y^4 - 625$
 (xxi) $p^5 - 16p$ (xxii) $16x^4 - 81$
 (xxiii) $x^4 - y^4$ (xxiv) $y^4 - 81$
 (xxv) $16x^4 - 625y^4$ (xxvi) $(a-b)^2 - (b-c)^2$
 (xxvii) $(x+y)^4 - (x-y)^4$ (xxviii) $x^4 - y^4 + x^2 - y^2$
 (xxix) $8a^3 - 2a$ (xxx) $x^2 - \frac{y^2}{100}$
 (xxxix) $9x^2 - (3y+z)^2$

Solution.

(i) We have,

$$x^2 - 9 = x^2 - 3^2 = (x-3)(x+3)$$

$$[\because a^2 - b^2 = (a-b)(a+b)]$$

(ii) We have,

$$4x^2 - 25y^2 = (2x)^2 - (5y)^2 = (2x-5y)(2x+5y)$$

(iii) We have,

$$4x^2 - 49y^2 = (2x)^2 - (7y)^2 = (2x-7y)(2x+7y)$$

(iv) We have,

$$\begin{aligned} 3a^2b^3 - 27a^4b &= 3a^2b(b^2 - 9a^2) = 3a^2b[b^2 - (3a)^2] \\ &= 3a^2b(b+3a)(b-3a) \end{aligned}$$

(v) We have,

$$\begin{aligned}28ay^2 - 175ax^2 &= 7a(4y^2 - 25x^2) \\ &= 7a[(2y)^2 - (5x)^2] = 7a(2y - 5x)(2y + 5x)\end{aligned}$$

(vi) We have,

$$9x^2 - 1 = (3x)^2 - 1^2 = (3x - 1)(3x + 1)$$

(vii) We have,

$$25ax^2 - 25a = 25a(x^2 - 1^2) = 25a(x - 1)(x + 1)$$

(viii) We have,

$$\frac{x^2}{9} - \frac{y^2}{25} = \left(\frac{x}{3}\right)^2 - \left(\frac{y}{5}\right)^2 = \left(\frac{x}{3} - \frac{y}{5}\right)\left(\frac{x}{3} + \frac{y}{5}\right)$$

(ix) We have,

$$\frac{2p^2}{25} - 32q^2 = 2\left(\frac{p^2}{25} - 16q^2\right) = 2\left[\left(\frac{p}{5}\right)^2 - (4q)^2\right] = 2\left(\frac{p}{5} + 4q\right)\left(\frac{p}{5} - 4q\right)$$

(x) We have,

$$49x^2 - 36y^2 = (7x)^2 - (6y)^2 = (7x - 6y)(7x + 6y)$$

(xi) We have,

$$y^3 - \frac{y}{9} = y\left(y^2 - \frac{1}{9}\right) = y\left[y^2 - \left(\frac{1}{3}\right)^2\right] = y\left(y + \frac{1}{3}\right)\left(y - \frac{1}{3}\right)$$

(xii) We have,

$$\frac{x^2}{25} - 625 = \left(\frac{x}{5}\right)^2 - (25)^2 = \left(\frac{x}{5} - 25\right)\left(\frac{x}{5} + 25\right)$$

(xiii) We have,

$$\begin{aligned}\frac{x^2}{8} - \frac{y^2}{18} &= \frac{1}{2}\left(\frac{x^2}{4} - \frac{y^2}{9}\right) = \frac{1}{2}\left[\left(\frac{x}{2}\right)^2 - \left(\frac{y}{3}\right)^2\right] \\ &= \frac{1}{2}\left(\frac{x}{2} + \frac{y}{3}\right)\left(\frac{x}{2} - \frac{y}{3}\right)\end{aligned}$$

(xiv) We have,

$$\frac{4x^2}{9} - \frac{9y^2}{16} = \left(\frac{2x}{3}\right)^2 - \left(\frac{3y}{4}\right)^2 = \left(\frac{2x}{3} + \frac{3y}{4}\right)\left(\frac{2x}{3} - \frac{3y}{4}\right)$$

(xv) We have,

$$\frac{x^3y}{9} - \frac{xy^3}{16} = xy\left(\frac{x^2}{9} - \frac{y^2}{16}\right) = xy\left[\left(\frac{x}{3}\right)^2 - \left(\frac{y}{4}\right)^2\right] = xy\left(\frac{x}{3} + \frac{y}{4}\right)\left(\frac{x}{3} - \frac{y}{4}\right)$$

(xvi) We have,

$$\begin{aligned}1331x^3y - 11y^3x &= (11)^3x^3y - 11y^3x = 11xy(11^2x^2 - y^2) \\ &= 11xy[(11x)^2 - y^2] = 11xy(11x + y)(11x - y)\end{aligned}$$

(xvii) We have,

$$\begin{aligned}\frac{1}{36}a^2b^2 - \frac{16}{49}b^2c^2 &= \left(\frac{ab}{6}\right)^2 - \left(\frac{4bc}{7}\right)^2 = \left(\frac{ab}{6} + \frac{4bc}{7}\right)\left(\frac{ab}{6} - \frac{4bc}{7}\right) \\ &= b^2\left(\frac{a}{6} + \frac{4c}{7}\right)\left(\frac{a}{6} - \frac{4c}{7}\right)\end{aligned}$$

(xviii) We have,

$$\begin{aligned}a^4 - (a - b)^4 &= (a^2)^2 - [(a - b)^2]^2 = [a^2 + (a - b)^2][a^2 - (a - b)^2] \\&= [a^2 + a^2 + b^2 - 2ab][a^2 - (a^2 + b^2 - 2ab)] \\&= [2a^2 + b^2 - 2ab][-b^2 + 2ab] \\&= (2a^2 + b^2 - 2ab)(2ab - b^2)\end{aligned}$$

(xix) We have,

$$\begin{aligned}x^4 - 1 &= (x^2)^2 - 1 = (x^2 + 1)(x^2 - 1) \\&= (x^2 + 1)(x + 1)(x - 1)\end{aligned}$$

(xx) We have,

$$\begin{aligned}y^4 - 625 &= (y^2)^2 - (25)^2 \\&= (y^2 + 25)(y^2 - 25) \\&= (y^2 + 25)(y^2 - 5^2) \\&= (y^2 + 25)(y + 5)(y - 5)\end{aligned}$$

(xxi) We have,

$$\begin{aligned}\rho^5 - 16\rho &= \rho(\rho^4 - 16) = \rho[(\rho^2)^2 - 4^2] \\&= \rho(\rho^2 + 4)(\rho^2 - 4) \\&= \rho(\rho^2 + 4)(\rho^2 - 2^2) \\&= \rho(\rho^2 + 4)(\rho + 2)(\rho - 2)\end{aligned}$$

(xxii) We have,

$$\begin{aligned}16x^4 - 81 &= (4x^2)^2 - 9^2 = (4x^2 + 9)(4x^2 - 9) \\&= (4x^2 + 9)[(2x)^2 - 3^2] \\&= (4x^2 + 9)(2x + 3)(2x - 3)\end{aligned}$$

(xxiii) We have,

$$\begin{aligned}x^4 - y^4 &= (x^2)^2 - (y^2)^2 \\&= (x^2 + y^2)(x^2 - y^2) \\&= (x^2 + y^2)(x + y)(x - y)\end{aligned}$$

(xxiv) We have,

$$\begin{aligned}y^4 - 81 &= (y^2)^2 - (9)^2 = (y^2 + 9)(y^2 - 9) \\&= (y^2 + 9)[(y)^2 - (3)^2] \\&= (y^2 + 9)(y + 3)(y - 3)\end{aligned}$$

(xxv) We have,

$$\begin{aligned}16x^4 - 625y^4 &= (4x^2)^2 - (25y^2)^2 = (4x^2 + 25y^2)(4x^2 - 25y^2) \\&= (4x^2 + 25y^2)[(2x)^2 - (5y)^2] \\&= (4x^2 + 25y^2)(2x + 5y)(2x - 5y)\end{aligned}$$

(xxvi) We have,

$$(a - b)^2 - (b - c)^2 = (a - b + b - c)(a - b - b + c)(a - c)(a - 2b + c)$$

(xxvii) We have,

$$\begin{aligned}(x + y)^4 - (x - y)^4 &= [(x + y)^2]^2 - [(x - y)^2]^2 \\&= [(x + y)^2 + (x - y)^2][(x + y)^2 - (x - y)^2]\end{aligned}$$

$$= (x^2 + y^2 + 2xy + x^2 + y^2 - 2xy)(x + y + x - y)(x + y - x + y)$$

$$= (2x^2 + 2y^2)(2x)(2y) = 2(x^2 + y^2)(2x)(2y) = 8xy(x^2 + y^2)$$

(xxviii) We have,

$$x^4 - y^4 + x^2 - y^2 = (x^2)^2 - (y^2)^2 + (x^2 - y^2) = (x^2 + y^2)(x^2 - y^2) + (x^2 - y^2)$$

$$= (x^2 - y^2)(x^2 + y^2 + 1) = (x + y)(x - y)(x^2 + y^2 + 1)$$

(xxix) We have,

$$8a^3 - 2a = 2a(4a^2 - 1)$$

$$= 2a[(2a)^2 - (1)^2] = 2a(2a + 1)(2a - 1)$$

(xxx) We have,

$$x^2 - \frac{y^2}{100} = x^2 - \left(\frac{y}{10}\right)^2 = \left(x + \frac{y}{10}\right)\left(x - \frac{y}{10}\right)$$

(xxxi) We have,

$$9x^2 - (3y + z)^2 = (3x)^2 - (3y + z)^2 = (3x + 3y + z)(3x - 3y - z)$$

Question. 93 The following expressions are the areas of rectangles. Find the possible lengths and breadths of these rectangles.

(i) $x^2 - 6x + 8$

(ii) $x^2 - 3x + 2$

(iii) $x^2 - 7x + 10$

(iv) $x^2 + 19x - 20$

(v) $x^2 + 9x + 20$

Solution.

(i) Given, area of a rectangle = $x^2 - 6x + 8$

Now, we have to find the possible length and breadth of the rectangle.

So, we factorise the given expression.

$$\text{i.e. } x^2 - 6x + 8 = x^2 - (4 + 2)x + 8 = x^2 - 4x - 2x + 8$$

$$= x(x - 4) - 2(x - 4) = (x - 4)(x - 2)$$

Since, area of a rectangle = Length \times Breadth. Hence, the possible length and breadth are $(x - 4)$ and $(x - 2)$.

(ii) We have,

$$\text{Area of rectangle} = x^2 - 3x + 2$$

$$= x^2 - (2 + 1)x + 2 = x^2 - 2x - x + 2$$

$$= x(x - 2) - 1(x - 2) = (x - 2)(x - 1)$$

\therefore The possible length and breadth are $(x - 2)$ and $(x - 1)$.

(iii) We have,

$$\text{Area of rectangle} = x^2 - 7x + 10$$

$$= x^2 - (5 + 2)x + 10 = x^2 - 5x - 2x + 10$$

$$= x(x - 5) - 2(x - 5) = (x - 5)(x - 2)$$

\therefore The possible length and breadth are $(x - 5)$ and $(x - 2)$.

(iv) We have,

$$\text{Area of rectangle} = x^2 + 19x - 20$$

$$= x^2 + (20 - 1)x - 20 = x^2 + 20x - x - 20$$

$$= x(x + 20) - 1(x + 20) = (x + 20)(x - 1)$$

\therefore The possible length and breadth are $(x + 20)$ and $(x - 1)$.

(v) We have, area of rectangle

$$= x^2 + 9x + 20$$

$$= x^2 + (5 + 4)x + 20 = x^2 + 5x + 4x + 20$$

$$= x(x + 5) + 4(x + 5) = (x + 5)(x + 4)$$

\therefore The possible length and breadth are $(x + 5)$ and $(x + 4)$.

Question. 94 Carry out the following divisions:

- (i) $51x^3y^2z + 17xyz$ (ii) $76x^3yz^3 + 19x^2y^2$
 (iii) $17ab^2c^3 + (-abc^2)$ (iv) $-121p^3q^3r^3 + (-11xy^2z^3)$

Solution.

(i) We have,

$$\begin{aligned} 51x^3y^2z + 17xyz &= \frac{51x^3y^2z}{17xyz} \\ &= \frac{17 \times 3 \times x \times x \times x \times y \times y \times z}{17 \times x \times y \times z} = 3x^2y \end{aligned}$$

(ii) We have,

$$\begin{aligned} 76x^3yz^3 + 19x^2y^2 &= \frac{76x^3yz^3}{19x^2y^2} \\ &= \frac{4 \times 19 \times x \times x \times x \times y \times z \times z \times z}{19 \times x \times x \times y \times y} = \frac{4xz^3}{y} \end{aligned}$$

(iii) We have,

$$17ab^2c^3 + (-abc^2) = \frac{17ab^2c^3}{-abc^2} = \frac{17 \times a \times b \times b \times c \times c \times c}{-a \times b \times c \times c} = -17bc$$

(iv) We have,

$$\begin{aligned} -121p^3q^3r^3 + (-11xy^2z^3) &= \frac{-121p^3q^3r^3}{-11xy^2z^3} \\ &= \frac{-11 \times 11 \times p \times p \times p \times q \times q \times q \times r \times r \times r}{-11 \times x \times y \times y \times z \times z \times z} = \frac{11p^3q^3r^3}{xy^2z^3} \end{aligned}$$

Question. 95 Perform the following divisions:

- (i) $(3pqr - 6p^2q^2r^2) \div 3pq$
 (ii) $(ax^3 - bx^2 + cx) \div (-dx)$
 (iii) $(x^3y^3 + x^2y^3 - xy^4 + xy) \div xy$
 (iv) $(-qrx + pryz - rxyz) \div (-xyz)$

Solution.

(i) We have,

$$\begin{aligned} (3pqr - 6p^2q^2r^2) \div 3pq &= \frac{3pqr - 6p^2q^2r^2}{3pq} = \frac{3pqr}{3pq} - \frac{6p^2q^2r^2}{3pq} \\ &= r - \frac{2 \times 3 \times p \times p \times q \times q \times r \times r}{3 \times p \times q} \\ &= r - 2pqr^2 \end{aligned}$$

(ii) We have,

$$\begin{aligned} (ax^3 - bx^2 + cx) + (-dx) &= \frac{ax^3 - bx^2 + cx}{-dx} \\ &= \frac{ax^3}{-dx} + \frac{bx^2}{dx} + \frac{cx}{-dx} = \frac{a \times x \times x \times x}{-d \times x} + \frac{b \times x \times x}{d \times x} + \frac{c \times x}{-d \times x} \\ &= -\frac{a}{d}x^2 + \frac{b}{d}x - \frac{c}{d} \end{aligned}$$

(iii) We have,

$$\begin{aligned} (x^3y^3 + x^2y^3 - xy^4 + xy) + xy \\ &= \frac{x^3y^3 + x^2y^3 - xy^4 + xy}{xy} = \frac{x^3y^3}{xy} + \frac{x^2y^3}{xy} - \frac{xy^4}{xy} + \frac{xy}{xy} \\ &= \frac{x \times x \times x \times y \times y \times y}{x \times y} + \frac{x \times x \times y \times y \times y}{x \times y} - \frac{x \times y \times y \times y \times y}{x \times y} + \frac{x \times y}{x \times y} \\ &= x^2y^2 + xy^2 - y^3 + 1 \end{aligned}$$

(iv) We have,

$$\begin{aligned} (-qxy + pryz - rxyz) + (-xyz) \\ &= \frac{-qxy + pryz - rxyz}{-xyz} = \frac{-qxy}{-xyz} + \frac{pryz}{-xyz} - \frac{rxyz}{-xyz} = \frac{qr}{z} - \frac{pr}{x} + r \end{aligned}$$

Question. 96 Factorise the expressions and divide them as directed.

(i) $(x^2 - 22x + 117) \div (x - 13)$

(ii) $(x^3 + x^2 - 132x) \div x(x - 11)$

(iii) $(2x^3 - 12x^2 + 16x) \div (x - 2)(x - 4)$

(iv) $(9x^2 - 4) \div (3x + 2)$

(v) $(3x^2 - 48) \div (x - 4)$

(vi) $(x^4 - 16) \div (x^3 + 2x^2 + 4x + 8)$

(vii) $(3x^4 - 1875) \div (3x^2 - 75)$

Solution.

(i) We have,

$$\begin{aligned} (x^2 - 22x + 117) \div (x - 13) \\ &= \frac{x^2 - 22x + 117}{x - 13} = \frac{x^2 - 13x - 9x + 117}{x - 13} = \frac{x(x - 13) - 9(x - 13)}{x - 13} \\ &= \frac{(x - 13)(x - 9)}{x - 13} = x - 9 \end{aligned}$$

(ii) We have,

$$\begin{aligned} (x^3 + x^2 - 132x) \div x(x - 11) \\ &= \frac{x^3 + x^2 - 132x}{x(x - 11)} = \frac{x(x^2 + x - 132)}{x(x - 11)} = \frac{x^2 + 12x - 11x - 132}{x - 11} \\ &= \frac{x(x + 12) - 11(x + 12)}{x - 11} = \frac{(x + 12)(x - 11)}{x - 11} \\ &= x + 12 \end{aligned}$$

(iii) We have,

$$\begin{aligned}(2x^3 - 12x^2 + 16x) + (x-2)(x-4) &= \frac{2x^3 - 12x^2 + 16x}{(x-2)(x-4)} = \frac{2x(x^2 - 6x + 8)}{(x-2)(x-4)} \\ &= \frac{2x(x^2 - 4x - 2x + 8)}{(x-2)(x-4)} \\ &= \frac{2x[x(x-4) - 2(x-4)]}{(x-2)(x-4)} = \frac{2x(x-4)(x-2)}{(x-2)(x-4)} = 2x\end{aligned}$$

(iv) We have,

$$\begin{aligned}(9x^2 - 4) + (3x + 2) &= \frac{9x^2 - 4}{3x + 2} = \frac{(3x)^2 - (2)^2}{3x + 2} \\ &= \frac{(3x + 2)(3x - 2)}{3x + 2} \quad [\because a^2 - b^2 = (a + b)(a - b)] \\ &= 3x - 2\end{aligned}$$

(v) We have,

$$\begin{aligned}(3x^2 - 48) + (x - 4) &= \frac{3x^2 - 48}{x - 4} = \frac{3(x^2 - 16)}{x - 4} \\ &= \frac{3(x^2 - 4^2)}{x - 4} \\ &= \frac{3(x + 4)(x - 4)}{x - 4} \quad [\because a^2 - b^2 = (a + b)(a - b)] \\ &= 3(x + 4)\end{aligned}$$

(vi) We have,

$$\begin{aligned}(x^4 - 16) + x^3 + 2x^2 + 4x + 8 &= \frac{x^4 - 16}{x^3 + 2x^2 + 4x + 8} = \frac{(x^2)^2 - 4^2}{x^2(x+2) + 4(x+2)} \\ &= \frac{(x^2 + 4)(x^2 - 4)}{(x^2 + 4)(x + 2)} = \frac{x^2 - 2^2}{x + 2} = \frac{(x + 2)(x - 2)}{x + 2} = x - 2\end{aligned}$$

(vii) We have,

$$\begin{aligned}(3x^4 - 1875) + (3x^2 - 75) &= \frac{3x^4 - 1875}{3x^2 - 75} = \frac{x^4 - 625}{x^2 - 25} = \frac{(x^2)^2 - (25)^2}{x^2 - 25} \\ &= \frac{(x^2 + 25)(x^2 - 25)}{(x^2 - 25)} = x^2 + 25\end{aligned}$$

Question. 97 The area of a square is given by $4x^2 + 12xy + 9y^2$. Find the side of the square.

Solution.

We have,

$$\text{Area of square} = 4x^2 + 12xy + 9y^2$$

So, we factorise the given expression.

$$\begin{aligned}\therefore 4x^2 + 12xy + 9y^2 &= (2x)^2 + 2 \times 2x \times 3y + (3y)^2 \quad [\because a^2 + 2ab + b^2 = (a + b)^2] \\ &= (2x + 3y)^2\end{aligned}$$

Since, area of a square having side length a is a^2 . Hence, side of the given square is $2x + 3y$.

Question. 98 The area of a square is $9x^2 + 24xy + 16y^2$. Find the side of the square.

Solution.

We have,

$$\begin{aligned}\text{Area of a square} &= 9x^2 + 24xy + 16y^2 = (3x)^2 + 2 \times 3x \times 4y + (4y)^2 \\ & \qquad \qquad \qquad [\because a^2 + 2ab + b^2 = (a + b)^2] \\ &= (3x + 4y)^2\end{aligned}$$

$$\therefore \text{The side of the square} = 3x + 4y \qquad [\because \text{area of square} = (\text{side})^2]$$

Question. 99 The area of a rectangle is $x^2 + 7x + 12$. If its breadth is $(x + 3)$, then find its length.

Solution.

Let the length of the rectangle be l

Given, area of a rectangle = $x^2 + 7x + 12$

and breadth = $x + 3$

We know that,

Area of rectangle = Length \times Breadth

$$\Rightarrow x^2 + 7x + 12 = l \times (x + 3)$$

$$\Rightarrow l = \frac{x^2 + 7x + 12}{x + 3} = \frac{x^2 + 4x + 3x + 12}{x + 3} = \frac{x(x + 4) + 3(x + 4)}{x + 3} = \frac{(x + 4)(x + 3)}{x + 3} = x + 4$$

Hence, the length of rectangle = $x + 4$

Question. 100 The curved surface area of a cylinder is $2\pi(y^2 - 7y + 12)$ and its radius is $(y - 3)$. Find the height of the cylinder (CSA of cylinder = ~~Formula does not parse~~)

Solution.

Let the height of cylinder be h .

Given, the curved surface area of a cylinder = $2\pi(y^2 - 7y + 12)$

and radius of cylinder = $y - 3$

We know that,

Curved surface area of cylinder = $2\pi rh$

$$\therefore 2\pi rh = 2\pi(y^2 - 7y + 12)$$

$$\Rightarrow 2\pi rh = 2\pi(y^2 - 4y - 3y + 12) = 2\pi[y(y - 4) - 3(y - 4)] = 2\pi(y - 3)(y - 4)$$

$$\Rightarrow 2\pi h = 2\pi(y - 4) \qquad [\because r = (y - 3), \text{ given}]$$

On comparing the both sides, we get $h = y - 4$

Hence, the height of the cylinder is $y - 4$.

Question. 101 The area of a circle is given by the expression $\pi x^2 + 6\pi x + 9\pi$. Find the radius of the circle.

Solution.

We have,

Area of a circle = $\pi x^2 + 6\pi x + 9\pi = \pi(x^2 + 6x + 9)$

$$\Rightarrow \pi r^2 = \pi(x^2 + 3x + 3x + 9) \qquad [\because \text{area of a circle} = \pi r^2, \text{ where } r \text{ is the radius}]$$

$$\Rightarrow \pi r^2 = \pi[x(x + 3) + 3(x + 3)] = \pi(x + 3)(x + 3) = \pi(x + 3)^2$$

$$\Rightarrow \pi r^2 = \pi(x + 3)^2$$

$$\text{On comparing both sides, } r^2 = (x + 3)^2 \Rightarrow r = x + 3$$

Hence, the radius of circle is $x + 3$.

Question.102 The sum of first n natural numbers is given by the expression $\frac{n^2}{2} + \frac{n}{2}$ Factorise this expression.

Solution.

We have, the sum of first n natural numbers

$$= \frac{n^2}{2} + \frac{n}{2}$$

Factorisation of given expression $= \frac{1}{2}(n^2 + n) = \frac{1}{2}n(n + 1)$ [taking n as common]

Question.103 The sum of $(x + 5)$ observations is $x^4 - 625$. Find the mean of the observations.

Solution.

We have, the sum of $(x + 5)$ observations $= x^4 - 625$

We know that, the mean of the n observations x_1, x_2, \dots, x_n is given by $\frac{x_1 + x_2 + \dots + x_n}{n}$.

\therefore The mean of $(x + 5)$ observations

$$\begin{aligned} &= \frac{\text{Sum of } (x + 5) \text{ observations}}{x + 5} = \frac{x^4 - 625}{x + 5} = \frac{(x^2)^2 - (25)^2}{x + 5} \\ &= \frac{(x^2 + 25)(x^2 - 25)}{x + 5} \quad [\because a^2 - b^2 = (a + b)(a - b)] \\ &= \frac{(x^2 + 25)[(x)^2 - (5)^2]}{x + 5} \\ &= \frac{(x^2 + 25)(x + 5)(x - 5)}{(x + 5)} = (x^2 + 25)(x - 5) \end{aligned}$$

Question.104 The height of a triangle is $x^4 + y^4$ and its base is $14xy$. Find the area of the triangle.

Solution.

Given, the height of a triangle and its base are $x^4 + y^4$ and $14xy$, respectively.

We know that, the area of a triangle $= \frac{1}{2} \times \text{Base} \times \text{Height} = \frac{1}{2} \times 14xy \times (x^4 + y^4)$

$$= 7xy(x^4 + y^4)$$

Question.105 The cost of a chocolate is Rs $(x + 4)$ and Rohit bought $(x + 4)$ chocolates. Find the total amount paid by him in terms of x . If $x = 10$, find the amount paid by him.

Solution.

Given, cost of a chocolate $= ₹ (x + 4)$

Rohit bought $(x + 4)$ chocolates.

\therefore The cost of $(x + 4)$ chocolates

$$\begin{aligned} &= \text{Cost of one chocolate} \times \text{Number of chocolates} = (x + 4)(x + 4) = (x + 4)^2 \\ &= x^2 + 8x + 16 \quad [\because (a + b)^2 = a^2 + 2ab + b^2] \end{aligned}$$

\therefore The total amount paid by Rohit $= ₹(x^2 + 8x + 16)$

Now, if $x = 10$. Then, the amount paid by Rohit $= 10^2 + 8 \times 10 + 16 = 100 + 80 + 16 = ₹ 196$

Question.106 The base of a parallelogram is $(2x + 3)$ units and the corresponding height is $(2x - 3)$ units. Find the area of the parallelogram in terms of x . What will be the area of a parallelogram of $x = 30$ units?

Solution.

We have, the base and the corresponding height of a parallelogram are $(2x + 3)$ units and $(2x - 3)$ units, respectively.

\therefore Area of a parallelogram $= \text{Base} \times \text{Height}$

$$\begin{aligned} &= (2x + 3) \times (2x - 3) = (2x)^2 - (3)^2 \quad [\because (a + b)(a - b) = a^2 - b^2] \\ &= (4x^2 - 9) \text{ sq units} \end{aligned}$$

Now, if $x = 30$. Then, the area of the parallelogram $= 4 \times (30)^2 - 9 = 400 - 9 = 391$ sq units

Question.107 The radius of a circle is $7ab - 7bc - 14ac$. Find the circumference of the circle, $(\pi = \frac{22}{7})$

Solution.

We have, radius of the circle = $7ab - 7bc - 14ac = r$

[say]

We know that,

\therefore The circumference of the circle = $2\pi r$

$$= 2 \times \frac{22}{7} \times (7ab - 7bc - 14ac)$$

$$= \frac{44}{7} \times 7(ab - bc - 2ac)$$

$$= 44[ab - c(b + 2a)]$$

Question.108 If $p + q = 12$ and $pq = 22$, then find $p^2 + q^2$.

Solution.

Given, $p + q = 12$ and $pq = 22$

Since,

$$(p + q)^2 = p^2 + q^2 + 2pq \quad [\text{using the identity, } (a + b)^2 = a^2 + b^2 + 2ab]$$

$$\therefore (12)^2 = p^2 + q^2 + 2 \times 22$$

$$\Rightarrow p^2 + q^2 = (12)^2 - 44$$

$$\Rightarrow p^2 + q^2 = 144 - 44 = 100$$

Question.109 If $a + b = 25$ and $a^2 + b^2$ then find ab .

Solution.

Given, $a + b = 25$ and $a^2 + b^2 = 225$

We know that,

$$(a + b)^2 = a^2 + b^2 + 2ab$$

[an algebraic identity]

$$\Rightarrow (25)^2 = 225 + 2ab$$

$$\Rightarrow 2ab = (25)^2 - 225$$

$$\Rightarrow 2ab = 625 - 225$$

$$\Rightarrow 2ab = 400$$

$$\Rightarrow ab = \frac{400}{2}$$

$$\Rightarrow ab = 200$$

Question.110 If $x - y = 13$ and $xy = 28$, then find $x^2 + y^2$.

Solution.

Given, $x - y = 13$ and $xy = 28$

Since,

$$(x - y)^2 = x^2 + y^2 - 2xy$$

[using the identity, $(a - b)^2 = a^2 + b^2 - 2ab$]

$$\therefore (13)^2 = x^2 + y^2 - 2 \times 28$$

$$\Rightarrow x^2 + y^2 = (13)^2 + 56$$

$$\Rightarrow x^2 + y^2 = 169 + 56$$

$$\Rightarrow x^2 + y^2 = 225$$

Question.111 If $m - n = 16$ and $m^2 + n^2 = 400$, then find mn .

Solution.

Given, $m - n = 16$ and $m^2 + n^2 = 400$.

Since, $(m - n)^2 = m^2 + n^2 - 2mn$
[using the identity, $(a - b)^2 = a^2 + b^2 - 2ab$]

$$\begin{aligned} \therefore (16)^2 &= 400 - 2mn \\ \Rightarrow 2mn &= 400 - (16)^2 \\ \Rightarrow 2mn &= 400 - 256 \\ \Rightarrow 2mn &= 144 \\ \Rightarrow mn &= \frac{144}{2} \\ \Rightarrow mn &= 72 \end{aligned}$$

Question.112 If $a^2 + b^2 = 74$ and $ab = 35$, then find $a + b$?

Solution.

Given, $a^2 + b^2 = 74$ and $ab = 35$

Since, $(a + b)^2 = a^2 + b^2 + 2ab$
[using the identity, $(a + b)^2 = a^2 + b^2 + 2ab$]

$$\begin{aligned} \therefore (a + b)^2 &= 74 + 2 \times 35 \\ \therefore (a + b)^2 &= 74 + 2 \times 35 \\ \Rightarrow (a + b)^2 &= 144 \\ \Rightarrow a + b &= \sqrt{144} && \text{[taking square root]} \\ \Rightarrow a + b &= 12 && \text{[rejecting -ve sign]} \end{aligned}$$

Question.113 Verify the following:

- (i) $(ab + bc)(ab - bc) + (bc + ca)(bc - ca) + (ca + ab)(ca - ab) = 0$
- (ii) $(a + b + c)(a^2 + b^2 + c^2 - ab - bc - ca) = a^3 + b^3 + c^3 - 3abc$
- (iii) $(p - q)(p^2 + pq + q^2) = p^3 - q^3$
- (iv) $(m + n)(m^2 - mn + n^2) = m^3 + n^3$
- (v) $(a + b)(a + b)(a + b) = a^3 + 3a^2b + 3ab^2 + b^3$
- (vi) $(a - b)(a - b)(a - b) = a^3 - 3a^2b + 3ab^2 - b^3$
- (vii) $(a^2 - b^2)(a^2 + b^2) + (b^2 - c^2)(b^2 + c^2) + (c^2 - a^2)(c^2 + a^2) = 0$
- (viii) $(5x + 8)^2 - 160x = (5x - 8)^2$
- (ix) $(7p - 13q)^2 + 364pq = (7p + 13q)^2$
- (x) $\left(\frac{3p}{7} + \frac{7}{6p}\right)^2 - \left(\frac{3p}{7} - \frac{7}{6p}\right)^2 = 2$

Solution.

$$\begin{aligned}
 \text{(i) Taking LHS} &= (ab + bc)(ab - bc) + (bc + ca)(bc - ca) + (ca + ab)(ca - ab) \\
 &= [(ab)^2 - (bc)^2] + [(bc)^2 - (ca)^2] + [(ca)^2 - (ab)^2] \\
 &\quad \text{[using the identity, } (a + b)(a - b) = a^2 - b^2\text{]} \\
 &= a^2b^2 - b^2c^2 + b^2c^2 - c^2a^2 + c^2a^2 - a^2b^2 = 0 \\
 &= \text{RHS} \quad \text{[cancelling the like terms having opposite signs]} \\
 &\quad \text{Hence verified.}
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) Taking LHS} &= (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ca) \\
 &= a(a^2 + b^2 + c^2 - ab - bc - ca) + b(a^2 + b^2 + c^2 - ab - bc - ca) \\
 &\quad + c(a^2 + b^2 + c^2 - ab - bc - ca) \\
 &\quad \text{[distributive law]} \\
 &= a^3 + ab^2 + ac^2 - a^2b - abc - a^2c + ba^2 + b^3 + bc^2 \\
 &\quad - b^2a - b^2c - bca + ca^2 + cb^2 + c^3 - cab - c^2b - c^2a \\
 &= a^3 + b^3 + c^3 - 3abc = \text{RHS} \\
 &\quad \text{Hence verified.}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) Taking LHS} &= (p - q)(p^2 + pq + q^2) \\
 &= p(p^2 + pq + q^2) - q(p^2 + pq + q^2) \\
 &= p^3 + p^2q + pq^2 - qp^2 - pq^2 - q^3 = p^3 - q^3 = \text{RHS} \\
 &\quad \text{Hence verified.}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv) Taking LHS} &= (m + n)(m^2 - mn + n^2) \\
 &= m(m^2 - mn + n^2) + n(m^2 - mn + n^2) \\
 &= m^3 - m^2n + mn^2 + nm^2 - mn^2 + n^3 = m^3 + n^3 = \text{RHS} \\
 &\quad \text{Hence verified.}
 \end{aligned}$$

$$\begin{aligned}
 \text{(v) Taking LHS} &= (a + b)(a + b)(a + b) \\
 &= (a + b)(a + b)^2 \\
 &= (a + b)(a^2 + b^2 + 2ab) \quad \text{[using the identity, } (a + b)^2 = a^2 + 2ab + b^2\text{]} \\
 &= a(a^2 + 2ab + b^2) + b(a^2 + 2ab + b^2) \\
 &= a^3 + 2a^2b + ab^2 + ba^2 + 2ab^2 + b^3 \\
 &= a^3 + 3a^2b + 3ab^2 + b^3 \quad \text{[adding like terms]} \\
 &= \text{RHS} \\
 &\quad \text{Hence verified.}
 \end{aligned}$$

$$\begin{aligned}
 \text{(vi) Taking LHS} &= (a - b)(a - b)(a - b) \\
 &= (a - b)(a - b)^2 \\
 &= (a - b)(a^2 - 2ab + b^2) \quad \text{[using the identity, } (a - b)^2 = a^2 - 2ab + b^2\text{]} \\
 &= a(a^2 - 2ab + b^2) - b(a^2 - 2ab + b^2) \\
 &= a^3 - 2a^2b + ab^2 - ba^2 + 2ab^2 - b^3 \\
 &= a^3 - 3a^2b + 3ab^2 - b^3 \quad \text{[adding like terms]} \\
 &= \text{RHS} \\
 &\quad \text{Hence verified.}
 \end{aligned}$$

$$\begin{aligned}
 \text{(vii) Taking LHS} &= (a^2 - b^2)(a^2 + b^2) + (b^2 - c^2)(b^2 + c^2) + (c^2 - a^2)(c^2 + a^2) \\
 &= (a^4 - b^4 + b^4 - c^4 + c^4 - a^4) \quad \text{[using the identity, } (a - b)(a + b) = a^2 - b^2\text{]} \\
 &= 0 = \text{RHS} \\
 &\quad \text{Hence verified.}
 \end{aligned}$$

(viii) Taking LHS = $(5x + 8)^2 - 160x$

$$= (5x)^2 + (8)^2 + 2 \times 5x \times 8 - 160x \quad [\text{using the identity, } (a + b)^2 = a^2 + 2ab + b^2]$$

$$= (5x)^2 + (8)^2 + 80x - 160x$$

$$= (5x)^2 + (8)^2 - 80x$$

$$= (5x)^2 + (8)^2 - 2 \times 5x \times 8$$

$$= (5x - 8)^2 \quad [\because a^2 + b^2 - 2ab = (a - b)^2]$$

$$= \text{RHS}$$

Hence verified.

(ix) Taking LHS = $(7p - 13q)^2 + 364pq$

$$= (7p)^2 + (13q)^2 - 2 \times 7p \times 13q + 364pq$$

$$= (7p)^2 + (13q)^2 - 182pq + 364pq$$

$$= (7p)^2 + (13q)^2 + 182pq$$

$$= (7p)^2 + (13q)^2 + 2 \times 7p \times 13q = (7p + 13q)^2 = \text{RHS}$$

Hence verified.

(x) Taking LHS = $\left(\frac{3p}{7} + \frac{7}{6p}\right)^2 - \left(\frac{3p}{7} - \frac{7}{6p}\right)^2$

$$= \left[\left(\frac{3p}{7} + \frac{7}{6p}\right) + \left(\frac{3p}{7} - \frac{7}{6p}\right) \right] \left[\left(\frac{3p}{7} + \frac{7}{6p}\right) - \left(\frac{3p}{7} - \frac{7}{6p}\right) \right]$$

[using the identity, $a^2 - b^2 = (a + b)(a - b)$]

$$= \left(\frac{3p}{7} + \frac{7}{6p} + \frac{3p}{7} - \frac{7}{6p}\right) \left(\frac{3p}{7} + \frac{7}{6p} - \frac{3p}{7} + \frac{7}{6p}\right) = \frac{6p}{7} \times \frac{14}{6p} = 2 = \text{RHS}$$

Hence verified.

Question.114 Find the value of a, if

(i) $8a = 35^2 - 27^2$

(ii) $9a = 76^2 - 67^2$

(iii) $pqa = (3p + q)^2 - (3p - q)^2$

(iv) $pq^2a = (4pq + 3q)^2 - (4pq - 3q)^2$

Solution.

(i) We have,

$$8a = 35^2 - 27^2$$

$$\Rightarrow 8a = (35 + 27)(35 - 27) \quad [\text{using the identity, } a^2 - b^2 = (a + b)(a - b)]$$

$$\Rightarrow 8a = 62 \times 8$$

$$\Rightarrow a = \frac{62 \times 8}{8}$$

$$\Rightarrow a = 62$$

(ii) We have,

$$9a = (76)^2 - (67)^2$$

$$\Rightarrow 9a = (76 + 67)(76 - 67) \quad [\text{using the identity, } a^2 - b^2 = (a + b)(a - b)]$$

$$\Rightarrow 9a = 143 \times 9$$

$$\Rightarrow a = \frac{143 \times 9}{9}$$

$$\Rightarrow a = 143$$

(iii) We have,

$$pqa = (3p + q)^2 - (3p - q)^2$$

$$\Rightarrow pqa = [(3p + q) + (3p - q)][(3p + q) - (3p - q)]$$

[using the identity, $a^2 - b^2 = (a + b)(a - b)$]

$$\Rightarrow pqa = [(3p + q + 3p - q)][3p + q - 3p + q]$$

$$\Rightarrow pqa = 6p \times 2q$$

$$\Rightarrow a = \frac{6p \times 2q}{pq} = \frac{(6 \times 2)pq}{pq}$$

$$\Rightarrow a = 12$$

(iv) We have,

$$\begin{aligned}pq^2a &= (4pq + 3q)^2 - (4pq - 3q)^2 \\ \Rightarrow &= [(4pq + 3q) + (4pq - 3q)][(4pq + 3q) - (4pq - 3q)] \\ &\quad \text{[using the identity, } a^2 - b^2 = (a + b)(a - b)\text{]} \\ &= (4pq + 3q + 4pq - 3q)(4pq + 3q - 4pq + 3q) \\ &= 8pq \times 6q \\ \Rightarrow & pq^2a = 48pq^2 \\ \Rightarrow & a = \frac{48pq^2}{pq^2} \\ \Rightarrow & a = 48\end{aligned}$$

Question.115 What should be added to $4c(-a + b + c)$ to obtain $3a(a + b + c) - 2b(a - b + c)$?

Solution.

Let x be added to the given expression

$4c(-a + b + c)$ to obtain $3a(a + b + c) - 2b(a - b + c)$

i.e. $x + 4c(-a + b + c) = 3a(a + b + c) - 2b(a - b + c)$

$$\begin{aligned}\Rightarrow & x = 3a(a + b + c) - 2b(a - b + c) - 4c(-a + b + c) \\ &= 3a^2 + 3ab + 3ac - 2ba + 2b^2 - 2bc + 4ca - 4cb - 4c^2\end{aligned}$$

$$\Rightarrow x = 3a^2 + ab + 7ac + 2b^2 - 6bc - 4c^2 \quad \text{[adding the like terms]}$$

Question.116 Subtract $b(b^2 + b - 7) + 5$ from $3b^2 - 8$ and find the value of expression obtained for $b = -3$.

Solution.

We have,

$$\begin{aligned}\text{Required difference} &= (3b^2 - 8) - [b(b^2 + b - 7) + 5] \\ &= 3b^2 - 8 - b(b^2 + b - 7) - 5 \\ &= 3b^2 - 8 - b^3 - b^2 + 7b - 5 = -b^3 + 2b^2 + 7b - 13\end{aligned}$$

Now, if $b = -3$

$$\begin{aligned}\text{The value of above expression} &= -(-3)^3 + 2(-3)^2 + 7(-3) - 13 \\ &= -(-27) + 2 \times 9 - 21 - 13 \\ &= 27 + 18 - 21 - 13 \\ &= 45 - 34 = 11\end{aligned}$$

Question.117 If $x - \frac{1}{x} = 7$, then find the value of $x^2 + \frac{1}{x^2}$.

Solution.

Given, $x - \frac{1}{x} = 7$

Since, $\left(x - \frac{1}{x}\right)^2 = x^2 + \frac{1}{x^2} - 2 \cdot x \cdot \frac{1}{x}$ [using the identity, $(a - b)^2 = a^2 + b^2 - 2ab$]

$$\therefore 7^2 = x^2 + \frac{1}{x^2} - 2$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 49 + 2$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 51$$

Question.118 Factorise $x^2 + \frac{1}{x^2} + 2 - 3x - \frac{1}{x}$.

Solution.

We have, $x^2 + \frac{1}{x^2} + 2 - 3x - \frac{3}{x}$

$$= x^2 + \frac{1}{x^2} + 2 \cdot x \cdot \frac{1}{x} - 3 \left(x + \frac{1}{x} \right)$$

$$= \left(x + \frac{1}{x} \right)^2 - 3 \left(x + \frac{1}{x} \right)$$

[using the identity, $a^2 + b^2 + 2ab = (a + b)^2$]

$$= \left(x + \frac{1}{x} \right) \left(x + \frac{1}{x} - 3 \right) \quad \left[\text{taking } \left(x + \frac{1}{x} \right) \text{ as common} \right]$$

Question.119 Factorise $p^4 + q^4 + p^2q^2$.

Solution.

We have, $p^4 + q^4 + p^2q^2$

$$= p^4 + q^4 + 2p^2q^2 - 2p^2q^2 + p^2q^2 \quad [\text{adding and subtracting } 2p^2q^2]$$

$$= p^4 + q^4 + 2p^2q^2 - p^2q^2$$

$$= [(p^2)^2 + (q^2)^2 + 2p^2q^2] - p^2q^2$$

[using the identity, $a^2 + b^2 + 2ab = (a + b)^2$]

$$= (p^2 + q^2)^2 - (pq)^2$$

$$= (p^2 + q^2 + pq)(p^2 + q^2 - pq) \quad [\text{using the identity, } a^2 - b^2 = (a + b)(a - b)]$$

Question.120 Find the value of

(i) $\frac{6.25 \times 6.25 - 1.75 \times 1.75}{4.5}$ (ii) $\frac{198 \times 198 - 102 \times 102}{96}$

Solution.

(i) We have, $\frac{6.25 \times 6.25 - 1.75 \times 1.75}{4.5} = \frac{(6.25)^2 - (1.75)^2}{4.5}$

$$= \frac{(6.25 + 1.75)(6.25 - 1.75)}{4.5} \quad [\text{using the identity, } a^2 - b^2 = (a + b)(a - b)]$$

$$= \frac{8 \times 4.5}{4.5} = 8$$

(ii) We have,

$$\frac{198 \times 198 - 102 \times 102}{96} = \frac{(198)^2 - (102)^2}{96} = \frac{(198 + 102)(198 - 102)}{96}$$

$$= \frac{300 \times 96}{96} = 300 \quad [\text{using the identity, } a^2 - b^2 = (a + b)(a - b)]$$

Question.121 The product of two expressions is $x^5 + x^3 + x$. If one of them is $x^2 + x + 1$, find the other.

Solution.

We have, product of two expressions $x^5 + x^3 + x$ and one is $x^2 + x + 1$.

Let the other expression be A. Then,

$$A \cdot (x^2 + x + 1) = x^5 + x^3 + x$$

$$\Rightarrow A = \frac{x^5 + x^3 + x}{x^2 + x + 1} = \frac{x(x^4 + x^2 + 1)}{x^2 + x + 1}$$

$$\Rightarrow A = \frac{x(x^4 + 2x^2 - x^2 + 1)}{x^2 + x + 1} = \frac{x(x^4 - 2x^2 + 1 - x^2)}{x^2 + x + 1}$$

[adding and subtracting x^2 in numerator term]


$$= \frac{x[(x^4 + 2x^2 + 1) - x^2]}{x^2 + x + 1} = \frac{x[(x^2 + 1)^2 - x^2]}{x^2 + x + 1}$$

$$= \frac{x(x^2 + 1 + x)(x^2 + 1 - x)}{x^2 + x + 1} \quad [\text{using the identity, } a^2 - b^2 = (a + b)(a - b)]$$

$$= x(x^2 + 1 - x)$$

Hence, the other expression is $x(x^2 - x + 1)$.

Question.122 Find the length of the side of the given square, if area of the square is 625sq units and then find the value of x.



Solution.

We have, a square having length of a side $(4x + 5)$ units and area is 625 sq units.

$$\therefore \text{Area of a square} = (\text{Side})^2$$

$$(4x + 5)^2 = 625$$

$$\Rightarrow (4x + 5)^2 = (25)^2 \quad [\text{taking square root both sides and neglecting } (-ve) \text{ sign}]$$

$$\Rightarrow 4x + 5 = 25$$


$$\Rightarrow 4x = 25 - 5$$

$$\Rightarrow 4x = 20$$

$$\Rightarrow x = 5$$

Hence, side = $4x + 5 = 4 \times 5 + 5 = 25$ units

Question.123 Take suitable number of cards given in the adjoining diagram [$G(x \times x)$ representing x^2 , $R(x \times 1)$ representing x and $Y(1 \times 1)$ representing 1] to factorise the following expressions, by arranging to cards in the form of rectangles: (i) $2x^2 + 6x + 4$ (ii) $x^2 + 4x + 4$. Factorise $2x^2 + 6x + 4$ by using the figure.




Calculate the area of figure.

Solution. The given information is incomplete for solution of this question.

Question.124 The figure shows the dimensions of a wall having a window and a door of a

room. Write an algebraic expression for the area of the wall to be painted.



Solution.

We have a wall of dimension $5x \times (5x + 2)$ having a window and a door of dimension $(2x \times x)$ and $(3x \times x)$, respectively.

Then, area of the window = $2x \times x = 2x^2$ sq units

Area of the door = $3x \times x = 3x^2$ sq units

and area of wall = $(5x + 2) \times 5x = (25x^2 + 10x)$ sq units

Now, area of the required part of the wall to be painted

= Area of the wall - (Area of the window + Area of the door)

= $25x^2 + 10x - (2x^2 + 3x^2)$

= $25x^2 + 10x - 5x^2 = 20x^2 + 10x$

= $2 \times 2 \times 5 \times x \times x + 2 \times 5 \times x$

= $2 \times 5 \times x(2x + 1) = 10x(2x + 1)$ sq units

Question.125 Match the expressions of column I with that of column II

Column I	Column II
(i) $(21x + 13y)^2$	(a) $441x^2 - 169y^2$
(ii) $(21x - 13y)^2$	(b) $441x^2 + 169y^2 + 546xy$
(iii) $(21x - 13y)(21x + 13y)$	(c) $441x^2 + 169y^2 - 546xy$
	(d) $441x^2 - 169y^2 + 546xy$

Solution.

(i) We have,

$$(21x + 13y)^2 = (21x)^2 + (13y)^2 + 2 \times 21x \times 13y$$

[using the identity, $(a + b)^2 = a^2 + b^2 + 2ab$]

$$= 441x^2 + 169y^2 + 546xy$$

(ii) $(21x - 13y)^2 = (21x)^2 + (13y)^2 - 2 \times 21x \times 13y$

[using the identity, $(a - b)^2 = a^2 + b^2 - 2ab$]

$$= 441x^2 + 169y^2 - 546xy$$

(iii) $(21x - 13y)(21x + 13y)$

$$= (21x)^2 - (13y)^2 = 441x^2 - 169y^2$$

[using the identity, $(a - b)(a + b) = a^2 - b^2$]

Hence, (i) \rightarrow (b), (ii) \rightarrow (c), (iii) \rightarrow (d)